Periodical
QD
271
.J6
Index
1980

ISSN 0021-9673

BLISHED
EKLY

CUMULATIVE AUTHOR AND SUBJECT INDEXES VOLUMES 181–200

Supplied free of charge to the 1980 subscribers of the Journal of Chromatography

RNAL OF

CHROMATOGRAPHY

ERNATIONAL JOURNAL ON CHROMATOGRAPHY, ELECTROPHORESIS AND RELATED METHODS

RSITY OF HAWAII
LIBRARY
2 0 PM '81

EDITOR, Michael Lederer (Switzerland)
ASSOCIATE EDITOR, K. Macek (Prague)
EDITORIAL BOARD
W. A. Aue (Halifax)
V. G. Berezkin (Moscow)
V. Betina (Bratislava)
A. Bevenue (Honolulu, HI)
P. Boulanger (Lille)
A. A. Boulton (Saskatoon)
G. P. Cartoni (Rome)
G. Duyckaerts (Liège)
L. Fishbein (Jefferson, AR)
A. Frigerio (Milan)
C. W. Gehrke (Columbia, MO)
E. Gil-Av (Rehovot)
G. Guiochon (Palaiseau)
I. M. Hais (Hradec Králové)
J. K. Haken (Kensington)
E. Heftmann (Berkeley, CA)
S. Hjertén (Uppsala)
E. C. Horning (Houston, TX)
Cs. Horváth (New Haven, CT)
J. F. K. Huber (Vienna)
A. T. James (Sharnbrook)
J. Janák (Brno)
E.sz. Kováts (Lausanne)
K. A. Kraus (Oak Ridge, TN)
E. Lederer (Gif-sur-Yvette)
A. Liberti (Rome)
H. M. McNair (Blacksburg, VA)
Y. Marcus (Jerusalem)
G. B. Marini-Bettòlo (Rome)
Č. Michalec (Prague)
R. Neher (Basel)
G. Nickless (Bristol)
J. Novák (Brno)
N. A. Parris (Wilmington, DE)
P. G. Righetti (Milan)
O. Samuelson (Göteborg)
G.-M. Schwab (Munich)
G. Semenza (Zürich)
L. R. Snyder (Tarrytown, NY)
A. Zlatkis (Houston, TX)
EDITORS, BIBLIOGRAPHY SECTION
K. Macek (Prague), J. Janák (Brno), Z. Deyl (Prague)
COORD. EDITOR, DATA SECTION
J. Gasparič (Hradec Králové)

ELSEVIER SCIENTIFIC PUBLISHING COMPANY
AMSTERDAM

ERIODICAL
QD
271
J6
Index
1980
.H. LIBRARY

JOURNAL OF CHROMATOGRAPHY

Scope. The *Journal of Chromatography* publishes papers on all aspects of chromatography, electrophoresis and related methods. Contributions consist mainly of research papers dealing with chromatographic theory, instrumental development and their applications. The section *Biomedical Applications*, which is under separate editorship, deals with the following aspects: developments in and applications of chromatographic and electrophoretic techniques related to clinical diagnosis (including the publication of normal values); screening and profiling procedures with special reference to metabolic disorders; results from basic medical research with direct consequences in clinical practice; combinations of chromatographic and electrophoretic methods with other physicochemical techniques such as mass spectrometry. In *Chromatographic Reviews*, reviews on all aspects of chromatography, electrophoresis and related methods are published.

Submission of Papers. Papers in English, French and German may be submitted, if possible in three copies. Manuscripts should be submitted to: The Editor of *Journal of Chromatography*, P.O. Box 681, 1000 AR Amsterdam, The Netherlands, or to: The Editor of *Journal of Chromatography, Biomedical Applications*, P.O. Box 681, 1000 AR Amsterdam, The Netherlands. Reviews are invited or proposed by letter to the Editors and will appear in *Chromatographic Reviews* or *Biomedical Applications*. An outline of the proposed review should first be forwarded to the Editors for preliminary discussion prior to preparation. Submission of an article is understood to imply that the article is original and unpublished and is not being considered for publication elsewhere. For copyright regulations, see below.

Subscription Orders. Subscription orders should be sent to: Elsevier Scientific Publishing Company, P.O. Box 211, 1000 AE Amsterdam, The Netherlands. The *Journal of Chromatography* and the *Biomedical Applications* section can be subscribed to separately.

Publication. The *Journal of Chromatography* (incl. *Biomedical Applications, Chromatographic Reviews* and *Cumulative Author and Subject Indexes, Vols. 201–210* and *211–220*) has 24 volumes in 1981. The subscription prices for 1981 are:

J. Chromatogr. (incl. *Chromatogr. Rev.* and *Cum. Indexes*) + *Biomed. Appl.* (Vols. 203–226):
Dfl. 3240.00 plus Dfl. 432.00 (postage) (total *ca.* US$ 1883.00)

J. Chromatogr. (incl. *Chromatogr. Rev.* and *Cum. Indexes*) only (Vols. 203–220):
Dfl. 2556.00 plus Dfl. 324.00 (postage) (total *ca.* US$ 1477.00)

Biomed. Appl. only (Vols. 221–226):
Dfl. 852.00 plus Dfl. 108.00 (postage) (total *ca.* US$ 492.25).

Journals are automatically sent by air mail to the U.S.A. and Canada at no extra costs, and to Japan, Australia and New Zealand with a small additional postal charge. Back volumes of the *Journal of Chromatography* (Vols. 1 through 202) are available at Dfl. 145.00 (plus postage). Claims for issues not received should be made within three months of publication of the issue. If not, they cannot be honoured free of charge. For customers in the U.S.A. and Canada wishing additional bibliographic information on this and other Elsevier journals, please contact Elsevier/North-Holland Inc., Journal Information Centre, 52 Vanderbilt Avenue, New York, NY 10164. Tel: (212) 867-9040.

Abstracts/Contents Lists published in Biochemical Abstracts, Biological Abstracts, Chemical Abstracts, Chemical Titles, Current Contents/Physical, Chemical & Earth Sciences, Current Contents/Life Sciences, Index Medicus, and Science Citation Index.

See page 3 of cover for Publication Schedule, Information for Authors, and information on the News Section and Advertisements.

© ELSEVIER SCIENTIFIC PUBLISHING COMPANY—1981

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system or transmitted in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, without the prior written permission of the publisher, Elsevier Scientific Publishing Company, P.O. Box 330, 1000 AH Amsterdam, The Netherlands.

Submission of an article for publication implies the transfer of the copyright from the author(s) to the publisher and entails the authors' irrevocable and exclusive authorization of the publisher to collect any sums or considerations for copying or reproduction payable by third parties (as mentioned in article 17 paragraph 2 of the Dutch Copyright Act of 1912 and in the Royal Decree of June 20, 1974 (S. 351) pursuant to article 16 b of the Dutch Copyright Act of 1912) and/or to act in or out of Court in connection therewith.

Special regulations for readers in the U.S.A. This journal has been registered with the Copyright Clearance Center, Inc. Consent is given for copying of articles for personal or internal use, or for the personal use of specific clients. This consent is given on the condition that the copier pays through the Center the per-copy fee stated in the code on the first page of each article for copying beyond that permitted by Sections 107 or 108 of the U.S. Copyright Law. The appropriate fee should be forwarded with a copy of the first page of the article to the Copyright Clearance Center, Inc., 21 Congress Street, Salem, MA 01970, U.S.A. If no code appears in an article, the author has not given broad consent to copy and permission to copy must be obtained directly from the author. All articles published prior to 1980 may be copied for a per-copy fee of US$ 2.25, also payable through the Center. This consent does not extend to other kinds of copying, such as for general distribution, resale, advertising and promotion purposes, or for creating new collective works. Special written permission must be obtained from the publisher for such copying.

Special regulations for authors in the U.S.A. Upon acceptance of an article by the journal, the author(s) will be asked to transfer copyright of the article to the publisher. This transfer will ensure the widest possible dissemination of information under the U.S. Copyright Law.

Printed in The Netherlands

Introducing Chromosorb LC-9. The specialist for carbohydrate analysis.

Typical applications: carbohydrates, high fructose syrup, sugars, amines, nucleotides, strongly polar compounds, nitro compounds, aromatic compounds, carboxylic acids, and polyhydroxy compounds.

New Chromosorb LC-9 joins the long line of fine quality column packings for liquid chromatography developed by Johns-Manville.

LC-9 is a microparticulate polar-bonded HPLC column packing prepared by chemically bonding an amino-type functional group on a Chromosorb LC-6 silica surface. 10μ. Typical physical properties: totally porous, irregularly shaped, high purity silica with high polarity, surface area of 350+ m^2g, average pore size of 110 angstroms and a bulk density (packed column) of 0.48 g/cc.

There are also eight other grades of Chromosorb LC supports and packings to choose from:

LC-1 An inert diatomite-based support compatible with all liquid phases, 37-44μ.

LC-2 A pellicular packing having active porous silica layers permanently bonded to an inert spherical solid glass core, 37-44μ.

LC-3 A pellicular packing with alumina bonded to the surface of diatomite, 37-44μ.

LC-4 Pellicular reverse phase packing with a monomolecular layer of octadecyltrichlorosilane chemically bonded to Chromosorb LC-2, 37-44μ.

LC-5 Pellicular reverse phase packing with a monomolecular layer of diphenyldichlorosilane chemically bonded to Chromosorb LC-2, 37-44μ.

LC-6 A fully porous microparticulate silica, 5μ and 10μ.

LC-7 A reverse phase microparticulate HPLC column packing material prepared by chemically bonding ODS to Chromosorb LC-6 via Si-O-Si-C type bond, 5μ and 10μ.

LC-8 A microparticulate polar-bonded phase HPLC column packing material prepared by chemically bonding a cyano-type group to a Chromosorb LC-6 silica surface, 5μ and 10μ.

For more information on specific performance and packing of Johns-Manville products or a handy guide to all 33 Johns-Manville chromatography products, contact Jack Denboske, Filtration and Minerals Division, Johns-Manville, Ken-Caryl Ranch, Denver, CO 80217, (303) 979-1000. Telex: 454404 JOHNMANVILLE DVR

JM Johns-Manville

TrAC

trends in analytical chemistry

A monthly professional magazine publishing short, critical reviews and news on trends and developments in analytical chemistry

Analytical chemistry has grown so rapidly and is applied to so many diverse problems that it is impossible for analytical chemists to have specialist knowledge of every available technique. However, it is important for them to be aware of techniques outside their own area and of what these techniques can achieve. They must also be in a position to select appropriate methods for solving the broad spectrum of problems encountered in practice. **Trends in Analytical Chemistry** will provide this information - its purpose is to promote communications about methodology amongst all scientists involved in chemical analysis.

Trends in Analytical Chemistry will

- publish short critical reviews that are written in such a way as to be intelligible to an interdisciplinary audience
- publish regular feature articles
- publish insights into norms, procedures and standards, related to the function, organization and operation of industrial, government or research laboratories
- serve as an invaluable teaching aid — it will be more up-to-date than any textbook
- publish articles on historical aspects of analytical chemistry
- report on meetings

There will also be a news section open for voluntary contributions, a book review section and a diary of forthcoming events.

Subscription Information:
1981-82: Volume 1 - issues will be published in March, October, November and December 1981 and thereafter monthly
LIBRARY EDITION - Volume 1
Comprising 16 issues and a compendium consisting of the archival material bound in hard covers with an index to facilitate storage and retrieval - US $133.25 / Dfl. 260.00*

SPECIAL INTRODUCTORY OFFER VALID UNTIL DECEMBER 1981

PERSONAL EDITION - Volume 1
(in 16 issues)
US $42.50 (U.S.A. and Canada)
£20.00 (U.K.)
Dfl. 91.50 (Europe)
Dfl. 95.00 (Rest of the World)
All subscriptions for the Personal Edition must be prepaid
Prices include air delivery worldwide

ELSEVIER

Free sample copies will be available on request

P.O. Box 211, 1000 AE Amsterdam, The Netherlands

52 Vanderbilt Ave., New York, N.Y. 10017, USA.

** The Dutch guilder price for the Library Edition is definitive. US $ price is subject to exchange rate fluctuation.*

A unique concept in liquid chromatography columns: 13cm of analytical column with a replaceable front end.

Brownlee Labs introduces a 13cm solution to the problem of short life and high replacement cost. We have combined a 3cm guard cartridge with a 10cm analytical cartridge to form the MPLC™ System. The direct coupling of these two cartridges, via the new union and holder, allows operation to 7,000 psi.

Longer column life lowers costs
Columns die because of sample and mobile phase contamination. Unlike conventional columns, however, the first 3cm of the MPLC System can be easily replaced at about 25% of the cost of a new column.

High performance
The 4.6mm i.d. cartridges are slurry packed with either 5 or 10μm sorbents.

Each 10cm cartridge is individually computer tested, and both the test chromatogram and performance data are supplied.

Versatile
The Brownlee MPLC concept of cartridge columns offers you a range of thirteen different sorbents in two lengths which can be coupled or used separately.

Order from a Brownlee Labs distributor, or write for more information: Brownlee Labs Inc., 2045 Martin Ave., Santa Clara, CA 95050, USA. Telephone 408/727 1346. Telex 171 156.

MPLC™ System. For economy and performance, no LC lab can afford to be without it.

Reference number **BL 13** A

13cm solution. **MPLC** System



Brownlee Labs

This journal now available on MICROFILM



Microfilm Specifications

- 16 mm or 35 mm: positive or negative film supplied in plastic labelled storage boxes: cartridge or cassette
- Polyester base silver halide film
- Reduction ratio 24 X
- Guaranteed standards — NBS resolution charts on each 100 ft reel
- 2A cine mode double page spread across the film

Back volumes up to the end of the 1980 subscription year may be obtained on microfilm

- **Annual Updates** — By summer of each year updates of all journals completing 4 or more volumes since the last update will become available on microfilm.
- **Complete Sets** — A complete set comprises all available volumes on microfilm for one title. A discount of 10% is included in the price quoted for complete sets.
- **Single Volumes** — Individual volumes may be purchased of any title
- **Continuation Orders** — Purchasers may enter a continuation order, and so receive automatically a proforma invoice for periodic updates. These orders will then be processed immediately upon receipt of payment.

Orders and information requests concerning back volumes on microfilm should be addressed exclusively to:

ELSEVIER SEQUOIA S.A.
P.O. Box 851
CH-1001 **Lausanne 1**
Switzerland

Reader service card

(use this card for more information on the news-section and advertisement pages)

I would like to receive, without any obligation, further information on
News items: _____
Advertisements: _____

I am especially interested in the fields as indicated below:

6040 Analytical Chemistry
6050 Chromatography
6150 Mass Spectrometry
6140 Spectroscopy
6240 Pharmaceutical Chemistry
6241 Medicinal Chemistry
6242 Clinical Chemistry
6260 Environmental Chemistry

PLEASE PRINT

Name: _____
Position: _____
Address: _____

THIS CARD IS REPLY PAID IF MAILED IN THE U.S.A.

Reader service card

(use this card for more information on the news-section and advertisement pages)

I would like to receive, without any obligation, further information on
News items: _____
Advertisements: _____

I am especially interested in the fields as indicated below:

6040 Analytical Chemistry
6050 Chromatography
6150 Mass Spectrometry
6140 Spectroscopy
6240 Pharmaceutical Chemistry
6241 Medicinal Chemistry
6242 Clinical Chemistry
6260 Environmental Chemistry

PLEASE PRINT

Name: _____
Position: _____
Address: _____

Reader service card

(use this card for more information on the news-section and advertisement pages)

I would like to receive, without any obligation, further information on
News items: _____
Advertisements: _____

I am especially interested in the fields as indicated below:

6040 Analytical Chemistry
6050 Chromatography
6150 Mass Spectrometry
6140 Spectroscopy
6240 Pharmaceutical Chemistry
6241 Medicinal Chemistry
6242 Clinical Chemistry
6260 Environmental Chemistry

PLEASE PRINT

Name: _____
Position: _____
Address: _____

Please stamp here

ELSEVIER SCIENTIFIC PUBLISHING COMPANY

News Section-Journal of Chromatography

P.O. Box 211

1000 AE AMSTERDAM - The Netherlands

No Postage Necessary if mailed in the United States

BUSINESS REPLY CARD

First Class Permit No. 6794 New York, N.Y.

Postage Will Be Paid By Addressee

ELSEVIER NORTH-HOLLAND, INC.
Attn: Promotion manager Chemistry
52 VANDERBILT AVENUE
NEW YORK, N Y 10164

Please stamp here

ELSEVIER SCIENTIFIC PUBLISHING COMPANY

News Section-Journal of Chromatography

P.O. Box 211

1000 AE AMSTERDAM - The Netherlands

JOURNAL
of
CHROMATOGRAPHY

INTERNATIONAL JOURNAL ON CHROMATOGRAPHY,
ELECTROPHORESIS AND RELATED METHODS

Cumulative Author and Subject Indexes
Volumes 181-200 (1980)

ELSEVIER SCIENTIFIC PUBLISHING COMPANY
AMSTERDAM
1980

Electronic data processing:
Büro für Satztechnik W. Meyer KG
Hanau, G.F.R.

© ELSEVIER SCIENTIFIC PUBLISHING COMPANY — 1981

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system or transmitted in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, without the prior written permission of the publisher, Elsevier Scientific Publishing Company, P.O. Box 330, 1000 AH Amsterdam, The Netherlands.

Submission of an article for publication implies the transfer of the copyright from the author(s) to the publisher and entails the authors' irrevocable and exclusive authorization of the publisher to collect any sums or considerations for copying or reproduction payable by third parties (as mentioned in article 17 paragraph 2 of the Dutch Copyright Act of 1912 and in the Royal Decree of June 20, 1974 (S. 351) pursuant to article 16 b of the Dutch Copyright Act of 1912) and/or to act in or out of Court in connection therewith.

Special regulations for readers in the U.S.A. This journal has been registered with the Copyright Clearance Center, Inc. Consent is given for copying of articles for personal or internal use, or for the personal use of specific clients. This consent is given on the condition that the copier pays through the Center the per-copy fee stated in the code on the first page of each article for copying beyond that permitted by Sections 107 or 108 of the U.S. Copyright Law. The appropriate fee should be forwarded with a copy of the first page of the article to the Copyright Clearance Center, Inc., 21 Congress Street, Salem, MA 01970, U.S.A. If no code appears in an article, the author has not given broad consent to copy and permission to copy must be obtained directly from the author. All articles published prior to 1980 may be copied for a per-copy fee of US$ 2.25, also payable through the Center. This consent does not extend to other kinds of copying, such as for general distribution, resale, advertising and promotion purposes, or for creating new collective works. Special written permission must be obtained from the publisher for such copying.

Special regulations for authors in the U.S.A. Upon acceptance of an article by the journal, the author(s) will be asked to transfer copyright of the article to the publisher. This transfer will ensure the widest possible dissemination of information under the U.S. Copyright Law.

Printed in The Netherlands

AUTHOR INDEX

VOLUMES 181—200

The volume number is placed in parentheses before the page number

A

Aaes-Jørgensen, T.
Specific HPLC method for estimation of the *cis*(*Z*)- and *trans*(*E*)-isomers of clopenthixol and a N-dealkyl metabolite (183) 239

Abbott, S.R., see Wehr, C.T. (185) 453

Abe, I. and Musha, S.
Resolution of amino acid enantiomers by glass capillary GC on easily prepared optically active stationary phases (200) 195

Abidi, S.L.
GLC of straight chain homologues of alkylbenzyldimethylammonium compounds (200) 216

Abraham, B., see Stan, H.-J. (195) 231

Ackermann, G., see Frey, H.-P. (198) 357

Adachi, S., see Ishida, M. (189) 421

Adam, M., see Svojtková, E. (183) 221

Adams, M.J.
GC adsorption studies on synthetic imogolite (188) 97

Adlakha, R.C. and Villanueva, V.R.
Automated ion-exchange chromatographic analysis of usual and unusual natural polyamines (187) 442

Adlard, E.R., Bowen, A.W. and Salmon, D.G.
Automatic system for the high-resolution GC analysis of gasoline-range hydrocarbon mixtures (186) 207

Agoh, T., Konishi, M. and Mori, Y.
Determination of peptido-aminobenzophenone (2-*o*-chlorobenzoyl-4-chloro-N-methyl-N'-glycylglycinanilide) by electron-capture GLC (182) 171

Agris, P.F., Tompson, J.G., Gehrke, C.W., Kuo, K.C. and Rice, R.H.
HPLC and MS of transfer RNA bases for isotopic abundance (194) 205

Ahmed, M., see Gault, M.H. (182) 465

Ahmed, M.S., Dobberstein, R.H. and Farnsworth, N.R.
Stevia rebaudiana. I. Use of *p*-bromophenacyl bromide to enhance UV detection of water-soluble organic acids (steviolbioside and rebaudioside B) in HPLC analysis (192) 387

Ahokas, J.T., Davies, C. and Ravenscroft, P.J.
Simultaneous analysis of disopyramide and quinidine in plasma by HPLC (183) 65

Åhs, U., see Hartvig, P. (183) 229

Aigner-Held, R. and Aue, W.A.
GC with *in situ* photolysis (189) 127

Aigner-Held, R., Aue, W.A. and Pickett, E.E.
LC on metal oxides with silica geometry (189) 139

Aigner-Held, R., see Aue, W.A. (189) 119

Airoldi, L., see Bidoli, F. (196) 314

Aitoku, A., see Shibamoto, T. (194) 277

Akagi, R., see Ubuka, T. (188) 442

Akhavan, I., Wrolstad, R.E. and Richardson, D.G.
Relative effectiveness of ion-exchange and lead acetate precipitation methods in isolating pear sugars and acids (190) 452

Akiki, C., Bourbouze, R., Luporsi, C. and Percheron, F.
Isolation of water-soluble plant glycoproteins by affinity chromatography on immobilised concanavalin A (in French) (188) 435

Al-Thamir, W.K., Purnell, J.H. and Laub, R.J.
Enhancement of GSC column performance by inert solid dilution (188) 79

Albert, K., see Bayer, E. (186) 497

Albet, C., Sánchez, M.G. and Colomé, J.
Determination of a new mucolytic drug Adamexina in biological liquids by photodensitometry (181) 504

Albro, P.W. and Parker, C.E.
General approach to the fractionation and class determination of complex mixtures of chlorinated aromatic compounds (197) 155

Alexander, G., see Garzó, G. (191) 253

Alexander, G., see Szathmáry, C. (191) 327

Allan, R.J., Goodman, H.T. and Watson, T.R.
Two HPLC determinations for mebendazole and its metabolites in human plasma using a rapid Sep Pak C18 extraction (183) 311

Allen, B.A. and Newman, R.A.
HPLC separation of clinically important folic acid derivatives using ion-pair chromatography (190) 241

Alpert, A.J. and Regnier, F.E.
Preparation of a porous microparticulate anion-exchange chromatography support for proteins (185) 375
Altenburg, K.
Dependence of the relative molar response of isomeric alkanes on molecular structure (in German) (194) 213
Alvinerie, M.
Reversed-phase HPLC of phenylbutazone in body fluids (181) 132
Amadò, R., see Waibel, R. (197) 86
Amato, G. d', see D'Amato, G. (196) 245
Ames, M.M., see Powis, G. (181) 95
Amiot, J. and Brisson, G.J.
Continuous automatic nitrogen determination for gel chromatography of protein enzymatic hydrolysates (193) 496
Amsterdam, E.A., see Gallaher, C. (181) 490
Anderson, G.M., Batter, D.K., Young, J.G., Shaywitz, B.A. and Cohen, D.J.
Simplified LC-electrochemical determination of norepinephrine and dopamine in rat brain (181) 453
Anderson, R.A., see Thomson, I. (188) 357
Anderson, T.J., see Lombrozo, L. (181) 1
Andersson, K.K., Balny, C., Douzou, P. and Bieth, J.G.
One-step purification of human leukocyte elastase by biospecific affinity chromatography at subzero temperatures (192) 236
Ando, I., see Ujimoto, K. (190) 161
Andronikashvili, T.G., Laperashvili, L.Y. and Kvernadze, T.K.
Use of volcanic slag as a support in GC (191) 323
Angelo, H.R., Christensen, J.M., Kristensen, M. and McNair, A.
GC method for the simultaneous determination of hydralazine and its acetylated metabolite in serum using a nitrogen-selective detector (183) 159
Anguille, J., see Masson, P. (192) 402
Anikeyev, V.I., see Volkov, S.M. (190) 445
Aniline, O., see Pitts, F.N. Jr. (193) 157
Anmo, T., see Seta, K. (181) 311
Antal, E., Mercik, S. and Kramer, P.A.
Technical considerations in the GC analysis of desipramine (183) 149
Anttila, M., see Nieminen, A.-L. (181) 11
Araki, M., see Tanaka, N. (199) 327
Araki, S., see Watabe, K. (192) 89
Araki, Y., see Shinozawa, S. (190) 489, (196) 463
Arancibia, E.L. and Catoggio, J.A.
GC study of solution and adsorption of hydrocarbons on glycols. I. Diethylene glycol and triethylene glycol (197) 135
Aratskova, A.A., see Kiselev, A.V. (195) 205
Arbin, A., Brink, H. and Vessman, J.
Alkylation of carboxylic acids by solid-liquid phase-transfer catalysis for determination by GC. II. (196) 255
Arbin, A. and Östelius, J.
Determination by electron-capture GC of mono- and di(2-ethylhexyl) phthalate in intravenous solutions stored in poly(vinyl chloride) bags (193) 405
Arfwidsson, A., see Poole, C.F. (199) 123
Arimori, S., see Hiramatsu, K. (182) 301
Armentrout, D.N., see Bouyoucos, S.A. (189) 61
Arnold, P., see Voelter, W. (199) 345
Arora, Y.K., see Bajaj, K.L. (196) 309
Arpino, P.J., Guiochon, G., Krien, P. and Devant, G.
Optimization of the instrumental parameters of a combined liquid chromatograph-mass spectrometer coupled by an interface for direct liquid introduction. I. Performance of the vacuum equipment (185) 529
Arx, E. von, see Von Arx, E. (193) 511
Ascalone, V.
Determination of chlordiazepoxide and its metabolites in human plasma by reversed-phase HPLC (181) 141
Ashes, J.R., Haken, J.K. and Mills, S.C.
GC of esters. XIII. Interrelationship of equivalent chain length *(ECL)* and retention index values of fatty esters (187) 297
Ashley, J.J., see Petersen, M.C. (183) 131
Assenza, S.P. and Brown, P.R.
Comparison of HPLC serum profiles of humans and dogs (181) 169
Assenza, S.P., see Hartwick, R.A. (186) 647
Astier, A. and Deutsch, A.M.
HPLC determination of hippuric acid in human urine. Preliminary results for normal urine levels (182) 88
Atkinson, D.J., see Linnett, L.M. (197) 1
Atrat, P., Deppmeyer, V. and Hörhold, C.
Efficient purification of a microbial steroid-1-dehydrogenase by electrophoretic desorption from the affinity matrix on a preparative scale (189) 279
Atwood, J.G., see Golay, M.J.E. (186) 353
Aue, W.A. and Aigner-Held, R.
Construction of a simple gas chromatograph with photolytic intersect (189) 119
Aue, W.A., Daniewski, M.M. and Pickett, E.E.
Synthesis and GC use of metal-containing surfaces (189) 13
Aue, W.A. and Kapila, S.
Studies and speculations on an alternative mechanism for the response of direct-current electron-capture detectors (188) 1
Aue, W.A. and Wickramanayake, P.P.
GC and LC on silica-supported chromia layers (197) 21
Aue, W.A. and Wickramanayake, P.P.
"Bonded phases" from cyclic organosilicons for GC and LC (200) 3
Aue, W.A., see Aigner-Held, R. (189) 127, (189) 139
Aue, W.A., see Flinn, C.G. (186) 299
Aue, W.A., see Kapila, S. (195) 17, (196) 397
Aue, W.A., see Siu, K.W.M. (189) 255, (192) 413
Aue, W.A., see Wickramanayake, P.P. (195) 25

Auld, W.H.R., see Marcantonio, L.A. (183) 118
Austin, J.H., see James, G.T. (195) 287
Avnir, D., see Morrison, H. (183) 83
Axelson, M. and Sjövall, J.
Strong non-polar cation exchangers for the separation of steroids in mixed chromatographic systems (186) 725
Ayala, J.A., Wentworth, W.E. and Chen, E.C.M.
Comparative investigation of radioactive sources for the electron capture detector (195) 1
Ayala, J.A., see Chen, E.C.M. (188) 89
Ayers, J.S., see Hearn, M.T.W. (185) 463
Ayres, J.W., see Block, J.H. (193) 111

B

Baars, A.M., see Guelen, P.J.M. (181) 497
Baba, S. and Kasuya, Y.
Synchronized accumulating radioisotope detector for radio GC (196) 144
Bachmann, H., see Jurenitsch, J. (189) 43
Bächmann, K., see Rudolph, J. (187) 319
Bächmann, K., see Vierkorn-Rudolph, B. (186) 219
Bachner, L., Boissel, J.-P. and Wajcman, H.
New sensitive technique for the quantitative analysis of initiation peptides (193) 491
Backman, L.
Protein-protein interactions studied by counter-current distribution. II. Test of the theoretical model using bovine serum albumin and L-tryptophan (196) 207
Baechler, J., see Gautam, S.R. (182) 482
Baerheim Svendsen, A., see Verpoorte, R. (184) 79
Bafus, D.A., see Zlatkis, A. (182) 137
Bagon, D.A. and Purnell, C.J.
Determination of airborne free monomeric aromatic and aliphatic isocyanates by HPLC (190) 175
Bailer, P., see Gips, H. (193) 322
Bailey, E. and Barron, E.J.
Determination of tranylcypromine in human plasma and urine using high-resolution GLC with nitrogen-sensitive detection (183) 25
Bailey, E., Farmer, P.B. and Lamb, J.H.
The enantiomer as internal standard for the quantitation of the alkylated amino acid S-methyl-L-cysteine in haemoglobin by GC-chemical ionisation MS with single ion detection (200) 145
Bajaj, K.L. and Arora, Y.K.
Alkaline chloramine-T reagent for the detection of phenolic compounds on thin-layer plates (196) 309
Baker, J.K., see Ma, C.-Y. (200) 163
Bakker, J.A., see Gelsema, W.J. (196) 51
Balasubramanian, R.
Trypan blue staining of proteins fractionated by acrylamide disc gel electrophoresis (189) 410
Balla, J., see János, É. (191) 239
Ballantine, J.A., Williams, K. and Morris, R.J.
Marine sterols. XIII. Evaluation of glass support-coated open tubular columns, prepared by the mercury-plug dynamic technique, for marine sterol analysis (198) 111
Ballerini, R., Chinol, M. and Ghelardoni, M.
Quantitative HPLC determination of Δ^4-3-ketosteroids in adrenocortical extracts (193) 413
Balny, C., see Andersson, K.K. (192) 236
Balogh, S., see Varga-Puchony, Z. (196) 185
Bancher, E., see Wurst, F. (188) 452
Band, P.R., see Besner, J.-G. (183) 346
Bangah, M., Jackman, G. and Bobik, A.
Determination of pindolol in human plasma by HPLC (183) 255
Bannier, A. and Brazier, J.L.
Determination of isofezolac in biological fluids by reversed-phase LCC (182) 369
Bao Miansheng, see Zhang Yukuei (197) 97
Bapat, B.V., see Naikwadi, K.P. (195) 309
Barbier, F., see Beke, R. (193) 504
Barchas, J.D., see Mefford, I.N. (181) 187
Barends, D.M., Van der Sandt, J.S.F. and Hulshoff, A.
Micro determination of gentamicin in serum by HPLC with UV detection (182) 201
Barendse, G.W.M., Van de Werken, P.H. and Takahashi, N.
HPLC of gibberellins (198) 449
Barford, R.A., Sliwinski, B.J. and Rothbart, H.L.
Interactions of proteins with bonded-phase ion exchangers (185) 393
Barkhan, P., see Haroon, Y. (200) 293
Barkley, R.M., see Sievers, R.E. (186) 3
Barlow, G.B., see Sampson, B. (183) 9
Barnes, A.D., see Robinson, J.L. (189) 145
Barnes, J.A., see Kennedy, J.F. (196) 379
Barnes, R.M., see Estes, S.A. (196) 265
Barnes, S., Pritchard, D.G., Settine, R.L. and Geckle, M.
Preparation and characterisation of permethylated derivatives of bile acids, and their application to GC analysis (183) 269
Barni, N., see Guerrini, A.M. (189) 440
Barrientos, A.R., see Cohen, J.L. (181) 478
Barron, E.J., see Bailey, E. (183) 25
Barrow, S.E., Taylor, A.A., Horning, E.C. and Horning, M.G.
HPLC separation and isolation of quinidine and quinine metabolites in rat urine (181) 219
Barry, C. and Pike, R.K.
Analysis of barban formulations by HPLC (195) 151

Barth, H.G. and Regnier, F.E.
High-performance gel permeation chromatography of water-soluble cellulosics (192) 275
Barthová, J., see Tichá, M. (194) 183
Bartik, T., Jánosi, A. and Simon, Á.
Qualitative and quantitative analysis of propylnaphthalenes by GC on silicone stationary phases (197) 92
Bartolucci, S.R., see Twomey, T.M. (183) 104
Basedow, A.M., Ebert, K.H., Ederer, H.J. and Fosshag, E.
Fractionation of polymers by gel permeation chromatography: an experimental and theoretical approach (192) 259
Baselt, R.C. and Franch, S.
Plasma and urine concentration of methapyrilene by nitrogen-phosphorus GLC (183) 234
Basile, B., see Middleditch, B.S. (199) 161
Bates, O.J., see Bradshaw, R.A. (187) 27
Batley, M., Packer, N.H. and Redmond, J.W.
HPLC of diglyceride *p*-nitrobenzoates. An approach to molecular analysis of phospholipids (198) 520
Battaglino, G., see Quercia, V. (193) 163
Batter, D.K., see Anderson, G.M. (181) 453
Bauer, M. and Untz, G.
Analysis of cinchona alkaloids by HPLC (in French) (192) 479
Baughman, R.A. Jr., see Lin, E.T. (183) 367
Baumgartner, P., see Székely, G. (186) 575
Bayer, E., Albert, K., Nieder, M., Grom, E. and Keller, T.
On-line coupling of HPLC and NMR (186) 497
Beaune, P., see Pileire, B. (182) 269
Beck, K.R., Leibowitz, B.J. and Ladisch, M.R.
Separation of methylol derivatives of imidazolidines, urea and carbamates by LC (190) 226
Beck, O. and Hesselgren, T.
Method for the determination of tryptophan in serum and cerebrospinal fluid (181) 100
Beijersbergen van Henegouwen, G.M.J., see Schüsler-van Hees, M.T.I.W. (196) 101
Beijnen, J.H., see Dekker, D. (193) 480
Beke, R., De Weerdt, G.A. and Barbier, F.
Separation of bile acids on Chromarods: a new qualitative and semi-quantitative TLC technique (193) 504
Belfroid, L., see Driessen, O. (181) 441
Belikov, V.A., see Shatz, V.D. (200) 105
Belisle, A.A., see Gay, M.L. (187) 153
Bellemare, G.R., see Roy, C. (197) 121
Belokleytseva, G.M., see Korol, A.N. (194) 145
Benecke, I., see König, W.A. (195) 292
Benet, L.Z., see Lin, E.T. (183) 367
Benjannet, S., see Seidah, N.G. (193) 291
Bens, G.A., see Van den Eeckhout, E. (193) 255
Benson, J.R., see Bradshaw, R.A. (187) 27
Berendsen, G.E. and De Galan, L.
Role of the chain length of chemically bonded phases and the retention mechanism in reversed-phase LC (196) 21
Berenson, G.S., see Radhakrishnamurthy, B. (192) 307
Berezkin, V.G., Budantseva, M.N. and Dows, E.
Chromatographic analysis of major trace components in solvents with concentration prior to sample injection (191) 309
Berezkin, V.G., Kolomiets, L.N., Korolev, A.A., Shmidel, Y.B. and Chizhkov, V.P.
Compound chromatography (191) 95
Berezkin, V.G., Sorokina, E.J., Shaligin, G.F., Litvinov, I.A. and Budantseva, M.N.
Electron-microscopic investigation of the sorbents used in GLC (193) 132
Berg, S., see Jonsson, A. (190) 97
Berlin, J., see Sasse, F. (194) 234
Bernard, N., Cuisinaud, G., Jozefczak, C., Seccia, M., Ferry, N. and Sassard, J.
Sensitive GC method for the determination in plasma of FM 24, 1-(2-*exo*-bicyclo[2,2,1]hept-2-ylphenoxy)-3-[(1-methylethyl)amino]-2-propanol, a new β-adrenoceptor blocking agent (183) 99
Bernard, N., see Cuisinaud, G. (181) 399
Bernasconi, M.G. and Casciola, M.
Surface ion exchange and adsorption of some dyes on α-$Zr(HPO_4)_2 \cdot H_2O$ microcrystals (195) 270
Berndt, W.O., see Phillips, R.D. (190) 419
Berry, V.V.
"Sequential isocratic step" LC: a high-throughput, problem-solving approach with sensitive, near-universal detection for wide polarity mixtures (199) 219
Bertani-Dziedzic, L., see Krstulovic, A.M. (186) 733
Besner, J.-G., Leclaire, R. and Band, P.R.
HPLC of 13-*cis*-retinoic acid and of endogenous retinol in human plasma (183) 346
Bethell, G.S., see Hearn, M.T.W. (185) 463
Betty, K.R., see Dickson, L.C. (190) 311
Beukers, H., see Driessen, O. (181) 441
Bevenue, A., see Ogata, J.N. (189) 425
Beveridge, T., see Niederberger, W. (182) 454
Bezard, J.A. and Ouedraogo, M.A.
Fractionation of oil triacylglycerols by reversed-phase HPLC (196) 279
Bianchini, J.P. and Gaydou, E.M.
Separation of polymethoxylated flavones by straight-phase HPLC (190) 233
Bianchini, P., Nader, H.B., Takahashi, H.K., Osima, B., Straus, A.H. and Dietrich, C.P.
Fractionation and identification of heparin and other acidic mucopolysaccharides by a new discontinuous electrophoretic method (196) 455
Bicchi, C. and Frattini, C.
Quantitative determination of minor components in essential oils: determination of pulegone in peppermint oils (190) 471
Bidlingmeyer, B.A., Deming, S.N., Price, W.P. Jr., Sachok, B. and Petrusek, M.

Retention mechanism for reversed-phase ion-pair LC (186) 419
Bidoli, F., Airoldi, L. and Pantarotto, C.
Quantitative determination of styrene-7β-oxide in blood by combined GC-multiple ion detection mass fragmentography (196) 314
Bielby, C.R., Gande, A.R., Morgan, E.D. and Wilson, I.D.
Practical aspects of the preparation and chromatography of the trimethylsilyl ethers of ecdysteroids (194) 43
Bielby, C.R., see Wilson, I.D. (194) 343
Biessels, H.W.A., see Huibregtse-Minderhout, L. (196) 425
Bieth, J.G., see Andersson, K.K. (192) 236
Billiet, H.A.H., Keehnen, P.D.M. and De Galan, L.
Single-pump solvent programmer for HPLC using synchronized valve switching (185) 515
Billiet, H.A.H., see Schoenmakers, P.J. (185) 179
Bills, D.D., see Heisler, E.G. (194) 89
Binder, H.
Separation of monosaccharides by HPLC: comparison of UV and RI detection (189) 414
Bishara, R.H., see Jakovljevic, I.M. (192) 425
Bishop, C.A., Harding, D.R.K., Meyer, L.J., Hancock, W.S. and Hearn, M.T.W.
HPLC of amino acids, peptides and proteins. XXI. The application of preparative reversed-phase HPLC for the purification of a synthetic underivatised peptide (192) 222
Bishop, G., see Vergin, H. (183) 383
Biswas, S.K., see Mandal, S.B. (188) 292
Bjornsson, T.D., Wolfram, K.M., Routledge, P.A. and Shand, D.G.
HPLC analysis of sulfinpyrazone and its metabolites in biological fluids (181) 417
Bjornsson, T.D., see Wolfram, K.M. (183) 57
Black, C., Douglas, D.M. and Tanzer, M.L.
Separation of cyanogen bromide peptides of collagen by means of HPLC (190) 393
Blackman, G.L., see Perkal, M. (196) 180
Blair, N.E. and Bonner, W.A.
Quantitative determination of D = L mixtures of optical enantiomers by GC (198) 185
Blair, W.R., see Parks, E.J. (185) 563
Blanken, W.M., see Gelsema, W.J. (196) 51, (198) 301
Blaser, A., see Gross, D. (198) 389
Blass, K.G., Briand, R.L., Ng, D.S. and Harold, S.
Miniature two-dimensional TLC separation of lecithin and sphingomyelin (182) 311
Blattná, J., see Pařízková, H. (191) 301
Bloch, J.M., see Wilhelm, J.C. (193) 329
Block, J.H., Ayres, J.W., Henry, D.R. and Levine, H.L.
Use of flow programming in paired-ion HPLC analysis of dosage forms containing dyphylline (193) 111
Blomberg, L. and Wännman, T.
Glass capillary columns coated with non-soluble methyl silicone gums (186) 159
Błotny, G., Kuśmierz, J., Maliński, E. and Szafranek, J.
GC and MS properties of ethyl esters of dithiocarbamic acids of primary and secondary amines (193) 61
Bloxham, D.P., see Corina, D.L. (198) 287
Bobik, A., see Bangah, M. (183) 255
Bobik, A., see Jackman, G.P. (182) 277
Bobleter, O., see Concin, R. (198) 131
Boček, P., Dolník, V., Deml, M. and Janák, J.
Separation and determination of the degradation products of tributyl phosphate by high-speed analytical isotachophoresis (195) 303
Boček, P., Ryšlavý, Z., Deml, M. and Janák, J.
High-speed isotachophoresis. Effect of driving current on the qualitative interpretation of isotachopherograms (191) 271
Boček, P., see Gebauer, P. (199) 81
Bochner, F., see Ross-Lee, L.M. (183) 175
Böcker, R.
Rapid analysis of doxycycline from biological samples by HPLC (187) 439
Bode, J., Schröter, H. and Maass, K.
Dissolvable disulphide-polyacrylamide gels for the electrophoretic analysis of chromosomal proteins and for affino-electrophoresis of thiol-proteins (190) 437
Bodily, K.C., see Klemm, L.H. (198) 1
Boer, F. de, see De Boer, F. (195) 407
Boeva, V.I., see Sakodynskii, K.I. (186) 227
Boisseau, J., see Mourot, D. (190) 486
Boissel, J.-P., see Bachner, L. (193) 491
Bollinger, M.J., see Sievers, R.E. (186) 3
Bombardelli, E., Bonati, A., Gabetta, B. and Martinelli, E.M.
GLC method for determination of ginsenosides in *Panax ginseng* (196) 121
Bonati, A., see Bombardelli, E. (196) 121
Bonazzi, P., see Marcucci, F. (198) 180
Bonchev, D., see Papazova, D. (188) 297
Bonner, F.T., see Pearsall, K.A. (200) 224
Bonner, W.A., see Blair, N.E. (198) 185
Bootsman, T.C., see Van Vliet, H.P.M. (185) 483
Borel, J.P., see Szymanowicz, A.G. (190) 457
Bos, P., see Hanekamp. H.B. (186) 489
Bosch, N. van den, see Van den Bosch, N. (183) 49
Bosisio, A.B., Loeherlein, C., Snyder, R.S. and Righetti, P.G.
Titration curves of proteins by combined isoelectric focusing-electrophoresis in highly porous polyacrylamide matrices (189) 317
Bouman, H., see Greving, J.E. (186) 683
Bourbouze, R., see Akiki, C. (188) 435
Bourne, D.W.A., see Gautam, S.R. (182) 482
Bouska, J.B. and Phillips, S.F.
Simple method for GLC analysis of polyethylene glycol 400 in biological fluids (183) 72

Bouyoucos, S.A. and Armentrout, D.N.
Determination of organophosphoric and organophosphorothioic acids in aqueous solutions by ion chromatography (189) 61
Boven, M. van, see Van Boven, M. (182) 435
Bowen, A.W., see Adlard, E.R. (186) 207
Bowers, W.D., see Clement, R.E. (189) 53, (190) 136
Bowker, M.J., Gladwin, R.P., Mills, D.J. and Stubbles, J.A.
Semi-automated HPLC method for the determination of diacetolol hydrochloride in medicated animal diets (200) 261
Bradshaw, R.A., Bates, O.J. and Benson, J.R.
Peptide separations on substituted polystyrene resins. Effect of cross-linkage (187) 27
Branfman, A.R., Bruni, R.J., Merrill, Y.N., Chadwick, M., Strong, J.M. and Ervin, T.J.
Quantitation of the antitumor agent N-(phosphonacetyl)-L-aspartic acid in human plasma and urine by GC-MS-selected ion monitoring (182) 163
Brater, C., see Seiwell, R. (182) 257
Bratzler, R.L., see Ivory, C.F. (198) 354
Brazeli, R., see Zlatkis, A. (182) 137
Brazell, R., see Poole, C.F. (187) 331
Brazier, J.L., see Bannier, A. (182) 369
Breckenridge, W.C., see Kuksis, A. (182) 1
Bredael, P. and Rietvelde, D.
Identification of the pyrolysis products of *cis*-decaline (in French) (192) 216
Bressolle, F., see Siouffi, A.M. (186) 563
Breter, H.-J., Zöllner, E.J. and Zahn, R.K.
Specificity of deoxyribonuclease hydrolysis determined by anion-exchange HPLC (193) 175
Briand, R.L., see Blass, K.G. (182) 311
Brinckman, F.E., Jewett, K.L., Iverson, W.P., Irgolic, K.J., Ehrhardt, K.C. and Stockton, R.A.
Graphite furnace atomic absorption spectrophotometers as automated element-specific detectors for HPLC. The determination of arsenite, arsenate, methylarsonic acid and dimethylarsinic acid (191) 31
Brinckman, F.E., see Parks, E.J. (185) 563
Brink, H., see Arbin, A. (196) 255
Brinkman, U.A.T. and De Vries, G.
Use of chemically bonded stationary phases in HPTLC. II (192) 331
Brinkman, U.A.T., De Vries, G. and Cuperus, R.
Stationary phases for HPLC (198) 421
Brinkman, U.A.T., see Lawrence, J.F. (185) 473
Brinkman, U.A.T., see Scholten, A.H.M.T. (199) 239
Brinkman, U.A.T., see Van Vliet, H.P.M. (185) 483
Brisson, G.J., see Amiot, J. (193) 496
Bristol, D.W.
Detection of trace organic impurities in binary solvent systems: a solvent purity test (188) 193
Brodie, R.R., Chasseaud, L.F., Walmsley, L.M., Soegtrop, H.H., Darragh, A. and O'Kelly, D.A.
Determination of the antispasmodic agent ethaverine in human plasma by HPLC (182) 379
Brooks, M.A. and Dixon, R.
Determination of estramustine and its 17-keto metabolite in plasma by HPLC (182) 387
Brötell, H.
Syringe and column adsorption of tertiary amines in GC (196) 489
Brown, A., see Middleditch, B.S. (195) 359
Brown, L.W., see Hundt, H.K.L. (183) 378
Brown, N.D., Scovill, J.P., Sleeman, H.K. and Doctor, B.P.
Determination of adiphenine hydrochloride and diphenylacetic acid by ion-pair HPLC (200) 267
Brown, N.D., Sleeman, H.K., Doctor, B.P. and Scovill, J.P.
Determination of aprophen and its hydrolytic by-products by ion-pair HPLC (195) 146
Brown, P.M. and Uppal, G.S.
Improved clean-up procedure for the determination of small residues of carbophenothion in mice (190) 201
Brown, P.R., see Assenza, S.P. (181) 169
Brown, P.R., see Halfpenny, A.P. (199) 275
Brown, P.R., see Hartwick, R.A. (186) 647, (186) 659
Brown, R.R., see Patterson, J.I. (182) 425
Broxterman, H.J., Noach, E.L. and De Vos, D.
Assay of HA-966 in rat plasma by capillary GLC with nitrogen-selective detection (183) 185
Brucker, J. de, see De Brucker, J. (187) 181
Bruins, A.P., see Hendriks, H. (190) 321
Bruni, R.J., see Branfman, A.R. (182) 163
Brunner, G., see Holloway, C.J. (188) 235
Brůnová, B., see Kuchař, M. (195) 329
Bruyn, C. de, see De Bruyn, C. (182) 317
Bryce, T.A. and Burrows, J.L.
Determination of oxpentifylline and a metabolite, 1-(5'-hydroxyhexyl)-3,7-dimethylxanthine, by GLC using a nitrogen-selective detector (181) 355
Brynda, E., Štrop, P., Mikeš, F. and Kálal, J.
Polarity of some chromatographic materials intended for the separation of biopolymers: spectroscopic label technique (196) 39
Buchanan, D.N., Fucek, F.R. and Domino, E.F.
Paired-ion HPLC assay for plasma choline (181) 329
Büchler, W.
The role of chromatography in Ciba-Geigy (Review) (184) 207
Buck, R.P., see Walters, R.R. (188) 61
Buckle, K.A., see Tangendjaja, B. (197) 274
Budantseva, M.N., see Berezkin, V.G. (191) 309, (193) 132

Budd, R.D.
GC separations of dialkyl barbiturate derivatives (192) 212
Budd, R.D., Yang, F.C. and Leung, W.J.
Mass screening and confirmation of methaqualone and its metabolites in urine by radioimmunoassay-TLC (190) 129
Budna, K.W., see Lankmayr, E.P. (198) 471
Buffa, M., Congiu, G., Lombard, A. and Tourn, M.L.
A homologous series of non-reducing oligosaccharides in *Artemisia absinthium* L. roots (200) 309
Bui-Nguyên, M.H.
Application of HPLC to the separation of ascorbic acid from isoascorbic acid (196) 163
Bunch, J.E. and Pellizzari, E.D.
Evaluation of chromatographic sorbents used in air pollution studies (186) 811
Burkhard, J., see Vodička, L. (198) 457, (200) 238
Burns, D.T., Glockling, F. and Harriott, M.
Comparative assessment of GLC and HPLC for the separation of tin tetraalkyls and alkyltin halides (200) 305
Burns, D.T. and Mackay, C.
Comparison of reversed-phase and adsorption modes of HPLC for the assay of fat soluble vitamins in multivitamin tablets (200) 300
Burns, D.T., MacKay, C. and Tillman, J.
Rationalisation of the chromatographic behaviour of vitamin D_2/D_3 and related compounds in adsorption HPLC (190) 141
Burnsed, G.E., see Heath, R.R. (189) 199
Burrows, J.L., see Bryce, T.A. (181) 355
Burton, G., see Nordlöv, H. (190) 221
Burtscher, E., see Concin, R. (198) 131
Busbee, D.L., see Marshall, M.V. (197) 217
Bush, B., see Smith, R.M. (181) 319
Bush, K., see Kirschbaum, J. (190) 481
Buskin, J.N., see Upton, R.A. (190) 119
Bye, A., see Lo, L.Y. (181) 473

C

Cacchi, S., see Gasparrini, F. (194) 239
Caccia, S., Guiso, G. and Zanini, M.G.
GLC determination of 4'-hydroxynomifensine in biological samples (190) 475
Cadet, J., Voituriez, L., Hahn, B.S. and Wang, S.Y.
Separation of cyclobutyl dimers of thymine and thymidine by HPLC and TLC (195) 139
Caglioti, L., see Gasparrini, F. (194) 239
Calabro, G., see Micali, G. (194) 245
Caldicott, A.B., see Minnikin, D.E. (188) 221
Caldwell, K.D., see Giddings, J.C. (185) 261
Campbell, B., see Pattinson, N. (187) 409
Canobbio, L., see Marcucci, F. (198) 180
Capp, M., see Nice, E.C. (185) 413
Capp, M., see O'Hare, M.J. (198) 23
Caprioli, R.M., see Lin, S.-N. (197) 31
Cardaci, V. and Ossicini, L.
Paper electrophoretic study of ion-pair formation. XIII. Behaviour of anionic trisoxalatocobalt(III) and trisoxalatochromium(III) in optically active electrolytes (198) 76
Cardwell, T.J., Marriott, P.J. and McDonough, P.S.
GC of volatile metal dialkyldithiophosphates (193) 53
Carlson, R.E. and Dolphin, D.
HPLC method for the analysis of isoflavones (198) 193
Carr, P.W.
Fundamental, predictive approach to dipole-dipole interactions based on the dipole moment of the solute and the dielectric constant of the solvent (194) 105
Carroll, F.I., see Lewin, A.H. (193) 371
Carruthers, M., see Nelson, L.M. (183) 295
Carson, V.J., see Jackman, G.P. (182) 277
Carter, D.E., see Karakaya, A. (195) 431
Cartier, P., see Pileire, B. (182) 269
Carver, J.R., see Henson, J.B. (198) 443
Casciola, M., see Bernasconi, M.G. (195) 270
Cashion, P.J., see Leigh, C.P.H. (192) 490
Casida, K.C., see Casida, M.E. (200) 35
Casida, M.E. and Casida, K.C.
Application of a unified theory of GC photoionization detector response to a 10.2-eV detector (200) 35
Cassady, J.M., see Doetsch, P.W. (189) 79
Cassidy, R.M., Harpur, R. and Elchuk, S.
Determination of multifunctional carboxylic acids in the presence of iron by GC (190) 188
Castagnoli, N. Jr., see De Ruyter, M.G.M. (183) 193
Castegnaro, M., see Walker, E.A. (187) 229
Castello, G. and D'Amato, G.
GC analysis of light hydrocarbons for the characterization of Porapak columns (196) 245
Catoggio, J.A., see Arancibia, E.L. (197) 135
Čatská, V., see Patzelová, V. (193) 137
Caude, M., see Foucault, A. (185) 345
Caude, M., see Gareil, P. (192) 53
Caude, M., see Guyon, F. (186) 677
Ćelap, M.B., Hodžić, I.M. and Janjić, T.J.
Resolution of neutral complexes of transition metals by stereoselective adsorption on optically active complexes. I. Partial resolution of tris(acetylacetonato)cobalt(III) and tris(acetylacetonato)chromium(III) on cobalt(III) complexes (198) 172
Ćelap, M.B., Vučković, G., Malinar, M.J., Janjić, T.J. and Radivojša, P.N.
Effect of the composition and structure of

cobalt(III) complexes on their R_F values obtained by TLC on silica gel (196) 59
Čeřovský, V., Tichá, M., Turková, J. and Labský, J.
Interaction of trypsin with immobilized *p*-aminobenzamidine derivatives studied by means of affinity electrophoresis (194) 175
Chadwick, M., see Branfman, A.R. (182) 163
Chafetz, L., see Philip, J. (192) 250
Chakrabarti, P., see Mandal, S.B. (188) 292
Chan, J.A., see Poehland, B.L. (190) 246
Chan, K.K., see Cohen, J.L. (181) 478
Chan, P.K., Siraj, M.Y. and Hayes, A.W.
HPLC analysis of the mycotoxin penicillic acid and its application to biological fluids (194) 387
Chan, W.Y., Seale, T. and Rennert, O.M.
Fluorescence micro disc electrophoresis in sodium dodecyl sulfate. A simple and sensitive method applicable for routine laboratory analysis (181) 259
Chandrarajan, J. and Klein, L.
Chromatographic separation and quantification of type I and type III collagens (182) 94
Chang, S.-L. and Kramer, W.G.
Improvements to a HPLC assay for allopurinol and oxipurinol in plasma (181) 286
Chaplin, R.P., see Nieass, C.S. (194) 335
Chapman, R.A., see Robinson, J.R. (193) 213
Charles, R. and Gil-Av, E.
N-Docosanoyl-L-valine-2(2-methyl)-*n*-heptadecylamide as a stationary phase for the resolution of optical isomers in GLC (195) 317
Chasseaud, L.F., see Brodie, R.R. (182) 379
Chauhan, J. and Darbre, A.
Determination of homovanillic, isohomovanillic and vanillylmandelic acids in human urine by means of glass capillary GLC with temperature-programmed electron-capture detection (183) 391
Chauhan, L.S., see Srivastava, S.P. (196) 225
Chawla, H.M., Gambhir, I. and Kathuria, L.
Separation and identification of some hydroxyacetophenones and their derivatives by TLC (188) 289
Cheetham, N.W.H. and Sirimanne, P.
Separation of partially methylated sugars by reversed-phase HPLC (196) 171
Chen, B.-K., see Melander, W.R. (185) 99
Chen, E.C.M., Wentworth, W.E. and Ayala, J.A.
Oxygen enhancement of the electron capture detector response of carbon dioxide (188) 89
Chen, E.C.M., see Ayala, J.A. (195) 1
Chen, E.C.M., see Wentworth, W.E. (186) 99
Chen Hu, L.-E., see Poole, C.F. (187) 331
Chen, S.S. and Hsu, D.-S.
Operational parameters of anion-exchange chromatography using AG MP-1 resin for rapid assay of adenine nucleotides (198) 500
Chen, S.S., see Hsu, D.-S. (192) 193
Chenery, R.J.
Use of bromine as an aid in the identification of unsaturated fatty alcohols and diols (189) 263
Chiang, T.-C.
GC-MS assay for low levels of retinoic acid in human blood (182) 335
Chinol, M., see Ballerini, R. (193) 413
Chiu, A., see Warsh, J.J. (183) 483
Chiu, C.C., see Poehland, B.L. (190) 246
Chizhkov, V.P., see Berezkin, V.G. (191) 95
Chornet, E., see Roy, C. (197) 121
Chou, M.W. and Yang, S.K.
Combined reversed-phase and normal-phase HPLC in the purification and identification of 7,12-dimethylbenz[*a*]anthracene metabolites (185) 635
Choudhary, G.
GLC determination of toxic diamines in permanent hair dyes (193) 277
Choudhary, G.
GLC determination of 2-mercaptoethanol (200) 211
Choudhary, V.R. and Sansare, S.D.
GC technique for temperature-programmed desorption of reversibly and irreversibly adsorbed species (192) 420
Chow, F., see Karmen, A. (199) 113
Chow, F., see Lam, S. (199) 295
Chow, F.K. and Grushka, E.
HPLC of nucleotides and nucleosides using outersphere and innersphere metal-solute complexes (185) 361
Chow, F.K., see Grushka, E. (199) 283
Choy, E.T. and Meisen, A.
GC detection of diethanolamine and its degradation products (187) 145
Chrétien, J.R., see Dubois, J.-E. (194) 121
Chrétien, M., see Seidah, N.G. (193) 291
Christensen, J.M., see Angelo, H.R. (183) 159
Christophersen, A.S. and Rasmussen, K.E.
Glass capillary column GC of barbiturates after flash-heater derivatization with dimethylformamide dimethylacetal (192) 363
Chriswell, C.D., see Richard, J.J. (199) 143
Churáček, J., see Jandera, P. (192) 1, (192) 19, (192) 37, (197) 181
Churchill, F.C. II and Ku, D.N.
Extractive alkylation of 5,2'-dichloro-4-nitrosalicylanilide (niclosamide) for GLC analysis (189) 375
Číhová, A., see Šmejkal, F. (197) 147
Claeys, A.E., see Lefevere, M.F. (186) 749
Clark, C.R. and Noggle, F.T. Jr.
LC identification of clorazepate in pharmaceutical products (188) 426
Clark, E.C., see Hundt, H.K.L. (182) 110, (183) 378
Clark, P.M.S., Kricka, L.J. and Whitehead, T.P.
Comparison of urinary excretion of UV-absorbing constituents in healthy subjects and patients with rheumatoid arthritis using analytical isotachophoresis (181) 347

Clark, T.C., see Honigberg, I.L. (181) 266
Clement, R.E., Eiceman, G.A., Karasek, F.W., Bowers, W.D. and Parsons, M.L.
Rapid analysis of chlorinated dioxins in complex organic mixtures by application of a GC-MS-calculator system with user-developed software (189) 53
Clement, R.E., Karasek, F.W., Bowers, W.D. and Parsons, M.L.
Correction for artifacts in the analysis of atmospheric aerosols (190) 136
Clement, R.E., see Dickson, L.C. (190) 311
Clementi, W.A., see Haegele, K.D. (187) 171
Clements, J.A., Hasson, K. and Smith, G.
Determination of isoprenaline by ion-pair HPLC (189) 272
Cochrane, W.P., see Greenhalgh, R. (188) 305
Cohen, D.J., see Anderson, G.M. (181) 453
Cohen, J.L., Hisayasu, G.H., Barrientos, A.R., Nayar, M.S.B. and Chan, K.K.
Reversed-phase HPLC analysis of methotrexate and 7-hydroxymethotrexate in serum (181) 478
Cohen, M.B., Maybaum, J. and Sadée, W.
Analysis of purine ribonucleotides and deoxyribonucleotides in cell extracts by HPLC (198) 435
Colantuoni, G.
Mass, charge, momentum and energy conservation in electrophoretic fractionation systems (Review) (184) 21
Cole, E.R., Crank, G. and Hai Minh, H.T.
Chromatography of 1,3-benzodioxole derivatives (193) 19
Colin, H., Martin, M. and Guiochon, G.
Extra-column effects in HPLC. I. Theoretical study of the injection problem (185) 79
Collins, D., see Pattinson, N. (187) 409
Colomé, J., see Albet, C. (181) 504
Colussi, D., see Sioufi, A. (182) 241
Colvin, M., see Strife, R.J. (182) 211
Concin, R., Burtscher, E. and Bobleter, O.
Chromatographic behaviour of aromatic compounds on Sephadex LH gels. Calibration of gel columns for determination of molecular weight distributions (198) 131
Congiu, G., see Buffa, M. (200) 309
Connole, E., see James, G.T. (195) 287
Connor, J.
The electron-capture detector. I. Theoretical model (200) 15
Cooke, M., Nickless, G. and Roberts, D.J.
Carbon skeleton-GC techniques and their applications (187) 47
Cooke, M. and Ober, A.G.
OV-17-QF-1 capillary column for organochlorine pesticide analysis (195) 265
Cooke, M. and Roberts, D.J.
Carbon skeleton capillary GC (193) 437
Cooke, M., see Ober, A.G. (196) 237
Cooper, G.K., see Corina, D.L. (198) 287
Čopíková, J., see Vrátný, P. (191) 313
Coq, B., Cretier, G. and Rocca, J.L.
Preparative LC: sample volume overload (186) 457
Corcia, A. di, see Di Corcia, A. (198) 347
Corina, D.L., Bloxham, D.P. and Cooper, G.K.
GC and GC-MS characterisation of methane thiolsulphonates carrying further functional groups (198) 287
Corradini, D. and Sinibaldi, M.
Gel filtration of soluble coloured metal hexacyanoferrate(II) compounds (187) 458
Costa, E., see Peralta, E. (190) 43
Côté, J., see Garneau, F.-X. (188) 445
Cotellessa, L., Riva, R., Salvà, P., Marcucci, F. and Mussini, E.
Quantitative determination of flufenamic acid in rat plasma and uterus by GC (192) 441
Cotterill, E.G.
Determination of residues of methazole and its metabolites, 1-(3,4-dichlorophenyl)-3-methylurea and 1-(3,4-dichlorophenyl)urea in soil by HPLC (197) 267
Cottrell, R.C., see Walters, D.G. (196) 501
Čoupek, J., see Mikeš, O. (192) 159
Court, W.E. and Iwu, M.M.
Chromogenic reactions of *Rauwolfia* alkaloids after separation by TLC (187) 199
Cousineau, J., see Green, J.R. (188) 439
Coutts, R.T., Dawe, R., Jones, G.R., Liu, S.-F. and Midha, K.K.
Analysis of perfluoroacyl derivatives of ephedrine, pseudoephedrine and analogues by GC and MS. I. Trifluoroacetyl derivatives (190) 53
Coutts, R.T., Hargesheimer, E.E. and Pasutto, F.M.
Application of a direct aqueous acetylation technique to the GC quantitation of nitrophenols and 1-naphthol in environmental water samples (195) 105
Cozamanis, I., see Reece, P.A. (181) 427
Crabbe, P.G. and Fryer, C.
Rapid quantitative analysis of solasodine, solasodine glycosides and solasodiene by HPLC (187) 87
Crampton, E.L., Glass, R.C., Marchant, B. and Rees, J.A.
Chemical ionisation mass fragmentographic measurement of dothiepin plasma concentrations following a single oral dose in man (183) 141
Crank, G., see Cole, E.R. (193) 19
Crathorne, B., Watts, C.D. and Fielding, M.
Analysis of non-volatile organic compounds in water by HPLC (185) 671
Cretier, G., see Coq, B. (186) 457
Crommen, J.
Reversed-phase ion-pair HPLC of drugs and related compounds using underivatized silica as the stationary phase (186) 705
Crommen, J.
Ion-pair chromatography in the low concentration range by use of highly absorbing counter ions. III. HPLC of

quaternary alkylammonium ions as ion pairs with naphthalene-2-sulphonate, using a silica support of low surface area (193) 225
Cronnelly, R., see De Ruyter, M.G.M. (183) 193
Crozier, A., Zaerr, J.B. and Morris, R.O. High-performance steric exclusion chromatography of plant hormones (198) 57
Cuisinaud, G., Ferry, N., Seccia, M., Bernard, N. and Sassard, J. Determination of nalidixic acid and its two major metabolites in human plasma and urine by reversed-phase HPLC (181) 399
Cuisinaud, G., see Bernard, N. (183) 99
Culvenor, C.C.J., see Frahn, J.L. (195) 379
Cummings, G., see Karmen, A. (199) 113
Cunningham, M.D., see Desiderio, D.M. (195) 369
Cuperus, R., see Brinkman, U.A.T. (198) 421
Curro, P., see Micali, G. (194) 245
Curtius, H.-C., Farner, H. and Rey, F. *In vivo* studies of the tryptophan-5-hydroxylase system. Quantitation of serotonin and tryptamine using GC-mass fragmentography (199) 171
Czech, B., see Ogierman, L. (198) 536

D

Dąbrowski, R., see Rayss, J. (188) 107
Dąbrowski, R., see Witkiewicz, Z. (200) 65
Dady, P.J., see Taylor, G.A. (183) 421
Daenens, P., see Van Boven, M. (182) 435
Dahlström, H., Leander, K., Lindström, B. and Molander, M. Quantitative determination of benzilonium bromide in plasma by GC-MS after oxidation to benzophenone (183) 511
Dale, G., see Tarbit, I.F. (181) 337
Dalferes, E.R. Jr., see Radhakrishnamurthy, B. (192) 307
D'Amato, G., see Castello, G. (196) 245
Damiani, D.E., Valles, E.M. and Gigola, C.E. GC determination of adsorption-desorption rates. Interaction of hydrogen on a nickel catalyst (196) 355
Damme, J.-C. van, see Van Damme, J.-C. (190) 401
Damon, C.E. and Pettitt, B.C. Jr. HPLC determination of denatonium benzoate in rapeseed oil (195) 243
Daniewski, M.M., see Aue, W.A. (189) 13
Danis, P.O., see Maya, L. (190) 145
Danzinger, R.G., see Torchia, M.G. (181) 120
Darbre, A., see Chauhan, J. (183) 391
Darke, D.J. and Roediger, W.E.W. Method for the measurement of hydroxylamine in colonic fluid using derivatisation and GC (181) 449
Darragh, A., see Brodie, R.R. (182) 379
Dautrevaux, M., see Lustenberger, P. (193) 451
Davies, C., see Ahokas, J.T. (183) 65
Davies, G., see Lindner, W. (185) 323
Davis, B.A., see Hui, K.-S. (192) 341
Davis, D.Y., see Honigberg, I.L. (181) 266
Davis, P.J. HPLC determination of glaucine and its metabolite, dehydroglaucine in microbial culture (193) 170
Davis, P.J. and Klein, A.E. HPLC separation of colchicine and its phenolic and N-desacetylated derivatives (188) 280
Dawe, R., see Coutts, R.T. (190) 53
Dawkins, J.V. and Yeadon, G. High-performance gel permeation chromatography of polystyrene with silica microspheres (188) 333
Dawson, R.M.C., see Johnson, R.C. (187) 235
Day, R.S. Quantitation of porphyrin esters on thin-layer chromatograms (198) 227
De Boer, F., see Huizing, H.J. (195) 407
De Brucker, J., see Nagels, L. (187) 181
De Bruyn, C., see Verheggen, T. (182) 317
De Galan, L., see Berendsen, G.E. (196) 21
De Galan, L., see Billiet, H.A.H. (185) 515
De Galan, L., see Schoenmakers, P.J. (185) 179
De Heaulme, M., see Pernes, J.-F. (181) 254
De Jong, A.W.J., Kraak, J.C., Poppe, H. and Nooitgedacht, F. Isotherm linearity and sample capacity in LC (193) 181
De Jong, G.J. The use of a pre-column for the direct HPLC determination of the antidepressants clovoxamine and fluvoxamine in plasma (183) 203
De Jong, W., see Tonnaer, J.A.D.M. (183) 303
De Leenheer, A.P., see Gelijkens, C.F. (183) 78, (194) 305
De Leenheer, A.P., see Lefevere, M.F. (186) 749
De Leenheer, A.P., see Nelis, H.J.C.F. (195) 35
De Ligny, C.L., see Gelsema, W.J. (196) 51, (198) 301
De Meo, M., see Wheeler, L.A. (183) 357
De Moerloose, P., see Van den Eeckhout, E. (193) 255
De Pooter, H., see Nagels, L. (187) 181
De Ruyter, M.G.M., Cronnelly, R. and Castagnoli, N. Jr. Reversed-phase, ion-pair LC of quaternary ammonium compounds. Determination of pyridostigmine, neostigmine and edrophonium in biological fluids (183) 193
De Silva, J.A.F., see Min, B.H. (183) 411
De Silva, J.A.F., see Strojny, N. (181) 272
De Vos, D., see Broxterman, H.J. (183) 185
De Vos, D., see Van den Bosch, N. (183) 49
De Vos, R.H., see Van der Poll, J.M. (187) 244
De Vries, C.P., Lomecky-Janousek, M. and Popp-Snijders, C.

Rapid quantitative assay of plasma 11-deoxycortisol and cortisol by HPLC for use in the metyrapone test (183) 87
De Vries, G., see Brinkman, U.A.T. (192) 331, (198) 421
De Weerdt, G.A., see Beke, R. (193) 504
De Zeeuw, R.A., see Drenth, B.F.H. (191) 109
De Zeeuw, R.A., see Greving, J.E. (186) 683
Dean, S., see Jack, D.B. (187) 277, (196) 189
Debeaupuis, J.P., see Lafont, P. (198) 481
Debeuf, C., see Nagels, L. (190) 411
Dębowski, J., see Kutner, W. (191) 47
Deelder, R.S., Linssen, H.A.J., Konijnendijk, A.P. and Van de Venne, J.L.M.
Retension mechanism in reversed-phase ion-pair chromatography of amines and amino acids on bonded phases (185) 241
Degenhardt, C.E.A.M. and Duvalois, W.
Simple device to facilitate the deactivation of a GC system by PEG 20M via the gas phase (192) 418
Dekker, D. and Beijnen, J.H.
Improved HPLC separation of decomposition products of prednisolone by adding sulphite to the mobile phase (193) 480
Delépine, B., see Mourot, D. (190) 486
Della Ragione, F., see Zappia, V. (189) 399
Dellacherie, E., see Hubert, P. (184) 325
Delver, B., see Vink, J. (181) 115
Deming, S.N., see Bidlingmeyer, B.A. (186) 419
Deming, S.N., see Kong, R.C. (199) 307
Deming, S.N., see Sachok, B. (199) 317
Demjaněnko, M., see Gaš, B. (192) 253
Deml, M., see Boček, P. (191) 271, (195) 303
Deml, M., see Gebauer, P. (199) 81
Deng Li-Ru
Comprehensive study of porous polymeric stationary phases for chromatography (186) 317
Deppmeyer, V., see Atrat, P. (189) 279
Desideri, P.G., see Lepri, L. (195) 65, (195) 187, (195) 339
Desiderio, D.M., Stein, J.L., Cunningham, M.D. and Sabbatini, J.Z.
HPLC and field desorption MS of hypothalamic oligopeptides (195) 369
Deutsch, A.M., see Astier, A. (182) 88
Devant, G., see Arpino, P.J. (185) 529
Dewaele, C. and Verzele, M.
HPLC of beer bitter acids (197) 189
Deyl, Z., see Macek, K. (193) 421
Deyl, Z., see Svojtková, E. (183) 221
Di Corcia, A., Samperi, R. and Severini, C.
GC column for the rapid determination of congeners in potable spirits (198) 347
Di Ferrante, D.T., Wilson, N.Y. and Leach, C.S.
Chromatographic method for the measurement of hydroxylysine, hydroxylysine glycosides and 3-methylhistidine in human urine (187) 271
Dickson, L.C., Clement, R.E., Betty, K.R. and Karasek, F.W.
Software improvements for GC-MS calculator system used in the analysis of trace organic compounds from environmental samples (190) 311
Dietrich, C.P., see Bianchini, P. (196) 455
Dilli, S. and Pillai, D.
Analysis of trace amounts of barbiturates in saliva (190) 113
Dilli, S. and Weekley, A.
Direct derivatization and GC determination of barbiturates in autopsy liver tissues (195) 349
Dimenna, G.P., Krick, T.P. and Segall, H.J.
Rapid HPLC isolation of monoesters, diesters and macrocyclic diester pyrrolizidine alkaloids from *Senecio jacobaea* and *Amsinckia intermedia* (192) 474
Dimov, N., see Papazova, D. (188) 297
Dirr, E., see Speck, M. (200) 313
DiSanzo, F.P., see Uden, P.C. (196) 403
Distler, W.
Modified preparation of 2-naphthacyl esters of fatty acids and their fluorometric detection in HPLC (in German) (192) 240
Dixon, R., see Brooks, M.A. (182) 387
Dizdaroglu, M. and Simic, M.G.
Separation of underivatized dipeptides by HPLC on a weak anion-exchange bonded phase (195) 119
Dizdaroglu, M., Simic, M.G. and Schott, H.
Separation and sequencing of the sequence isomers of pyrimidine deoxypentanucleoside tetraphosphates by HPLC (188) 273
Dobashi, A., see Hara, S. (186) 543
Dobberstein, R.H., see Ahmed, M.S. (192) 387
Doctor, B.P., see Brown, N.D. (195) 146, (200) 267
Doctor, V.M., see Holzer, G. (194) 410
Dodeur, M. and Jacquet, M.A.
Interactions of insolubilized lectins with membrane glycoproteins in presence of detergents (195) 197
Doerr, R.C., Parris, N. and Parks, O.W.
Determination of sulfanilic acid in the presence of sulfanilamide and some sulfa drugs by reversed-phase ion-pair HPLC (196) 498
Doetsch, P.W., Cassady, J.M. and McLaughlin, J.L.
Cactus alkaloids. XL. Identification of mescaline and other β-phenethylamines in *Pereskia, Pereskiopsis* and *Islaya* by use of fluorescamine conjugates (189) 79
Dolan, J.W. and Snyder, L.R.
Coated-open-tubular chromatography with flow segmentation. II. Experimental study and optimization for size-exclusion separation (185) 57
Dolan, J.W., see Gant, J.R. (185) 153
Dolan, J.W., see Snyder, L.R. (185) 43
Dolník, V., see Boček, P. (195) 303
Dolphin, D., see Carlson, R.E. (198) 193
Dolphin, D., see Silverman, R.B. (194) 273

Domino, E.F. and Domino, S.E.
GC-MS analysis of polybrominated biphenyl constituents of Firemaster FF-1 (197) 258
Domino, E.F., see Buchanan, D.N. (181) 329
Domino, S.E., see Domino, E.F. (197) 258
Done, A.K., see Miceli, J.N. (183) 250
Dongen, W. van, see Van Dongen, W. (187) 181
Donini, P., see Guerrini, A.M. (189) 440
Donnet, J.-B., see Kessaissia, Z. (196) 481
Doolittle, R.E., see Heath, R.R. (189) 199
Doran, D., see Smith, R.M. (181) 319
Dorrbecker, B., see Kullberg, M.P. (187) 264
Douglas, D.M., see Black, C. (190) 393
Doumas, J., see Maume, B.F. (186) 581
Dousek, F.P., see Smolková, E. (191) 61
Douzou, P., see Andersson, K.K. (192) 236
Dowling, R.H., see Podesta, M.T. (182) 293
Downing, D.T. and Stranieri, A.M.
Correction for deviation from the Lambert-Beer law in the quantitation of thin-layer chromatograms by photodensitometry (192) 208
Dows, E., see Berezkin, V.G. (191) 309
Drenth, B.F.H. and De Zeeuw, R.A.
HPLC in the elucidation of the metabolic fate of butoprozine. Screening for unknown metabolites in the rat (191) 109
Dressler, M., see Teplý, J. (191) 221
Dreux, M., see Lafosse, M. (188) 315, (193) 9
Driessen, O., Beukers, H., Belfroid, L. and Emonds, A.
Analysis of methotrexate by isotachophoresis (181) 441
Driscoll, J.N., see Jaramillo, L.F. (186) 637
Driskell, J.A., see Lim, K.L. (188) 285
Drozd, J., see Rijks, J.A. (186) 167
Du Plessis, L.M., see Van Niekerk, P.J. (187) 436
Dua, V.K., see Srivastava, S.P. (196) 225
Dubois, J.-E., Chrétien, J.R., Soják, L. and Rijks, J.A.
Topological analysis of the behaviour of linear alkenes up to tetradecenes in GLC on squalane (194) 121
Ducharme, D., see Lawrence, A.H. (194) 434
Dufek, P. and Pacáková, V.
Comparison of the HPLC behaviour of *s*-triazine derivatives on various stationary phases (187) 341
Dufek, P., Pacáková, V. and Tesařová, E.
Separation and behaviour of *s*-triazine derivatives on a NH_2-chemically bonded stationary phase by HPLC (191) 115
Duksin, D., see Mahoney, W.C. (198) 506
Dumas, T.
Determination of formaldehyde, acetaldehyde and associated components in solution and in vapours by GC (200) 206
Dupuis, M.D. and Hill, H.H. Jr.
Survey of selected hydrides as doping agents for a hydrogen-atmosphere flame-ionization detector (195) 211
Durand, J.P. and Petroff, N.
Determination of benzo[*a*]pyrene and other polyaromatic hydrocarbons in petroleum oils by direct LC (190) 85
Duranti, M.
Program for processing amino acid data with a programmable pocket calculator (194) 69
Dutt, M.C. and Poh, T.T.
Use of ninhydrin as a spray reagent for the detection of some basic drugs on thin-layer chromatograms (195) 133
Duvalois, W., see Degenhardt, C.E.A.M. (192) 418

E

Eadie, M.J., see Ross-Lee, L.M. (183) 175
Ebert, K.H., see Basedow, A.M. (192) 259
Echols, R.E., Harris, J. and Miller, R.H. Jr.
Modified procedure for determining vitamin B_1 by GC (193) 470
Edelson, J., see Kullberg, M.P. (187) 264
Ederer, H.J., see Basedow, A.M. (192) 259
Edlund, P.O.
Determination of clonidine in human plasma by glass capillary GC with electron-capture detection (187) 161
Eeckhout, E. van den, see Van den Eeckhout, E. (193) 255
Egly, J.-M., see Ochoa, J.L. (188) 257
Egri, L., Egri, L.L., Takács, J.M. and Kralik, D.C.
Error in the determination of the specific retention volume. Specific retention volumes of *n*-alkanes on Apolane-87 stationary phase as functions of column temperature (198) 85
Egri, L.L., see Egri, L. (198) 85
Eguchi, S., see Kono, A. (182) 125
Ehrhardt, K.C., see Brinckman, F.E. (191) 31
Ehrsson, H., see Eksborg, S. (185) 583
Eiceman, G.A. and Karasek, F.W.
Retention behavior of chlorinated benzenes, chlorinated phenols, Aldrin, and Aroclor 1242 on Tenax-GC, Chromosorb 101, Florisil, and Carbopack C HT (200) 115
Eiceman, G.A., see Clement, R.E. (189) 53
Einarsson, R., see Renck, B. (197) 278
Ek, K., see Righetti, P.G. (184) 415
Ekman, L., Lindström, B., Nilsson, G. and Wibell, L.
Determination of benoxaprofen in plasma and urine by LC (182) 478
Eksborg, S., Ehrsson, H. and Lönroth, U.
LC determination of tetracycline in plasma and urine (185) 583
Elchuk, S., see Cassidy, R.M. (190) 188
Elema, E.T., see Jellema, R. (189) 406
Elsohly, M.A., see Ma, C.-Y. (200) 163

Emond, J.P. and Pagé, M.
An approach to ampholyte-displacement chromatography (200) 57
Emonds, A., see Driessen, O. (181) 441
Ende, M., Pfeifer, P. and Spiteller, G.
Contamination by use of Extrelut columns (in German) (183) 1
Endo, M., Imamichi, H., Moriyasu, M. and Hashimoto, Y.
Microdetermination of stimulant drugs in urine by HPLC (196) 334
Eng Tho Gan, I., see Schier, G.M. (182) 232
Engelhardt, H. and Mathes, D.
High-performance exclusion chromatography of water-soluble polymers with chemically bonded stationary phases (185) 305
Engelhardt, H., see Martin, A.J.P. (186) 15
Englmaier, P.
Trimethylsilyl esters of plant acids and their application in GLC. Preparation, kinetics of silylation and solvent effects on quantitative reaction and stability of the derivatives (in German) (194) 33
Engström, N., Hellgren, L. and Vincent, J.
Instant copy technique for documentation of thin-layer chromatograms (189) 284
Enkhardt, U. and Gräser, H.
Separation of benzoic acid derivatives from plant extracts on Sephadex G-10 columns (in German) (198) 373
Enomoto, J., see Shiono, S. (193) 243
Erickson, C.K., see Goldman, M.E. (190) 217
Ericson, J., see Lesko, L.J. (182) 226
Eriksen, P.B., see Knudsen, P.J. (187) 373
Eriksson, B.-M., Persson, B.-A. and Lindberg, M.
Determination of apomorphine in plasma and brain tissue by ion-pair extraction and LC (185) 575
Erixson, E., see Westerlund, D. (185) 593
Ernst, G.F. and Van der Kaaden, A.
HPLC analysis of furazolidone in liver and kidney (198) 526
Ervik, M., Kylberg-Hanssen, K. and Lagerström, P.-O.
Electron-capture-GC determination of atenolol in plasma and urine, using a simplified procedure with improved selectivity (182) 341
Ervin, H.E. and McGinnis, G.D.
Analysis of pentachlorophenol in waste water using HPLC (190) 203
Ervin, T.J., see Branfman, A.R. (182) 163
Eskin, N.A.M., see Hoehn, E. (194) 251
Eskola, P., see Tolan, J.W. (190) 367
Esmans, E., see Nagels, L. (190) 411
Estes, S.A., Poirier, C.A., Uden, P.C. and Barnes, R.M.
GC with plasma emission spectroscopic detection of Friedel-Crafts catalyzed alkyl group redistribution products among Si, Ge, Sn and Pb atoms (196) 265
Ettre, L.S.
Relative retention expressions in chromatography (198) 229
Evans, G.W., see Johnson, P.E. (188) 405
Everaerts, F., see Mikkers, F. (182) 496
Everaerts, F., see Verheggen, T. (182) 317
Evershed, R.P., see Morgan, E.D. (186) 605
Evrard, E., see Malvoisin, E. (186) 81
Evtushenko, E.V., Vakhrusheva, N.M. and Ovodov, Y.S.
Preparative LCC of methyl ethers of methyl α-L-rhamnopyranoside (196) 331

F

Facchini, V., Streeter, A.J. and Timbrell, J.A.
Determination of hydralazine and its acetylated metabolites in urine by GC and HPLC (187) 218
Facchini, V. and Timbrell, J.A.
Determination of hydralazine metabolites: 4-hydrazino-phthalazin-1-one and N-acetylhydrazinophthalazin-1-one by GC and *s*-triazolo[3,4-*a*]phthalazine and phthalazinone by HPLC (183) 167
Fagerlund, C., see Hartvig, P. (181) 17
Faiman, M.D., see Jensen, J.C. (181) 407
Fairclough, R.J., see Lauren, D.R. (200) 288
Fales, H.M., see Jaouni, T.M. (182) 473
Farmer, P.B., see Bailey, E. (200) 145
Farner, H., see Curtius, H.-C. (199) 171
Farnsworth, N.R., see Ahmed, M.S. (192) 387
Farnworth, E.R., see Kramer, J.K.G. (198) 279
Farooqui, A.A.
Purification of enzymes by heparin-Sepharose affinity chromatography (Review) (184) 335
Faupel, M., Felix, H.R. and Von Arx, E.
Fast and simple method for the separation of intermediates and cofactors involved in the biosynthesis of cephalosporin C using chemically bonded C_{12} reversed-phase TLC (193) 511
Fehsenfeld, F.C., see Sievers, R.E. (186) 3
Felix, H.R., see Faupel, M. (193) 511
Fell, A.F., Plag, S.M. and Neil, J.M.
Stability-indicating assay for azathioprine and 6-mercaptopurine by reversed-phase HPLC (186) 691
Felt, V. and Hušek, P.
Effect of derivative structure on flame-ionization detector response of amino acid oxazolidinones (197) 226
Fenger, M., see Knudsen, P.J. (187) 373
Fenimore, D.C. and Meyer, C.J.
Contact spotting. A new approach to sample application in TLC (186) 555
Feraud, J.P., see Gareil, P. (192) 53

Ferrante, D.T. di, see Di Ferrante, D.T. (187) 271
Ferry, N., see Bernard, N. (183) 99
Ferry, N., see Cuisinaud, G. (181) 399
Fielding, M., see Crathorne, B. (185) 671
Filonenko, G.V., see Korol, A.N. (194) 145
Finegold, S.M., see Wheeler, L.A. (183) 357
Fink, D.W., see Tolan, J.W. (190) 367
Finlayson, A.J., see MacKenzie, S.L. (187) 239
Finn, J.M., see Pirkle, W.H. (192) 143
Firpo-Pamies, G., see Gassiot-Matas, M. (187) 1
Fitt, L.E., Hassler, W. and Just, D.E.
A rapid and high-resolution method to determine the composition of corn syrups by LC (187) 381
Flanagan, R.J., Storey, G.C.A. and Holt, D.W.
Rapid HPLC method for the measurement of amiodarone in blood plasma or serum at the concentrations attained during therapy (187) 391
Flatmark, T., see Haavik, J. (198) 511
Fleuren, H.L.J., Verwey-van Wissen, C.P.W. and Van Rossum, J.M.
Quantitative GC determination of two oxidized metabolites of the diuretic mefruside in human urine, plasma and red blood cells (182) 179
Flinn, C.G. and Aue, W.A.
GC detection of organogermanium compounds by surface or gas-phase luminescence (186) 299
Floberg, S., Lanbeck, K. and Lindström, B.
Determination of methimazole in plasma using GC-MS after extractive alkylation (182) 63
Florentz, K., see Knudsen, P.J. (187) 373
Foard, D.E., see Lin, K.D. (195) 385
Foglietti, M.J., see Weber, M. (188) 377
Foley, J., see Kirschbaum, J. (200) 282
Fontaine, N., see Szymanowicz, A.G. (190) 457
Formstecher, P., see Lustenberger, P. (193) 451
Fornstedt, N.
HPLC determination of 5-hydroxyindole-3-acetic acid in urine using Sephadex G-10 for isolation (181) 456
Fosshag, E., see Basedow, A.M. (192) 259
Foucault, A., Caude, M. and Oliveros, L.
Ligand-exchange chromatography of enantiomeric amino acids on copper-loaded chiral bonded silica gel and of amino acids on copper(II)-modified silica gel (185) 345
Foucault, A., see Guyon, F. (186) 677
Fouchard, R.C., see Kramer, J.K.G. (198) 279
Frahn, J.L., Culvenor, C.C.J. and Mills, J.A.
Preparative separation of the pyrrolizidine alkaloids, intermedine and lycopsamine, as their borate complexes (195) 379
Franc, J. and Kosíková, E.
Determination of some anions separated by TLC (187) 462
Franch, S., see Baselt, R.C. (183) 234
Francis, J.D., see Nair, B.R. (195) 158
Frattini, C., see Bicchi, C. (190) 471
Fredriksson, S.
Decreased accuracy in isotachophoretic analysis due to septum bleed (188) 262
Fredriksson, S.
Complexation with ferric ions: a potential source of error in the isotachophoretic determination of oxalate (188) 266
Freedman, A.N.
The photoionization detector. Theory, performance and application as a low-level monitor of oil vapour (190) 263
Frei, R.W., see Hanekamp, H.B. (186) 489
Frei, R.W., see Lawrence, J.F. (185) 473
Frei, R.W., see Scholten, A.H.M.T. (199) 239
Frei, R.W., see Van Vliet, H.P.M. (185) 483
Freinkel, N., see Johnson, R.C. (187) 235
Frey, H.-P. and Ackermann, G.
Correction of R_F values according to Galanos and Kapoulas (in German) (198) 357
Frey, J., see Perier, C. (182) 155
Fričová, V., see Kramlová, M. (193) 515
Friedman, M.A., see Unger, P.D. (187) 429
Friedman, Z., Smith, H.W. and Hancock, W.S.
Rapid HPLC method for quantitation of 3-methylhistidine (182) 414
Friel, P.N., see Raisys, V.A. (183) 441
Fritz, D.F., Sahil, A. and Kováts, E. sz.
Study of the adsorption effects at the surface of poly(ethylene glycol)-coated column packings (186) 63
Fritz, J.S., see Gjerde, D.T. (186) 509, (187) 35, (188) 391
Fritz, J.S., see Richard, J.J. (199) 143
Fryer, C., see Crabbe, P.G. (187) 87
Fucek, F.R., see Buchanan, D.N. (181) 329
Fujihara, M., see Iizima, N. (193) 464
Fujita, K., see Ito, S. (187) 418
Fujita, T., see Kimura, M. (188) 417
Fukuhara, K., see Matsuki, Y. (188) 177
Fukunaga, A., Hatta, Y., Ishibashi, M. and Miyazaki, H.
GC of dimethylalkylsilyl ether derivatives of bile acid methyl esters (190) 339
Furukawa, M. and Yokoyama, T.
Determination of free toluene diisocyanate in polyurethane prepolymers by HPLC (198) 212

G

Gabetta, B., see Bombardelli, E. (196) 121
Gáborjányi, R., see János, É. (191) 239
Gabryś, J. and Konecki, J.
GLC of free amino acids in the hyaloplasm of rat cerebral, cerebellar and ocular tissues, and in skeletal and heart muscle (182) 147

Gagné, D. and Lodge, B.A.
Analysis of dexamethasone sodium phosphate formulations by HPLC (193) 160
Gal, J., Marcell, P.D. and Tarascio, C.M.
HPLC micro-assay for chloramphenicol in human blood plasma and cerebrospinal fluid (181) 123
Galácz, J., see Szathmáry, C. (191) 327
Galan, L. de, see De Galan, L. (185) 179, (185) 515, (196) 21
Galensa, R. and Herrmann, K.
Analysis of flavonoids by HPLC (189) 217
Gallaher, C., Henderson, G.L., Low, R.I., Amsterdam, E.A. and Mason, D.T.
Comparison of HPLC with an enzyme multiplied immunoassay technique for the analysis of serum procainamide and N-acetylprocainamide (181) 490
Galletti, P., see Zappia, V. (189) 399
Galoux, M., see Van Damme, J.-C. (190) 401
Gambhir, I., see Chawla, H.M. (188) 289
Gande, A.R., see Bielby, C.R. (194) 43
Ganjian, I., Kubo, I. and Kubota, T.
Rapid and micro identification of biologically active diterpenes in *Rabdosia umbrosus* var. *excisinflexus* (Labiatae) by reversed-phase HPLC (200) 250
Ganschow, I., see Halket, J.M. (192) 434
Gant, J.R., Dolan, J.W. and Snyder, L.R.
Systematic approach to optimizing resolution in reversed-phase LC with emphasis on the role of temperature (185) 153
Garaj, J., see Krupčík, J. (191) 207
Garattini, E., see Gazzotti, G. (188) 400
Garbagna, L., see Marcucci, F. (198) 180
Gareil, P., Personnaz, L., Feraud, J.P. and Caude, M.
Study of non-linear elution in preparative LC (in French) (192) 53
Garland, W.A., see Min, B.H. (183) 411
Garneau, F.-X., Côté, J., Harvey, O. and Simard, J.-L.
Fractionation of steroid digitonides by TLC and CC on silica gel (188) 445
Garzó, G., Alexander, G. and Till, A.
GC determination of 5-alkyl-uracils and 5-alkyl-deoxyuridines using flash methylation and glass capillary columns (191) 253
Gaš, B., Demjaněnko, M. and Vacík, J.
High-frequency contactless conductivity detection in isotachophoresis (192) 253
Gasparič, J.
Use of reversed-phase TLC to asses hydrophobicity in studies of quantitative structure-activity relationships for drugs (196) 391
Gasparrini, F., Cacchi, S., Caglioti, L., Misiti, D. and Giovannoli, M.
Intermediate- and large-scale reversed-phase preparative HPLC on an axially compressed column: a facile, quantitative separation of 7α- and 7β-methyl-17β-acetoxy-3-oxoandrost-4-enes (194) 239
Gassen, H.G., see Jost, W. (185) 403
Gassiot-Matas, M. and Firpo-Pamies, G.
Relationships between GC retention index and molecular structure (187) 1
Gast, C.H., Kraak, J.C., Poppe, H. and Maessen, F.J.M.J.
Capacilities of on-line element-specific detection in HPLC using an inductively coupled argon plasma emission source detector (185) 549
Gattavecchia, E. and Tonelli, D.
TLC of some 5-nitroimidazoles of pharmaceutical interest (193) 340
Gaudio, L.A., see Taylor, R.F. (187) 212
Gault, M.H., Ahmed, M., Tibbo, N., Longerich, L. and Sugden, D.
HPLC method for isolation of tritiated digoxin and metabolites in urine (182) 465
Gautam, S.R., Nahum, A., Baechler, J. and Bourne, D.W.A.
Determination of papaverine in plasma and urine by HPLC (182) 482
Gavrilova, T.B., Kiselev, A.V. and Roshchina, T.M.
Chromatography of hydrocarbons on molybdenum sulphide (192) 323
Gawell, B.-M. and Larsson, B.
Determination of styrene in foods by reversed-phase HPLC (198) 198
Gay, M.L., Belisle, A.A. and Patton, J.F.
Quantification of petroleum-type hydrocarbons in avian tissue (187) 153
Gaydou, E.M., see Bianchini, J.P. (190) 233
Gayot, G., see Mourot, D. (190) 486
Gazda, K., Kamiński, M., Klawiter, J., Kowalczyk, J.S., Makuch, B., Prusiewicz, K. and Śledzińska, B.
Influence of particle parameters on the efficiency of LC systems (191) 9
Gazdag, M., see Szepesi, G. (191) 101
Gazzotti, G., Garattini, E. and Salmona, M.
Improved GC method for measuring phenylethylene glycol. Application to the determination of styrene monooxygenase and epoxide hydrase activities (188) 400
Gebauer, P., Boček, P., Deml, M. and Janák, J.
Isotachophoresis of kinetically labile complexes (199) 81
Geckle, M., see Barnes, S. (183) 269
Geher, K., see Kuksis, A. (182) 1
Gehrke, C.W., Kuo, K.C. and Zumwalt, R.W.
Chromatography of nucleosides (188) 129
Gehrke, C.W., see Agris, P.F. (194) 205
Geissler, H.E., see Schumacher, A. (181) 512
Gelijkens, C.F. and De Leenheer, A.P.
Simple method for isolating free urinary catecholamines by boric acid gel chromatography (183) 78
Gelijkens, C.F. and De Leenheer, A.P.
Reversed-phase LC of 5-fluorouracil nucleosides and nucleotides in the presence of quaternary ammonium ions (194) 305
Gelpí, E., see Martínez, E. (186) 619
Gelpi, E., see Tusell, J.M. (181) 295
Gelsema, W.J., De Ligny, C.L. and Blanken, W.M.

Gel chromatographic comparison of the molecular weight distributions of Ampholine, Servalyte and Pharmalyte carrier ampholytes used in isoelectric focusing (198) 301
Gelsema, W.J., De Ligny, C.L., Blanken, W.M., Hamer, R.J., Roozen, A.M.P. and Bakker, J.A.
Study of the feasibility of separation of carrier ampholytes from peptides by hydrophobic interaction chromatography on octyl-Sepharose (196) 51
Gemeiner, P., see Mislovičová, D. (194) 95
Gerich, J.E., see Haymond, M.W. (183) 403
Gfeller, J.C. and Stockmeyer, M.
HPLC column switching technique in the analysis of medicated feed for an automated clean-up procedure (198) 162
Ghaemi, Y. and Wall, R.A.
Hydrophobic chromatography with dynamically coated stationary phases. III. Non-ionic surfactant effects (198) 397
Ghatge, B.B., see Naikwadi, K.P. (195) 309
Ghelardoni, M., see Ballerini, R. (193) 413
Ghias-Ud-Din, M., Olson, E.B. Jr. and Rankin, J.
Separation and quantitation of [^{14}C]prostaglandin E_1 from lung effluent metabolites by HPLC (192) 463
Gianazza, E. and Righetti, P.G.
Size and charge distribution of macromolecules in living systems (193) 1
Gianazza, E., see Righetti, P.G. (184) 415
Gibbs, V., see Kirschbaum, J. (200) 282
Giddings, J.C., Myers, M.N., Caldwell, K.D. and Pav, J.W.
Steric field-flow fractionation as a tool for the size characterization of chromatographic supports (185) 261
Giddings, J.C., Myers, M.N. and Janča, J.
Retention characteristics of various polymers in thermal field-flow fractionation (186) 37
Gigola, C.E., see Damiani, D.E. (196) 355
Gil-Av, E., see Charles, R. (195) 317
Gilbert, J., Shepherd, M.J., Startin, J.R. and McWeeny, D.J.
GC determination of vinylidene chloride monomer in packaging films and in foods (197) 71
Gilbert, J., Shepherd, M.J. and Wallwork, M.A.
Determination of oligomers in vinyl chloride polymers by steric exclusion chromatography (193) 235
Gilbert, J. and Startin, J.R.
GC-MS of the flophemesyl derivatives of a series of fatty acid methyl ester chlorohydrins (189) 86
Gilbert, M.T., see Knox, J.H. (186) 405
Gilpin, R.K. and Sisco, W.R.
Effect of temperature on precision of retention measurements in LC (194) 285
Ginneken, C.A.M. van, see Van Ginneken, C.A.M. (182) 252, (182) 487, (182) 492
Giovannoli, M., see Gasparrini, F. (194) 239
Gips, H., Korte, K., Meinecke, B. and Bailer, P.
Separation of C_{21}-, C_{19}- and C_{18}-steroids on Sephadex LH-20 microcolumns (193) 322
Gjerde, D.T. and Fritz, J.S.
Chromatographic separation of metal ions on macroreticular anion-exchange resins of a low capacity (188) 391
Gjerde, D.T., Fritz, J.S. and Schmuckler, G.
Anion chromatography with low-conductivity eluents (186) 509
Gjerde, D.T., Schmuckler, G. and Fritz, J.S.
Anion chromatography with low-conductivity eluents. II (187) 35
Glad, M., Ohlson, S., Hansson, L., Månsson, M.-O. and Mosbach, K.
High-performance liquid affinity chromatography of nucleosides, nucleotides and carbohydrates with boronic acid-substituted microparticulate silica (200) 254
Gladwin, R.P., see Bowker, M.J. (200) 261
Glajch, J.L., Kirkland, J.J., Squire, K.M. and Minor, J.M.
Optimization of solvent strength and selectivity for reversed-phase LC using an interactive mixture-design statistical technique (199) 57
Glass, R.C., see Crampton, E.L. (183) 141
Glazunova, L.D., see Sakodynskii, K.I. (186) 227
Glockling, F., see Burns, D.T. (200) 305
Glöckner, G.
Determination of resolution in gel permeation chromatography (191) 279
Glover, D.J., see Hoffsommer, J.C. (195) 435
Godeneche, D., see Michelot, J. (188) 431
Godse, D.D., see Warsh, J.J. (183) 483
Goehl, T.J., see Shiu, G.K. (181) 127
Goins, W. Jr., see Marcelin, G. (187) 57
Golander, Y. and Sternson, L.A.
Paired-ion chromatographic analysis of tamoxifen and two major metabolites in plasma (181) 41
Golay, M.J.E.
The HETP of an unretained component of fluid flow in a coiled round pipe (186) 341
Golay, M.J.E.
The dynamic diffusion constant within fluid flow in an open straight tube with an elliptical cross-section (196) 349
Golay, M.J.E. and Atwood, J.G.
Early phases of the dispersion of a sample injected in Poiseuille flow (186) 353
Goldman, M.E., Hamm, H. and Erickson, C.K.
Determination of melatonin by HPLC with electrochemical detection (190) 217
Golovnya, R.V. and Misharina, T.A.
Determination of the energy of intermolecular interactions in GC from retention indices (190) 1
Gonzalez, M.A., see Marshall, M.V. (197) 217
Goodfellow, M., see Minnikin, D.E. (188) 221
Goodman, H.T., see Allan, R.J. (183) 311
Goodwin, G.H., see Mathew, C.G.P. (198) 80
Gorbunoff, M.J.
Purification of ovomucoid by hydroxyapatite chromatography (187) 224

Goryayev, V.M., see Volkov, S.M. (190) 445
Gossard, F., see Seidah, N.G. (193) 291
Graaff, P.R., see Raisys, V.A. (183) 441
Grabovac, I., see Morris, C.E.M. (189) 259
Gracza, L. and Ruff, P.
Simple method for separation and quantitative determination of kawalactones by HPLC (in German) (193) 486
Graham, S.W., see Wright, B.W. (199) 355
Gräser, H., see Enkhardt, U. (198) 373
Greco, A.V., see Mingrone, G. (183) 277
Green, E.S., see Mendenhall, C.L. (190) 197
Green, J.R. and Cousineau, J.
Purification of phlorizin by CC on Sephadex LH-20 with aqueous propan-2-ol (188) 439
Greenhalgh, R. and Cochrane, W.P.
Use of chlorpyrifos as an evaluation standard for GC detectors (188) 305
Greenland, R., see Wenk, G. (183) 261
Grego, B., see Hearn, M.T.W. (185) 429
Gregorová, I., see Tomsová, Z. (200) 221
Greving, J.E., Bouman, H., Jonkman, J.H.G., Westenberg, H.G.M. and De Zeeuw, R.A.
Analysis of quaternary ammonium compounds and basic drugs based on ion-pair adsorption HPLC (186) 683
Grgurinovich, N. and Miners, J.O.
Simple, rapid procedure for the determination of valproate and ethosuximide in plasma by GLC (182) 237
Gribnau, F.W.J., see Van Loenhout, J.W.A. (182) 487
Griest, W.H. and Martin, T.W.
Stimulated sequential dehydroxy-fluorination: a safe, convenient method for GC analysis of phosphate diesters with possible application to hydrolysates of organophosphorus pesticides and nerve gases (192) 173
Griffin, A.C., see Marshall, M.V. (197) 217
Grimsrud, E.P. and Miller, D.A.
Non-linear responses of the electron-capture detector to alkyl monochlorides (192) 117
Grimsrud, E.P., Miller, D.A., Stebbins, R.G. and Kim, S.H.
Alteration of the electron-capture detector response to polycyclic aromatic hydrocarbons by oxygen doping of the carrier gas (197) 51
Grimsrud, E.P., see Miller, D.A. (190) 133
Grob, K.
GC stationary phases analysed by capillary GC (198) 176
Grob, K. Jr. and Neukom, H.P.
Factors affecting the accuracy and precision of cold on-column injections in capillary GC (189) 109
Grob, K. Jr. and Neukom, H.P.
Should the septum part of vaporizing injectors be kept at lower temperatures? (198) 64
Grob, R.L. and Leasure, J.B.
Study of the kinetics for a catalyzed reaction using GC (197) 129
Groh, R. and Halász, I.
Separation of C_6 cyclic hydrocarbons before the inert peak on microporous silica: a case of negative adsorption (199) 23
Grom, E., see Bayer, E. (186) 497
Gross, D., Gutekunst, H., Blaser, A. and Hamböck, H.
Peak identification in capillary GC by simultaneous flame ionization detection and ^{14}C-detection (198) 389
Gross, R.W. and Sobel, B.E.
Isocratic HPLC separation of phosphoglycerides and lysophosphoglycerides (197) 79
Gruber, V.F., see Walker, R.W. (181) 85
Grumadas, A.J., see Poshkus, D.P. (191) 169
Grushka, E. and Chow, F.K.
HPLC of nucleotide-Mg(II) complexes. Separation and mechanism of separation (199) 283
Grushka, E. and Mott, S.D.
Chromatographic peak shape. III. Influence of kinetics parameters on solute profiles (186) 117
Grushka, E., see Chow, F.K. (185) 361
Grzybowski, J., Lamparczyk, H., Nasal, A. and Radecki, A.
Relationship between the retention indices of phenols on polar and non-polar stationary phases (196) 217
Gübitz, G.
Derivatization of fatty acids with 1-chlormethylisatin for HPLC (187) 208
Guelen, P.J.M., Baars, A.M., Vree, T.B., Nijkerk, A.J. and Vermeer, J.M.
Rapid and sensitive determination of chlorthalidone in blood, plasma and urine of man using HPLC (181) 497
Guelen, P.J.M., see O'Reilly, W.J. (183) 492
Guentert, T.W., Rakhit, A., Upton, R.A. and Riegelman, S.
An integrated approach to measurements of quinidine and metabolites in biological fluids (183) 514
Guentert, T.W., see Upton, R.A. (190) 119
Guerrini, A.M., Barni, N. and Donini, P.
Chromatographic separation and identification of short-chain acid-soluble polyphosphates from *Saccharomyces cerevisiae* (189) 440
Guiochon, G.
Preparation and operation of LC columns of very high efficiency (185) 3
Guiochon, G.
A hazard in high-performance liquid chromatography (189) 108
Guiochon, G., see Arpino, P.J. (185) 529
Guiochon, G., see Colin, H. (185) 79
Guiochon, G., see Siouffi, A.M. (186) 563
Guiso, G., see Caccia, S. (190) 475
Gutekunst, H., see Gross, D. (198) 389

Guyon, F., Foucault, A. and Caude, M.
Ligand-exchange chromatography of small peptides on copper(II)-modified silica gel: application to the study of the enzymatic degradation of methionine-enkephalin (186) 677
Gvozdovitch, T.N., see Kiselev, A.V. (195) 205
Gyllenhaal, O. and Hartvig, P.
Electron capture GC of sulphonamides. Effects of structure and temperature on detector response (189) 351
Gyllenhaal, O., Johansson, L. and Vessman, J.
GC of epinephrine and norepinephrine after derivatization with chloroformates in aqueous media (190) 347
Gyllenhaal, O., see Hartvig, P. (181) 17

H

Haas, R.E., see Van der Meer, C. (182) 121
Haavik, J. and Flatmark, T.
Rapid and sensitive assay of tyrosine 3-monooxygenase activity by HPLC using the native fluorescence of DOPA (198) 511
Haddad, P.R., see Low, G.K.C. (198) 235
Haegele, K.D., Skrdlant, H.B., Talseth, T., McNay, J.L., Shepherd, A.M.M. and Clementi, W.A.
Quantitative analysis of hydralazine pyruvic acid hydrazone, the major plasma metabolite of hydralazine (187) 171
Hageman, J.H., see Maestas, R.R. (189) 225
Hager, D.
Theoretical considerations of molecular sieve effects (187) 285
Hahn, B.S., see Cadet, J. (195) 139
Hai Minh, H.T., see Cole, E.R. (193) 19
Hailey, D.M., see Johnston, M.A. (189) 241
Hais, I.M.
Two-dimensional (Letter to the Editor) (187) 466
Haken, J.K. and Srisukh, D.
Elimination of the reference stationary phase in classification schemes using the molecular retention index (199) 199
Haken, J.K., Srisukh, D. and Vernon, F.
Calcium complex grease as a low-polarity stationary phase (190) 291
Haken, J.K. and Vernon, F.
Evaluation of a thermally stable hydrocarbon as a non-polar base stationary phase for use in Rohrschneider-McReynolds-type schemes (186) 89
Haken, J.K., see Ashes, J.R. (187) 297
Haken, J.K., see Wainwright, M.S. (184) 1, (188) 246, (198) 156
Hakusui, H., Suzuki, W. and Sano, M.
Quantitative analysis of oxepinac in human plasma, urine and saliva by GC-mass fragmentography (182) 47
Hal, H.J.M. van, see Van Hal, H.J.M. (181) 25, (181) 115
Halász, I., see Groh, R. (199) 23
Halász, I., see Hofmann, K. (199) 3
Halász, I., see Martin, A.J.P. (186) 15
Hale, W.B., see Stocklinski, A.W. (195) 281
Halfpenny, A.P. and Brown, P.R.
Optimized assay for purine nucleoside phosphorylase by reversed-phase HPLC (199) 275
Halket, J.M.
Numerical analysis of chromatographic-spectrometric data. Purity of partially resolved peaks and pattern separation of the pure component spectra (186) 443
Halket, J.M., Ganschow, I. and Lisboa, B.P.
GC-MS properties of boronate esters of 24*R*,25-dihydroxycholecalciferol (192) 434
Halket, J.M. and Lisboa, B.P.
Simple mass chromatographic procedure for the detection and identification of sterols using derivative correlations (189) 267
Halkiewicz, J., see Radecki, A. (187) 363
Hall, D.R., see Lester, R. (190) 35
Hallmark, M.R., see Shaikh, B. (195) 392
Hallmark, R.K., see Shaikh, B. (195) 392
Halpern, B.C., see Stea, B. (188) 363
Halpern, R.M., see Stea, B. (188) 363
Hama, T., see Kojima, K. (189) 233
Hamano, T., Mitsuhashi, Y. and Matsuki, Y.
Improved GC method for the quantitative determination of secondary amines as sulphonamides formed by reaction with benzenesulphonyl chloride (190) 462
Hamböck, H., see Gross, D. (198) 389
Hamer, R.J., see Gelsema, W.J. (196) 51
Hamilton, D.J.
GC measurement of volatility of herbicide esters (195) 75
Hamm, H., see Goldman, M.E. (190) 217
Hammer, J., see Sasse, F. (194) 234
Hammers, W.E., see Özcimder, M. (187) 307
Hammond, E.W., see Smith, E.C. (188) 205
Hančil, V., see Rosenbaum, M. (191) 157
Hancock, W.S., see Bishop, C.A. (192) 222
Hancock, W.S., see Friedman, Z. (182) 414
Hancock, W.S., see Hearn, M.T.W. (185) 429, (185) 463
Hanekamp. H.B., Bos, P. and Frei, R.W.
Design and selective application of a dropping mercury electrode amperometric detector in column LC (186) 489
Hanna, S., Insler, M., Zapata, R. and Lachman, L.
HPLC analysis of naloxone hydrochloride in injectable solutions (200) 277
Hannestad, U. and Sörbo, B.
Derivatization of dithiols and certain monothiols with phenylarsine oxide for GC (200) 171
Hanrahan, J.M. and Paterson, A.R.
Adsorption-desorption GC-IR

determination of trace disulfur decafluoride in sulfur hexafluoride (193) 265
Hanrahan, J.M., see Smith, J.S. (193) 271
Hansen, S.H., see Helboe, P. (189) 249
Hansson, L., see Glad, M. (200) 254
Hanukoglu, I. and Jefcoate, C.R.
Pregnenolone separation from cholesterol using Sephadex LH-20 minicolumns (190) 256
Hanya, T., see Matsumoto, G. (193) 89, (194) 199
Hara, S. and Dobashi, A.
LC resolution of enantiomers on normal-phase chiral amide-bonded silica gel. Retentions of optically active α-amino acid derivatives on N-acyl homologues of L-valylaminopropylsilanized silica phases (186) 543
Hara, S. and Ohsawa, A.
Design of binary solvent systems for LSC on silica gel columns using mono- and difunctional steroids as solutes (200) 85
Hara, Y., see Kono, A. (182) 125
Harada, K., see Shibamoto, T. (194) 277
Harapat, S.R. and Kates, R.E.
HPLC analysis of verapamil. II. Simultaneous quantitation of verapamil and its active metabolite, norverapamil (181) 484
Harding, D.R.K., see Bishop, C.A. (192) 222
Hardwick, J. and Hechtman, P.
High-capacity method for purification of human liver hexosaminidase B using hydrophobic chromatography (190) 385
Hargesheimer, E.E., see Coutts, R.T. (195) 105
Harold, S., see Blass, K.G. (182) 311
Haroon, Y., Shearer, M.J. and Barkhan, P.
Resolution of phylloquinone (vitamin K_1), phylloquinone 2,3-epoxide, 2-chloro-phylloquinone and their geometric isomers by HPLC (200) 293
Harpur, R., see Cassidy, R.M. (190) 188
Harrap, K.R., see Taylor, G.A. (183) 421
Harriott, M., see Burns, D.T. (200) 305
Harris, J., see Echols, R.E. (193) 470
Harris, J.A., see Horrocks, J.P. (197) 109
Harsányi, I., see Pauncz, J.K. (195) 251
Hartvig, P., Åhs, U. and Wickström, G.
Determination of propiomazine and its N-demethyl metabolite in plasma by GC with alkali flame ionization detection (183) 229
Hartvig, P., Fagerlund, C. and Gyllenhaal, O.
Electron-capture GC of plasma sulphonylureas after extractive methylation (181) 17
Hartvig, P., see Gyllenhaal, O. (189) 351
Hartwick, R.A., Assenza, S.P. and Brown, P.R.
Identification and quantitation of nucleosides, bases and other UV-absorbing compounds in serum, using reversed-phase HPLC. I. Chromatographic methodology (186) 647
Hartwick, R.A., Krstulovic, A.M. and Brown, P.R.
Identification and quantitation of nucleosides, bases and other UV-absorbing compounds in serum, using reversed-phase HPLC. II. Evaluation of human sera (186) 659
Harvey, D.J.
Dimethoxymethylsilyl ethers: stable silyl derivatives for GC and MS (196) 156
Harvey, O., see Garneau, F.-X. (188) 445
Harzer, K.
Detection of glibornurid in serum by reversed-phase HPLC (in German) (183) 115
Hashimoto, M., see Nobuhara, Y. (190) 251
Hashimoto, T., see Kato, Y. (190) 297, (190) 305, (193) 29, (193) 311, (193) 458
Hashimoto, Y., see Endo, M. (196) 334
Hassan-Ali, S., see Witzgall, H. (198) 70
Hassler, W., see Fitt, L.E. (187) 381
Hasson, K., see Clements, J.A. (189) 272
Hastings, J.R.B., see Walker, J.M. (189) 106
Hata, M., see Kimura, M. (183) 467
Hatch, R.T., see Radjai, M.K. (196) 319
Hatta, Y., see Fukunaga, A. (190) 339
Hautoniemi, L., see Nieminen, A.-L. (181) 11
Hayashi, C., see Yamaguchi, Y. (182) 430
Hayashida, H., see Inayama, S. (194) 85
Hayes, A.W., see Chan, P.K. (194) 387
Hayes, A.W., see Phillips, R.D. (190) 419
Haymond, M.W., Howard, C.P., Miles, J.M. and Gerich, J.E.
Determination of leucine flux in vivo by GC-MS utilizing stable isotopes for trace and internal standard (183) 403
Hazzard, C.Y., see Hoffsommer, J.C. (195) 435
Hearn, M.T.W., Bethell, G.S., Ayers, J.S. and Hancock, W.S.
Application of 1,1'-carbonyldiimidazole-activated agarose for the purification of proteins. II. The use of an activated matrix devoid of additional charged groups for the purification of thyroid proteins (185) 463
Hearn, M.T.W., Grego, B. and Hancock, W.S.
HPLC of amino acids, peptides and proteins. XX. Investigation of the effect of pH and ion-pair formation on the retention of peptides on chemically-bonded hydrocarbonaceous stationary phases (185) 429
Hearn, M.T.W., see Bishop, C.A. (192) 222
Heath, R.R., Burnsed, G.E., Tumlinson, J.H. and Doolittle, R.E.
Separation of a series of positional and geometrical isomers of olefinic aliphatic primary alcohols and acetates by capillary GC (189) 199
Heaulme, M. de, see De Heaulme, M. (181) 254
Hechtman, P., see Hardwick, J. (190) 385
Heemstra, S., see Terweij-Groen, C.P. (181) 385
Heftmann, E., see Hunter, I.R. (198) 363
Heftmann, E., see Lin, J.-T. (190) 169
Heimler, D., see Lepri, L. (195) 65, (195) 187, (195) 339
Heisler, E.G., Siciliano, J., Stinson, E.E., Osman, S.F. and Bills, D.D.
HPLC determination of major mycotoxins produced by *Alternaria* molds (194) 89

Hekman, P., Porskamp, P.A.T.W., Ketelaars, H.C.J. and Van Ginneken, C.A.M.
Rapid HPLC method for the determination of probenecid in biological fluids (182) 252
Hekman, P. and Van Ginneken, C.A.M.
Rapid determination of renal contrast media in biological fluids by means of HPLC (182) 492
Helboe, P., Olsen, O. and Sørensen, H.
Separation of glucosinolates by HPLC (197) 199
Helboe, P., Thomsen, M. and Hansen, S.H.
Improved HPLC method for the comparison of heptaene macrolide antibiotics (189) 249
Heldt, U. and Köser, H.J.K.
Different bases for the GC retention index system (192) 107
Heller, W., Schallies, M. and Schmidt, K.
GLC-MS studies on extracts and distillates from Posidonomia shales (186) 843
Hellgren, L., see Engström, N. (189) 284
Helmich, O. and Hradec, J.
Isolation of 14-methylhexadecanoic acid from wool fat by preparative GLC (193) 153
Henderson, D.E., see Uden, P.C. (196) 403
Henderson, G.L., see Gallaher, C. (181) 490
Hendriks, H. and Bruins, A.P.
Study of three types of essential oil of *Valeriana officinalis* L. s.l by combined GC-negative ion chemical ionization MS (190) 321
Henke, B. and Westerlund, D.
Hydrophobic chromatography and bioanalysis of some polar pyridine derivatives used as antilipolytic agents (187) 189
Henriksen, U.B. and Jelnes, J.E.
Experimental taxonomy of *Biomphalaria* (gastropoda: planorbidae). I. Methods for experimental taxonomic studies on *Biomphalaria* carried out by horizontal starch gel electrophoresis and staining of twelve enzymes (188) 169
Henry, D.R., see Block, J.H. (193) 111
Henson, J.B., Carver, J.R., Wilson, J.B. and Huisman, T.H.J.
Carboxymethyl-cellulose microchromatography for the quantitation of hemoglobin Bart's (γ_4) and its use in the detection of the α-thalassemia conditions (198) 443
Herbut, G., see Kowalczyk, J.S. (196) 11
Hercules, D.M., see Wright, B.W. (199) 355
Herkner, K. and Swoboda, W.
Concentration-dependent changes in Methylene Units in steroid analysis (in German) (197) 232
Hermansson, J.
Separation of steroid glucuronides by reversed-phase LCC (194) 80
Hernández, R., see Sosa, J.M. (189) 169
Herrmann, K., see Galensa, R. (189) 217
Herrmann, K., see Schulz, J.M. (195) 85, (195) 95
Herzel, F.
Determination of 2,6-dichlorobenzonitrile residues in soil and water (in German) (193) 320
Hesselgren, T., see Beck, O. (181) 100
Heyes, W.F., Salmon, J.R. and Marlow, W.
HPLC separation of the N- and S-oxides of fluphenazine and fluphenazine decanoate (194) 416
Heyns, K., see Röper, H. (193) 381
Hibi, K., Ishii, D. and Tsuda, T.
Alumina and support-coated open-tubular columns in open-tubular micro-capillary LC (189) 179
Hignite, C., see Preskorn, S.H. (197) 246
Hikida, K., see Sadanaga, T. (183) 246
Hilbers, H., see Zuidema, J. (182) 445
Hilbers, H.W., see Zuidema, J. (182) 130
Hill, H.H. Jr., see Dupuis, M.D. (195) 211
Hill, H.H. Jr., see Osman, M. (186) 273
Hindle, R., see Sundaram, K.M.S. (194) 100
Hines, H.B., see Middleditch, B.S. (195) 359
Hirai, Y., see Yoza, N. (196) 471
Hiramatsu, K., Nozaki, H. and Arimori, S.
Lipid content of human platelets quantitated by TLC in combination with flame ionization detection (182) 301
Hiramatsu, M., see Ikeda, M. (183) 41
Hirano, S., see Nobuhara, Y. (190) 251
Hirata, Y., Lin, P.T., Novotný, M. and Wightman, R.M.
Small-volume electrochemical detector for microcolumn LC (181) 287
Hirata, Y. and Novotný, M.
Techniques of capillary LC (186) 521
Hirose, N., see Tamegai, T. (193) 483
Hirota, K., see Ikeda, M. (183) 41
Hirschhorn, R., see Ratech, H. (183) 499
Hirschy, L.M., see Marcelin, G. (187) 57
Hisayasu, G.H., see Cohen, J.L. (181) 478
Hiscocks, P.G., see Seamark, D.A. (197) 271
Hjalmarsson, S.-G., see Moberg, U. (181) 147
Hjertén, S., Öfverstedt, L.-G. and Johansson, G.
Free displacement electrophoresis (isotachophoresis): an absolute determination of the Kohlrausch functions and their use in interaction studies (194) 1
Hlavay, J., see Varga-Puchony, Z. (196) 185
Hlavay, J., see Vigh, G. (193) 432
Hobbs, D.C., see Twomey, T.M. (183) 104
Hodžić, I.M., see Ćelap, M.B. (198) 172
Hoehn, E., Kim, S.I., Eskin, N.A.M. and Ismail, F.
Chromogenic reagent for vicine and convicine on thin-layer plates (194) 251
Hoellinger, H., see Scherrmann, J.M. (196) 342
Hoener, B.-A. and Wolff, J.L.
HPLC assay for the metabolites of nitrofurantoin in plasma and urine (182) 246
Hoes, M.J.A., see O'Reilly, W.J. (183) 492
Hoeven, T.A. van der, see Van der Hoeven, T.A. (196) 494
Hoffelt, J., see Scherrmann, J.M. (196) 342

Hoffman, H.T. Jr.
Determination of furfural in spent sulfite liquor by GC (194) 228
Hoffsommer, J.C., Glover, D.J. and Hazzard, C.Y.
Quantitative analysis of polynitrophenols in water in the micro- to nanogram range by reversed-phase ion-pair LC (195) 435
Hofmann, K. and Halász, I.
Mass transfer in ideal and geometrically deformed open tubes. III. Deformed metal and plastic tubes (199) 3
Hokse, H.
Analysis of cyclodextrins by HPLC (189) 98
Holan, Z., Votruba, J. and Vlasáková, V.
New method of chitin determination based on deacetylation and GLC assay of liberated acetic acid (190) 67
Holczinger, L., see Kremmer, T. (191) 287
Holdiness, M.R., Rosen, M.T., Justice, J.B. and Neill, D.B.
GC-MS determination of dopamine in subregions of rat brain (198) 329
Holdren, M., see Osman, M. (186) 273
Hollister, L.E., see Lombrozo, L. (181) 1
Holloway, C.J., Husmann-Holloway, S. and Brunner, G.
New method for the quantitative assay of the enzymatic hydrolysis of UDPglucuronic acid using analytical capillary isotachophoresis (188) 235
Holt, D.W., see Flanagan, R.J. (187) 391
Holtlund, J., see Kristensen, T. (192) 494
Holzer, G., Oró, J., Smith, S.J. and Doctor, V.M.
Separation of monosaccharides as their alditol acetates by capillary column GLC (194) 410
Holzer, G., Oró, J. and Tornabene, T.G.
GC-MS analysis of neutral lipids from methanogenic and thermoacidophilic bacteria (186) 795
Honda, S., Tanimitsu, N. and Kakehi, K.
GC analysis of aliphatic and aromatic aldehydes as trimethylsilylated dithioacetals of 2-mercaptoethanol (194) 191
Hong, J., see Peralta, E. (190) 43
Honigberg, I.L., Stewart, J.T., Clark, T.C. and Davis, D.Y.
Non-extractive fluorometric measurement of *p*-aminosalicylic acid in plasma by ion-pairing techniques and HPLC (181) 266
Hoogmartens, J., see Roets, E. (194) 262
Hooper, W.D., see Ross-Lee, L.M. (183) 175
Hörhold, C., see Atrat, P. (189) 279
Hori, H., see Inayama, S. (194) 85
Horikawa, R., see Miyazaki, S. (181) 177
Horino, M., see Kakita, K. (181) 248
Horký, K., see Tomsová, Z. (200) 221
Horning, E.C., see Barrow, S.E. (181) 219
Horning, M.G. and Lertratanangkoon, K.
HPLC separation of carbamazepine metabolites excreted in rat urine (181) 59
Horning, M.G., see Barrow, S.E. (181) 219
Horrocks, J.P., Harris, J.A., Phillips, C.S.G. and Scott, K.F.
Quantitative aspects of heater displacement chromatography (197) 109
Horváth, C., Melander, W. and Nahum, A.
Measurement of association constants for complexes by reversed-phase HPLC (186) 371
Horváth, C., see Melander, W. (199) 35
Horváth, C., see Melander, W.R. (185) 99, (185) 111, (185) 129
Horváth, C., see Nahum, A. (192) 315
Hostettmann, K., Hostettmann-Kaldas, M. and Sticher, O.
Application of droplet counter-current chromatography to the isolation of natural products (186) 529
Hostettmann-Kaldas, M., see Hostettmann, K. (186) 529
House, D.W., see Pirkle, W.H. (192) 143
Howard, A.G., Pizzie, R.A. and Tickell, S.M.
TLC of S-methylcysteine sulphoxide (kale anaemia factor) in the presence of common amino acids (189) 436
Howard, C.P., see Haymond, M.W. (183) 403
Hradec, J., see Helmich, O. (193) 153
Hrdina, P.D., see Rovei, V. (182) 349
Hsü, C.-Y., see Wu, H.-L. (193) 476
Hsu, D.-S. and Chen, S.S.
Simple anion-exchange chromatography for the determination of adenine nucleotides by using AG MP-1 resin (192) 193
Hsu, D.-S., see Chen, S.S. (198) 500
Hsu, F., see Poole, C.F. (199) 123
Hsu, L.-C., see Wu, H.-L. (193) 476
Huang, C.T.L., Szczepanik-van Leeuwen, P.A. and Nichols, B.L.
New solvent systems for separation of free and conjugated bile acids. III. Separation of bile acid methyl ester acetates (196) 150
Huber, J.F.K., see Jonker, R.J. (186) 285
Hubert, P. and Dellacherie, E.
Use of water-soluble biospecific polymers for the purification of proteins (Review) (184) 325
Hubert, P. and Porath, J.
Metal chelate affinity chromatography. I. Influence of various parameters on the retention of nucleotides and related compounds (198) 247
Hübner, G., see Kramlová, M. (193) 515
Hui, K.-S., Salschutz, M., Davis, B.A. and Lajtha, A.
Separation of alkylaminonaphthylene-sulfonyl peptides and amino acids by HPLC. Methods for measuring melanotropin inhibiting factor breakdown (192) 341
Huibregtse-Minderhout, L., Van der Kerk-Van Hoof, A.C., Wijkens, P., Biessels, H.W.A. and Salemink, C.A.
Glass capillary column GC method for simultaneous quantitative determination of insect juvenile hormones at the picogram level: a comparitive study of various halogenated derivatives (196) 425
Huikeshoven, H., see Zuidema, J. (182) 130

Huisman, T.H.J., see Henson, J.B. (198) 443
Huizing, H.J., De Boer, F. and Malingré, T.M.
Chloranil, a sensitive detection reagent for pyrrolizidine alkaloids on thin-layer chromatograms (195) 407
Hulshoff, A., Neijt, J.P., Smulders, C.F.A., Van Loenen, A.C. and Pinedo, H.M.
Determination of hexamethylmelamine and metabolites in plasma or serum by GLC with a nitrogen-sensitive detector (181) 363
Hulshoff, A., Roseboom, H. and Renema, J.
Improved detectability of barbiturates in HPLC by pre-column labelling and UV detection (186) 535
Hulshoff, A., see Barends, D.M. (182) 201
Hulsman, J., see Mikkers, F. (182) 496
Hundt, H.K.L., Brown, L.W. and Clark, E.C.
Determination of cinnarizine in plasma by HPLC (183) 378
Hundt, H.K.L., Clark, E.C. and Van der Linde, H.C.
TLC method for the quantitative analysis of L-tryptophan in human plasma (182) 110
Hundt, H.K.L., see Van der Meer, M.J. (181) 282
Hunter, C., see Sugden, K. (192) 228
Hunter, I.R., Walden, M.K. and Heftmann, E.
HPLC of *Solanum* and *Veratrum* alkaloids (198) 363
Hunter, I.R., see Lin, J.-T. (190) 169
Hušek, P., see Felt, V. (197) 226
Husmann-Holloway, S., see Holloway, C.J. (188) 235
Hutchinson, I.G., see Minnikin, D.E. (188) 221
Hutte, R.S., see Sievers, R.E. (186) 3
Hwang, D.L., see Lin, K.D. (195) 385
Hylin, J.W., see Ogata, J.N. (189) 425

I

Iizima, N., Fujihara, M. and Nagumo, T.
Application of a sintered silica gel plate to the TLC of carbohydrates (193) 464
Ikeda, M., Kawase, M., Hiramatsu, M., Hirota, K. and Ohmori, S.
Improved GC method of determining diclofenac in plasma (183) 41
Imai, K., see Miyazaki, S. (181) 177
Imamichi, H., see Endo, M. (196) 334
Imazu, M., see Inayama, S. (194) 85
Inayama, S., Hori, H., Shibata, T., Ozawa, Y., Yamagami, K., Imazu, M. and Hayashida, H.
Simple and rapid separation of certain prostaglandins by reversed-phase HPLC (194) 85
Inoue, H., see Uobe, K. (193) 83
Insler, M., see Hanna, S. (200) 277
Ioffe, B.V., Vitenberg, A.G. and Tsibul'skaya, I.A.
Application of head-space analysis to the study of volatile organic impurity concentrates. GC determination of aromatic hydrocarbons in atmospheric air (186) 851
Irgolic, K.J., see Brinckman, F.E. (191) 31
Irwin, W.J., see Williams, K.J. (194) 217
Ishibashi, M., see Fukunaga, A. (190) 339
Ishibashi, T., Kawabata, T. and Tanabe, H.
GC determination of N-nitrosoamino acids (195) 416
Ishida, M., Suyama, K. and Adachi, S.
Background contamination by phthalates commonly encountered in the chromatographic analysis of lipid samples (189) 421
Ishii, D., Tsuda, T. and Takeuchi, T.
Studies of open-tubular micro-capillary LC. IV. Soda-lime glass columns treated with alkaline solution (185) 73
Ishii, D., see Hibi, K. (189) 179
Ishii, D., see Takeuchi, T. (190) 150, (192) 127
Ishikura, S., see Onodera, S. (200) 137
Ishimura, A., see Shiono, S. (193) 243
Ishizuka, T., see Tanaka, K. (190) 77
Ismail, F., see Hoehn, E. (194) 251
Isobe, K., see Onishi, S. (182) 105
Isobe, T., see Shimono, T. (197) 59
Ito, K., see Yoza, N. (196) 471
Ito, M., see Tsukida, K. (192) 395
Ito, S. and Fujita, K.
Detection of thiols on thin-layer chromatograms with 3,5-di-*tert.* -butyl-1,2-benzoquinone-iron(III) chloride (187) 418
Ito, T., see Matsuki, Y. (188) 177
Ito, Y.
New horizontal flow-through coil planet centrifuge for counter-current chromatography. I. Principle of design and analysis of acceleration (188) 33
Ito, Y.
New horizontal flow-through coil planet centrifuge for counter-current chromatography. II. The apparatus and its partition capabilities (188) 43
Ito, Y.
Toroidal coil planet centrifuge for counter-current chromatography (192) 75
Ito, Y.
Preparative counter-current chromatography with a slowly rotating glass coil (196) 295
Ito, Y. and Putterman, G.J.
New horizontal flow-through coil planet centrifuge for counter-current chromatography. III. Separation and purification of dinitrophenyl amino acids and peptides (193) 37
Itoh, S., see Onishi, S. (182) 105
Itokawa, Y., see Kimura, M. (188) 417
Iverson, W.P., see Brinckman, F.E. (191) 31
Ivory, C.F.
Continuous flow electrophoresis. The crescent phenomena revisited. I. Isothermal effects (195) 165

Ivory, C.F. and Bratzler, R.L.
Error incurred in gel permeation chromatography by using the elution peak volume in lieu of the elution mean volumes in the calculation of K_{av} (198) 354
Iwai, K., see Suzuki, T. (198) 217
Iwakawa, J., Kobatake, H., Suzuki, I. and Kushida, H.
Fluorodensitometric microdetermination of reducing sugars on thin-layer chromatograms (193) 333
Iwasaki, H., Shimada, K. and Tarutani, T.
Gel chromatographic study of the polymerization of silicic acid in acid solutions (198) 429
Iwu, M.M., see Court, W.E. (187) 199

J

Jack, D.B., Dean, S. and Kendall, M.J.
Detection of some antihypertensive drugs and their metabolites in urine by TLC. Five commonly used beta blockers and hydralazine (187) 277
Jack, D.B., Dean, S., Kendall, M.J. and Laugher, S.
Detection of some antihypertensive drugs and their metabolites in urine by TLC. II. A further five beta blockers and dihydralazine (196) 189
Jack, D.B., see Quarterman, C.P. (183) 92
Jack, D.B., see Schweizer, A. (195) 421
Jackman, G., see Bangah, M. (183) 255
Jackman, G.P., Carson, V.J., Bobik, A. and Skews, H.
Simple and sensitive procedure for the assay of serotonin and catecholamines in brain by HPLC using fluorescence detection (182) 277
Jacob, K., see Meyer, H.D. (199) 339
Jacob, K., see Vogt, W. (186) 197, (199) 191
Jacquet, M.A., see Dodeur, M. (195) 197
Jahangir, L.M. and Samuelson, O.
Chromatography in aqueous solution on styrene-divinylbenzene resins (193) 197
Jakobsen, P., see Treppendahl, S. (189) 276
Jakovljevic, I.M. and Bishara, R.H.
Resolution of ethylphenols following *in situ* reaction on a thin-layer plate with Fast Blue B Salt (192) 425
James, G.T., Thach, A.B., Connole, E., Austin, J.H. and Rinehart, R.
Nucleic acid bases and derivatives: detection by Dns derivatization TLC (195) 287
Janado, M., Yano, Y., Nakamori, H. and Nishida, T.
Anomalous partition behaviour of sodium dodecyl sulphate monomer in Sephadex gels with high dextran concentrations. Possible role of water of hydration in the gel phase (193) 345
Janado, M., see Yano, Y. (200) 125
Janák, J., see Boček, P. (191) 271, (195) 303
Janák, J., see Gebauer, P. (199) 81
Janák, J., see Novák, J. (199) 209
Janák, J., see Soják, L. (191) 199, (195) 43
Janča, J.
Concentration effects in gel permeation chromatography. VI. Non-Newtonian behaviour of polymer solutions (187) 21
Janča, J., see Giddings, J.C. (186) 37
Jandera, P. and Churáček, J.
Gradient elution in LC. XI. Influence of the adjustable gradient parameters on the chromatographic behaviour of sample compounds (192) 1
Jandera, P. and Churáček, J.
Gradient elution in LC. XII. Optimization of conditions for gradient elution (192) 19
Jandera, P. and Churáček, J.
Reversed-phase LC of aromatic sulphonic acids and other strongly polar compounds without addition of an ion-pairing counter-ion (197) 181
Jandera, P., Churáček, J. and Svoboda, L.
Gradient elution in LC. XIII. Instrumental errors in gradient elution chromatography (192) 37
Janicki, C.A., see Ko, C.Y. (190) 429
Janini, G.M., Manning, W.B., Zielinski, W.L. Jr and Muschik, G.M.
GLC separation of bile acids and steroids on a nematic liquid crystal (193) 444
Janini, G.M., see Zielinski, W.L. Jr. (186) 237
Janjić, T.J., see Ćelap, M.B. (196) 59, (198) 172
János, É., Balla, J., Tyihák, E. and Gáborjányi, R.
GLC analysis of dimedone derivatives of formaldehyde and other aliphatic aldehydes on capillary columns (191) 239
Jánosi, A., see Bartik, T. (197) 92
Jansen, M.J.T., see Lankelma, J. (182) 35
Jansen, P.L.M. and Tangerman, A.
Separation and characterization of bilirubin conjugates by HPLC (182) 100
Jansson, L., Nilsson, B. and Lindgren, R.
Quantitation of serum tocopherols by HPLC with fluorescence detection (181) 242
Jansta, J., see Smolková, E. (191) 61
Jaouni, T.M., Leon, M.B., Rosing, D.R. and Fales, H.M.
Analysis of verapamil in plasma by LC (182) 473
Jaramillo, L.F. and Driscoll, J.N.
Analysis of therapeutic and commonly abused drugs in serum and urine by GLC using a photoionization detector (186) 637
Jardine, I. and Weidner, M.M.
Approach to the quantitative analysis of nucleotides by GC-MS (182) 395
Jardine, I., see Strife, R.J. (182) 211
Jaroniec, J.A., see Jaroniec, M. (188) 27

Jaroniec, M., Różyło, J.K., Jaroniec, J.A. and Ościk-Mendyk, B.
Liquid adsorption chromatography with mixed mobile phases. IV. A new equation for the capacity ratio involving the solute-solvent interactions (188) 27
Jefcoate, C.R., see Hanukoglu, I. (190) 256
Jefferies, T.M., see Riley, C.M. (185) 197
Jelínková, M., see Kuchař, M. (195) 329
Jellema, R., Elema, E.T. and Malingré, T.M.
Optical brighteners as TLC detection reagents for glycoalkaloids and steroid alkaloids in *Solanum* species. II. Blankophor BA 267%, BBU neu and KU, and Tinopal CBS-X and 5 BMS-X (189) 406
Jelnes, J.E., see Henriksen, U.B. (188) 169
Jender, H.G., see Schwenn, J.D. (193) 285
Jennings, W., Settlage, J.A., Miller, R.J. and Raabe, O.G.
New approach to chromatographic optimization in glass capillary GC (186) 189
Jensen, J.C. and Faiman, M.D.
Determination of disulfiram and metabolites from biological fluids by HPLC (181) 407
Jerauld, R.S., see Wheeler, L.A. (183) 357
Jernejčič, M.
Study of the interactions between substituted phenols and alcohols and dibenzyl sulphoxide by means of GC (198) 495
Jesdinsky, D., see Nau, H. (182) 71
Jewett, K.L., see Brinckman, F.E. (191) 31
Johansson, G., see Hjertén, S. (194) 1
Johansson, L., see Gyllenhaal, O. (190) 347
Johansson, L., see Poole, C.F. (194) 365
Johns, E.W., see Mathew, C.G.P. (198) 80
Johns, E.W., see Smith, B.J. (200) 200
Johns, E.W., see Walker, J.M. (189) 106
Johnson, K.J., see Robinson, J.L. (189) 145
Johnson, P.E. and Evans, G.W.
Binding of zinc and copper to some gel filtration media (188) 405
Johnson, R.C., Dawson, R.M.C. and Freinkel, N.
Paper ionophoresis of pancreatic islet phosphate esters in a magnesium-containing buffer (187) 235
Johnston, M.A., Smith, W.J., Kennedy, J.M., Lea, A.R. and Hailey, D.M.
Reversed-phase HPLC analysis of cinchona alkaloids in pharmaceuticals (189) 241
Jones, A.D., see Smith, E.C. (188) 205
Jones, C.E.R. and Vanderborgh, N.E.
Elucidation of geomatrices by laser pyrolysis-GC and pyrolysis-MS (186) 831
Jones, G.J.L., see Kuksis, A. (182) 1
Jones, G.R., see Coutts, R.T. (190) 53
Jong, A.W.J. de, see De Jong, A.W.J. (193) 181
Jong, G.J. de, see De Jong, G.J. (183) 203
Jong, W. de, see De Jong, W. (183) 303
Jonker, R.J., Poppe, H. and Huber, J.F.K.
In situ loading and reloading of GC columns with stationary liquids (186) 285
Jonkman, J.H.G., see Greving, J.E. (186) 683
Jonsson, A. and Berg, S.
Determination of 1,2-dibromoethane, 1,2-dichloroethane and benzene in ambient air using porous polymer traps and GC-MS analysis with selected ion monitoring (190) 97
Jordan, P.M., see Nordlöv, H. (190) 221
Jorgenson, J.W., see Smith, S.L. (187) 111
Jost, W., Unger, K.K., Lipecky, R. and Gassen, H.G.
Application of a weakly basic dimethylamino-modified silica ion exchanger to the separation of oligonucleotides (185) 403
Joy, K.W., Shay, C. and McLimont, M.J.
Effect of some organic buffers on the estimation of aspartic acid and resolution in amino acid analysis (194) 76
Jozefczak, C., see Bernard, N. (183) 99
Jukofsky, D., Verebey, K. and Mulé, S.J.
Qualitative differentiation between cocaine, lidocaine and cocaine-lidocaine mixtures ("rock-cocaine") using TLC (198) 534
Julák, J., Tureček, F. and Miková, Z.
Identification of characteristic branched-chain fatty acids of *Mycobacterium kansasii* and *gordonae* by GC-MS (190) 183
Jurand, J., see Knox, J.H. (186) 763
Jurenitsch, J. and Kampelmühler, I.
Rapid determination of nonylic acid vanillylamide and other capsaicinoids in Capsicum fruits and extracts by means of Ag^+-complexation HPLC (in German) (193) 101
Jurenitsch, J. and Leinmüller, R.
Quantification of nonylic acid vanillylamide and other capsaicinoids in the pungent principle of Capsicum fruits and preparations by GLC on glass capillary columns (189) 389
Jurenitsch, J., Lumper, G. and Bachmann, H.
Determination of free short-chain monocarboxylic acids from naturally occurring esters by GLSC on micropacked columns (in German) (189) 43
Just, D.E., see Fitt, L.E. (187) 381
Justice, J.B., see Holdiness, M.R. (198) 329

K

Kaaden, A. van der, see Van der Kaaden, A. (198) 526
Kaay, M. van der, see Van der Kaay, M. (181) 227
Kahle, V. and Tesařík, K.
Separation of saccharides and their anomers by HPLC (191) 121
Kakehi, K., see Honda, S. (194) 191

Kakita, K.
Release of insulin analogues in man, stimulated with glucose. In vitro and in vivo studies (182) 419
Kakita, K., Horino, M., Oyama, H., Tenku, A., Matsumura, S. and Nishida, S.
Gel chromatographic separation of insulin analogues in human serum (181) 248
Kakuno, K., see Yamamoto, S. (194) 399
Kálal, J., see Brynda, E. (196) 39
Kálal, J., see Lukás, J. (194) 297
Kalász, H., see Mincsovics, E. (191) 293
Kaljurand, M. and Küllik, E.
Comparison between conventional and cross-correlation GC for evolved gas analysis (186) 145
Kalmikova, T.A., see Lazaris, A.Y. (198) 337
Kamiński, M., see Gazda, K. (191) 9
Kampelmühler, I., see Jurenitsch, J. (193) 101
Kanaske, K., see Lombrozo, L. (181) 1
Kangas, L., see Nieminen, A.-L. (181) 11
Kaniansky, D., Madajová, V., Zelenský, I. and Stankoviansky, S.
Role of the charge number of the counter-ionic constituent in the separation of anions by isotachophoresis (194) 11
Kaniewska, T. and Wejman, W.
GC determination of bromo and fluoro derivatives of benzodiazepine in human body fluids (182) 81
Kaniwa, M.-A., see Nakamura, A. (196) 133
Kapila, S., Vogt, C.A. and Aue, W.A.
Reaction and response in electron-capture detectors (196) 397
Kapila, S., Vogt, C.R. and Aue, W.A.
Effects of dimension and density in electron capture detectors (195) 17
Kapila, S., see Aue, W.A. (188) 1
Kaplowitz, N., see Reeve, J. (194) 424
Karácsony, E.M., see Szabó, A. (193) 500
Karakaya, A. and Carter, D.E.
HPLC of glucuronide and sulfate conjugates using ion-pair chromatography (195) 431
Karasek, F.W., see Clement, R.E. (189) 53, (190) 136
Karasek, F.W., see Dickson, L.C. (190) 311
Karasek, F.W., see Eiceman, G.A. (200) 115
Karger, B.L., see Lindner, W. (185) 323
Karger, B.L., see McCormick, R.M. (199) 259
Karino, I., see Shiono, S. (193) 243
Karmen, A., Lam, S., Cummings, G. and Chow, F.
Specific detection in GLC effluents with on-line HPLC optical detectors (199) 113
Karmen, A., see Lam, S. (199) 295
Karobath, M., see Lapin, A. (193) 95
Karski, S., see Paryjczak, T. (188) 254
Kaspar, H.F. and Tiedje, J.M.
Response of electron-capture detector to hydrogen, oxygen, nitrogen, carbon dioxide, nitric oxide and nitrous oxide (193) 142
Kasuya, Y., see Baba, S. (196) 144
Kataoka, H., see Yamamoto, S. (194) 399
Kates, R.E., see Harapat, S.R. (181) 484
Kathuria, L., see Chawla, H.M. (188) 289
Kato, T., see Kojima, K. (189) 233
Kato, Y., Komiya, K., Sasaki, H. and Hashimoto, T.
Separation range and separation efficiency in high-speed gel filtration on TSK-GEL SW columns (190) 297
Kato, Y., Komiya, K., Sasaki, H. and Hashimoto, T.
High-speed gel filtration of proteins in sodium dodecyl sulphate aqueous solution on TSK-GEL SW type (193) 29
Kato, Y., Komiya, K., Sasaki, H. and Hashimoto, T.
Comparison of TSK-GEL PW type and SW type in high-speed aqueous gel-permeation chromatography (193) 311
Kato, Y., Komiya, K., Sasaki, H. and Hashimoto, T.
High-speed gel filtration of proteins in 6 *M* guanidine hydrochloride on TSK-GEL SW columns (193) 458
Kato, Y., Komiya, K., Sawada, Y., Sasaki, H. and Hashimoto, T.
Purification of enzymes by high-speed gel filtration on TSK-GEL SW columns (190) 305
Katoh, K. and Miyazaki, H.
Assay method for urokinase activity by capillary-tube isotachophoresis using a synthetic substrate (188) 383
Katoh, K., see Someno, T. (188) 185
Katz, E., see Ogan, K. (188) 115
Kawabata, T., see Ishibashi, T. (195) 416
Kawabe, K., see Tamegai, T. (193) 483
Kawada, T., see Suzuki, T. (198) 217
Kawade, N., see Onishi, S. (182) 105
Kawai, S., see Kobayashi, K. (187) 413
Kawai, S., see Takayama, Y. (197) 240
Kawalek, J.C., see Shaikh, B. (195) 392
Kawano, Y., Yanagihara, K., Miyamoto, T. and Yamamoto, I.
Examination of the conversion products of pyrethrins and allethrin formulations exposed to sunlight by GC and MS (198) 317
Kawase, M., see Ikeda, M. (183) 41
Keehnen, P.D.M., see Billiet, H.A.H. (185) 515
Keller, T., see Bayer, E. (186) 497
Kempf, J., see Ochoa, J.L. (188) 257
Kemula, W., see Kutner, W. (191) 47
Kendall, M.J., see Jack, D.B. (187) 277, (196) 189
Kendall, M.J., see Quarterman, C.P. (183) 92
Kendall, M.J., see Schweizer, A. (195) 421
Kennedy, J.F., Barnes, J.A. and Matthews, J.B.
Model studies of the reactions and non-specific adsorptive properties of cyanogen bromide-activated macroporous agarose (196) 379
Kennedy, J.M., see Johnston, M.A. (189) 241
Kennedy, J.M., see Lea, A.R. (198) 41
Kerk-Van Hoof, A.C. van der, see Van der Kerk-Van Hoof, A.C. (196) 425
Kerns, E., see Simmonds, P.G. (186) 785
Kerremans, L., see Vanderhaeghe, H. (193) 119

Kessaissia, Z., Papirer, E. and Donnet, J.-B.
GC study of grafted alkyl chains used as stationary phases (196) 481
Ketelaars, H.C.J., see Hekman, P. (182) 252
Ketelaars, H.C.J., see Van Loenhout, J.W.A. (182) 487
Khripach, V.A., see Volkov, S.M. (190) 445
Kiguchi, S., see Ubuka, T. (188) 442
Kim, S.H., see Grimsrud, E.P. (197) 51
Kim, S.I., see Hoehn, E. (194) 251
Kimura, M., Fujita, T., Nishida, S. and Itokawa, Y.
Differential fluorimetric determination of picogram levels of thiamine, thiamine monophosphate, diphosphate and triphosphate using HPLC (188) 417
Kimura, M., Kobayashi, K., Hata, M., Matsuoka, A., Kitamura, H. and Kimura, Y.
Determination of gliclazide in human serum by HPLC using an anion-exchange resin (183) 467
Kimura, Y., see Kimura, M. (183) 467
King, L.J., see Williams, P.L. (186) 595
Kinuta, M., see Ubuka, T. (188) 442
Kirby, B.D., see Wheeler, L.A. (183) 357
Kirkland, J.J.
Size-exclusion HPLC of inorganic colloids (185) 273
Kirkland, J.J., see Glajch, J.L. (199) 57
Kirschbaum, J., Gibbs, V., Foley, J. and Perlman, S.
HPLC of the oral anti-asthmatic agent pirquinozol and its active metabolite (200) 282
Kirschbaum, J., Poet, R., Bush, K. and Petrie, G.
HPLC of the topical anti-inflammatory steroid halcinonide (190) 481
Kirson, I., see Tishbee, A. (195) 425
Kiselev, A.V., Aratskova, A.A., Gvozdovitch, T.N. and Yashin, Y.I.
Retention behaviour of *o*-, *m*- and *p*-isomers of benzene derivatives on a silica gel hydroxylated surface in LC (195) 205
Kiselev, A.V., see Gavrilova, T.B. (192) 323
Kitamura, H., see Kimura, M. (183) 467
Klawiter, J., see Gazda, K. (191) 9
Kleijn, E. van der, see Van der Kleijn, E. (182) 35
Klein, A.E., see Davis, P.J. (188) 280
Klein, F.K., see Maybaum, J. (188) 149
Klein, L., see Chandrarajan, J. (182) 94
Klemm, D. and Klemm, E.
TLC of nitrobenzylarenes (in German) (188) 448
Klemm, E., see Klemm, D. (188) 448
Klemm, L.H., Shabtai, J. and Bodily, K.C.
GC of some alkenyl- and cycloalkenylnaphthalenes on Bentone 34, silicone fluid DC 550, and Apiezon L phases (198) 1
Kleyn, E. van der, see Van der Kleyn, E. (183) 492
Kneczke, M.
Determination of pilocarpine, physostigmine, its degradation product rubreserine and preservatives by HPLC (198) 529
Knedel, M., see Meyer, H.D. (199) 339
Knight, B.I. and Skellern, G.G.
Measurement of the formation of paracetamol and *p*-nitrophenol glucuronides *in vitro*, by ion-pair HPLC (192) 247
Knight, P.W., see Rapaka, R.S. (196) 512
Knox, J.H. and Gilbert, M.T.
Kinetic optimization of straight open-tubular LC (186) 405
Knox, J.H. and Jurand, J.
Mechanism of reversed-phase separation of tetracyclines by HPLC (186) 763
Knox, J.H. and McLennan, F.
Band dispersion in high-performance exclusion chromatography (185) 289
Knudsen, P.J., Eriksen, P.B., Fenger, M. and Florentz, K.
HPLC of hyaluronic acid and oligosaccharides produced by bovine testes hyaluronidase (187) 373
Knuutinen, J. and Paasivirta, J.
TLC of chlorinated guaiacols (194) 55
Ko, C.Y. and Janicki, C.A.
Purity determination of zomepirac sodium dihydrate by HPLC (190) 429
Kobatake, H., see Iwakawa, J. (193) 333
Kobayashi, K., Tanaka, M. and Kawai, S.
GC determination of low-molecular-weight carbonyl compounds in aqueous solution as their O-(2,3,4,5,6-pentafluorobenzyl) oximes (187) 413
Kobayashi, K., see Kimura, M. (183) 467
Kobayashi, T.
GLC separation of geometric isomers of unsaturated fatty acid methyl esters using a glass capillary column (194) 404
Kocourek, J., see Turková, R. (192) 408
Kodama, H., Yamamoto, M. and Sasaki, K.
Isotachophoretic analysis of some sulfur-containing amino acids in human urine (183) 226
Kodama, H., see Mikasa, H. (190) 501
Kojima, K., Hama, T., Kato, T. and Nagatsu, T.
Rapid chromatographic purification of dipeptidyl peptidase IV in human submaxillary gland (189) 233
Kojima, S., see Nakamura, A. (196) 133
Kolarz, B.N., see Wojaczyńska, M. (196) 75
Kolman, Z., see Michalec, Č. (190) 498
Kolomiets, L.N., see Berezkin, V.G. (191) 95
Komers, R., see Rosenbaum, M. (191) 157
Komiya, K., see Kato, Y. (190) 297, (190) 305, (193) 29, (193) 311, (193) 458
Kondo, K.
Analytical studies of maridomycin. III. Spectrophotometric determination of maridomycins and 9-propionylmaridomycins as iodine complexes after TLC (190) 493
Konecki, J., see Gabryś, J. (182) 147

Kong, R.C., Sachok, B. and Deming, S.N.
Combined effects of pH and surface-active-ion concentration in reversed-phase LC (199) 307
Kong, R.C., see Sachok, B. (199) 317
König, W.A. and Benecke, I.
GC separation of chiral 2-hydroxy acids and 2-alkyl-substituted carboxylic acids (195) 292
König, W.A. and Sievers, S.
Structural requirements for enantioselectivity in GC of chiral α-hydroxy acids (200) 189
Konijnendijk, A.P., see Deelder, R.S. (185) 241
Konishi, M., see Agoh, T. (182) 171
Kono, A., Hara, Y., Eguchi, S., Tanaka, M. and Matsushima, Y.
Determination of two new metabolites of 1-hexylcarbamoyl-5-fluorouracil in biomedical specimens by HPLC (182) 125
Korol, A.N., Belokleytseva, G.M. and Filonenko, G.V.
Support treated with Carbowax for a standard, non-polar packing in GLC (194) 145
Korol, A.N., see Mikhaylenko, V.P. (191) 231
Korolev, A.A., see Berezkin, V.G. (191) 95
Korte, K., see Gips, H. (193) 322
Köser, H.J.K., see Heldt, U. (192) 107
Kosíková, E., see Franc, J. (187) 462
Kostenbauder, H.B., see Oseekey, K.B. (182) 459
Kouřilová, D., see Krejčí, M. (191) 3
Kovachev, G.L., see Milkova, T.S. (194) 429
Kováts, E. sz., see Fritz, D.F. (186) 63
Kováts, E. sz., see Riedo, F. (186) 47
Kowalczyk, J.S. and Herbut, G.
Influence of temperature on separation processes in adsorption LC systems (196) 11
Kowalczyk, J.S., see Gazda, K. (191) 9
Kozuharov, S.
Coating support-coated open tubular capillary columns with Silar 10C or Alltech CS-10 and Silica T40 for separation of isomers of fatty acid methyl esters (198) 153
Kraak, J.C., see De Jong, A.W.J. (193) 181
Kraak, J.C., see Gast, C.H. (185) 549
Kraak, J.C., see Terweij-Groen, C.P. (181) 385
Kralik, D.C., see Egri, L. (198) 85
Králík, O., see Rittich, B. (197) 43
Kramer, J.K.G., Fouchard, R.C. and Farnworth, E.R.
Effect of solvents on the resolution of neutral lipids on chromarods (198) 279
Kramer, P.A., see Antal, E. (183) 149
Kramer, W.G., see Chang, S.-L. (181) 286
Kraml, J., see Kramlová, M. (193) 515
Kraml, J., see Přistoupil, T.I. (196) 142
Kramlová, M., Přistoupil, T.I., Ulrych, S., Fričová, V., Kraml, J. and Hübner, G.
Electrofocusing of stroma-free hemolyzates of human erythrocytes; comparison of fresh, stored and chemically modified samples (193) 515
Kramlová, M., see Přistoupil, T.I. (196) 142
Kraus, G., see Seifert, K. (192) 97
Krause, R.T.
Resolution, sensitivity and selectivity of a HPLC post-column fluorometric labeling technique for determination of carbamate insecticides (185) 615
Krause, W.
Determination of plasma mepindolol levels by HPLC and electrochemical detection (181) 67
Krejčí, M., Kouřilová, D., Vespalec, R. and Šlais, K.
Measurement of exclusion volumes of packed columns by means of electrokinetic detection (191) 3
Krejčí, M., Tesařík, K. and Pajurek, J.
Open tubular columns in LC (191) 17
Kremmer, T. and Holczinger, L.
Investigation of vinca alkaloid-plasma membrane interactions by detergent gel chromatography (191) 287
Kresse, M., Schley, J. and Müller-Oerlinghausen, B.
Reliable routine method for determination of perazine in serum by TLC with an internal standard (183) 475
Krick, T.P., see Dimenna, G.P. (192) 474
Krick, T.P., see Rao, G.H.R. (196) 506
Kricka, L.J., see Clark, P.M.S. (181) 347
Krien, P., see Arpino, P.J. (185) 529
Kristensen, M., see Angelo, H.R. (183) 159
Kristensen, T. and Holtlund, J.
Chromatography of chromatin proteins on Cibacron Blue F3G-A-agarose (192) 494
Kříž, J., see Kuraš, M. (191) 319
Kříž, J., see Punčochářová, J. (191) 81
Kříž, J., see Vodička, L. (198) 457, (200) 238
Kropinski, A.M., see Ryan, E.A. (195) 127
Krstulovic, A.M., Zakaria, M., Lohse, K. and Bertani-Dziedzic, L.
Diagnosis of neural crest tumors by reversed-phase HPLC determination of urinary catcholamine metabolites (186) 733
Krstulovic, A.M., see Hartwick, R.A. (186) 659
Krupčík, J., Leclercq, P.A., Garaj, J. and Šimová, A.
Analysis of alkylated mixtures of polychlorinated biphenyls by capillary GC-MS (191) 207
Krupčík, J., see Soják, L. (190) 283, (191) 199, (195) 43
Krýsl, S., see Smolková-Keulemansová, E. (184) 347
Ku, D.N., see Churchill, F.C. II (189) 375
Kubín, M., see Vozka, S. (193) 128
Kubinski, H.
Fractionation of ^{32}P-labeled animal RNA on Sephadex-Sepharose columns (197) 236
Kubo, I., see Ganjian, I. (200) 250
Kubota, T., see Ganjian, I. (200) 250
Kucera, P.
Design and use of short microbore columns in LC (198) 93
Kucera, P., see Scott, R.P.W. (185) 27, (186) 475

Kuchař, M., Rejholec, V., Brůnová, B. and Jelínková, M.
R_M values from reversed-phase TLC as parameters of lipophilicity in quantitative structure-activity relationships in four series of arylaliphatic acids (195) 329
Kucsman, Á., see Szókán, G. (198) 207
Kuehn, G.D., see Maestas, R.R. (189) 225
Kuga, S.
New cellulose gel for chromatography (195) 221
Kuhlenkamp, J., see Reeve, J. (194) 424
Kuksis, A., Myher, J.J., Geher, K., Shaikh, N.A., Breckenridge, W.C., Jones, G.J.L. and Little, J.A.
Comparative determination of plasma phospholipids by automated GLC and manual colorimetric phosphorus methods (182) 1
Kullberg, M.P., Dorrbecker, B., Lennon, J., Rowe, E. and Edelson, J.
HPLC analysis of amrinone and its N-acetyl derivative in plasma. Pharmacokinetics of amrinone in the dog (187) 264
Küllik, E., see Kaljurand, M. (186) 145
Kumanotani, J., see Oshima, R. (192) 452
Kumanotani, J., see Yamauchi, Y. (198) 49
Kumps, A. and Mardens, Y.
Improved GLC method for the simultaneous determination of phenobarbital, phenytoin, carbamazepine and primidone in biological fluids (182) 116
Kuniak, Ľ., see Mislovičová, D. (194) 95
Kuo, K.C., see Agris, P.F. (194) 205
Kuo, K.C., see Gehrke, C.W. (188) 129
Kuraš, M., Kříž, J., Tříska, J. and Vodička, L.
Preparative separation of C_9 and C_{10} monoaromatic hydrocarbon mixtures by HPLC on silica gel (191) 319
Kurihara, H., see Ujimoto, K. (190) 161
Kushida, H., see Iwakawa, J. (193) 333
Kuśmierz, J., see Błotny, G. (193) 61
Kutner, W., Dębowski, J. and Kemula, W.
Polarographic detection for HPLC using a flow-through detector (191) 47
Kuwahara, A., see Nakamura, K. (192) 351, (196) 85
Kvernadze, T.K., see Andronikashvili, T.G. (191) 323
Kylberg-Hanssen, K., see Ervik, M. (182) 341

L

Laar, G.L.M. van de, see Van de Laar, G.L.M. (183) 459
Labský, J., see Čeřovský, V. (194) 175
Labský, J., see Tichá, M. (194) 183
Lachman, L., see Hanna, S. (200) 277
Ladisch, M.R., see Beck, K.R. (190) 226
Lafont, P. and Debeaupuis, J.P.
Photodensitometric determination of "PR-toxin", metabolite of *Penicillium roqueforti* (in French) (198) 481
Lafosse, M. and Dreux, M.
Study of mixed solvents: squalane-stearonitrile. I. Comparison of results by GC and surface tensions (in French) (188) 315
Lafosse, M. and Dreux, M.
GLC determination of equilibrium constants of interactions between solutes and free nitrile groups (in French) (193) 9
Laganà, A., Petronio, B.M. and Rotatori, M.
Concentration and determination of polycyclic aromatic hydrocarbons in aqueous samples on graphitized carbon black (198) 143
Lagemaat, D. van de, see Van de Lagemaat, D. (197) 263
Lagerström, P.-O., see Ervik, M. (182) 341
Lajtha, A., see Hui, K.-S. (192) 341
Lajunen, K., Purokoski, S. and Pitkänen, E.
Qualitative TLC separation of 1,5-anhydroglucitol in the presence of other carbohydrates on silica gel impregnated with borate buffer (187) 455
Lake, B.G., see Walters, D.G. (196) 501
Laker, M.F.
Estimation of neutral sugars and sugar alcohols in biological fluids by GLC (Review) (184) 457
Lam, J., see Picman, A.K. (189) 187
Lam, S., Chow, F. and Karmen, A.
Reversed-phase HPLC resolution of D- and L-Dns-amino acids by mixed chelate complexation (199) 295
Lam, S., see Karmen, A. (199) 113
Lamb, J.H., see Bailey, E. (200) 145
Lambert, P.W. and Roos, B.A.
Paired-ion reversed-phase HPLC of human and rat calcitonin (198) 293
Lammens, J., see Verzele, M. (186) 435
Lamparczyk, H., see Grzybowski, J. (196) 217
Lanbeck, K., Lindström, B. and Wibell, L.
HPLC determination of indoprofen in plasma and urine (182) 262
Lanbeck, K., see Floberg, S. (182) 63
Lankelma, J., Van der Kleijn, E. and Jansen, M.J.T.
Determination of 5-methyltetrahydrofolic acid in plasma and spinal fluid by HPLC, using on-column concentration and electrochemical detection (182) 35
Lankmayr, E.P., Budna, K.W. and Nachtmann, F.
Separation of enantiomeric iodinated thyronines by LC of diastereomers (198) 471
Lanouette, M. and Pike, R.K.
Analysis of aminocarb formulations by HPLC (190) 208
Laperashvili, L.Y., see Andronikashvili, T.G. (191) 323

Lapin, A. and Karobath, M.
Simple and rapid mass fragmentographic method for the determination of glycine in brain tissue (193) 95
Larivière, N., see Seidah, N.G. (193) 291
Larsson, B., see Gawell, B.-M. (198) 198
Larsson, L., Mårdh, P.-A., Odham, G. and Westerdahl, G.
Detection of tuberculostearic acid in biological specimens by means of glass capillary GC-electron and chemical ionization MS, utilizing selected ion monitoring (182) 402
Laub, R.J., see Al-Thamir, W.K. (188) 79
Laudat, M.H., see Pileire, B. (182) 269
Laugher, S., see Jack, D.B. (196) 189
Lauren, D.R. and Fairclough, R.J.
Separation and purification of metabolites of the mycotoxin, sporidesmin, by HPLC (200) 288
Lawrence, A.H. and Ducharme, D.
Simple rapid TLC method for the detection of optical brighteners in polymers (194) 434
Lawrence, A.L., see Middleditch, B.S. (195) 359
Lawrence, J.F., Brinkman, U.A.T. and Frei, R.W.
Continuous post-column ion-pair extraction detection of some basic organic compounds in normal-phase chromatography (185) 473
Lawrence, J.F., Panopio, L.G. and McLeod, H.A.
Direct analysis of the herbicide barban in wheat products by reversed-phase LC (195) 113
Lazaris, A.Y., Schmujlovich, S.M. and Kalmikova, T.A.
Determination of residual vinyl chloride in poly(vinyl chloride) resins (198) 337
Lea, A.G.H.
Reversed-phase gradient HPLC of procyanidins and their oxidation products in ciders and wines, optimised by Snyder's procedures (194) 62
Lea, A.R., Kennedy, J.M. and Low, G.K.-C.
Analysis of hydrocortisone acetate ointments and creams by HPLC (198) 41
Lea, A.R., see Johnston, M.A. (189) 241
Leach, C.S., see Di Ferrante, D.T. (187) 271
Leander, K., see Dahlström, H. (183) 511
Leasure, J.B., see Grob, R.L. (197) 129
Leclaire, R., see Besner, J.-G. (183) 346
Leclercq, P.A., see Krupčík, J. (191) 207
Leclercq, P.A., see Soják, L. (191) 187
Leclercq, P.A., see Uralets, V.P. (194) 135
Lederer, M.
Simple and fast separation of the iodotyrosines by TLC (194) 270
Ledvinová-Reinišová, J., see Michalec, Č. (190) 498
Lee, C.H., see Mauro, D. (187) 421
Lee, C.Y., see Wissemann, K.W. (192) 232
Lee, G.J.-L. and Tieckelmann, H.
HPLC separation of unsaturated disaccharides derived from heparan sulfate and heparin (195) 402
Lee, M.L. and Wright, B.W.
Preparation of glass capillary columns for GC (Review) (184) 235
Lee, M.L., see Merli, F. (199) 371
Lee, M.L., see Wright, B.W. (199) 355
Lee, T.D., see Ramsey, J.D. (184) 185
Leenheer, A.P. de, see De Leenheer, A.P. (183) 78, (186) 749, (194) 305, (195) 35
Lefevere, M.F., De Leenheer, A.P. and Claeys, A.E.
HPLC assay of vitamin K in human serum (186) 749
Leibowitz, B.J., see Beck, K.R. (190) 226
Leigh, C.P.H. and Cashion, P.J.
Rapid separation of nucleoside mono-, di- and triphosphates on ion-exclusion/exchange columns (192) 490
Leinmüller, R., see Jurenitsch, J. (189) 389
Lenda, K. and Svenneby, G.
Rapid HPLC determination of amino acids in synaptosomal extracts (198) 516
Lennon, J., see Kullberg, M.P. (187) 264
Leon, M.B., see Jaouni, T.M. (182) 473
Leonard, K., see Preskorn, S.H. (197) 246
LePage, J.N., see Lindner, W. (185) 323
Lepri, L., Desideri, P.G. and Heimler, D.
Reversed-phase and soap TLC of amino acids (195) 65
Lepri, L., Desideri, P.G. and Heimler, D.
Reversed-phase and soap TLC of peptides (195) 187
Lepri, L., Desideri, P.G. and Heimler, D.
Reversed-phase and soap TLC of phenols (195) 339
Leray, C., see Raffin, J.P. (196) 323
Lertratanangkoon, K., see Horning, M.G. (181) 59
Leskawa, K.C., Saito, M. and Rosenberg, A.
Use of deoxycholate in gel filtration chromatography of synaptosomal membrane glycoproteins (193) 316
Lesko, L.J., Ericson, J., Ostheimer, G. and Marion, A.
Simultaneous determination of bupivacaine and 2,6-pipecoloxylidide in serum by GLC (182) 226
Lester, R. and Hall, D.R.
4-(*p*-methoxycinnamyloxy)-4'-methoxyazobenzene: a nematic liquid crystal for the GLC analysis of the stereochemistry of lepidopterous sex pheromones and related unsaturated fatty alcohols and derivatives (190) 35
Leung, W.J., see Budd, R.D. (190) 129
Lévesque, D. and Mallet, V.N.
GLC analysis of aminocarb and seven derivatives (200) 228
Levine, H.L., see Block, J.H. (193) 111
Levy, R.L.
Confirmatory identification of Susceptibility Card antibiotics by ion-pair and reversed-phase HPLC (192) 467
Lewin, A.H., Parker, S.R. and Carroll, F.I.
Positive identification and quantitation of isomeric cocaines by HPLC (193) 371

Lewis, J. H.
Qualitative screening procedure for the detection of benzoyl ecgonine in the urine of greyhounds (196) 337
Li Haochun, see Lu Peichang (184) 215
Li, P. P., see Warsh, J. J. (183) 483
Li Wan Po, A., see Williams, K. J. (194) 217
Li Xiouzhen, see Zhang Yukuei (197) 97
Lichty, D. G., see Markham, G. (192) 429
Lie Ken Jie, M. S. F.
Fatty acids. XIX. A quantitative treatment of saturated triglycerides by reversed-phase HPLC (192) 457
Liebich, H. M., Pickert, A., Stierle, U. and Wöll, J.
GC-MS of saturated and unsaturated dicarboxylic acids in urine (199) 181
Ligny, C. L. de, see De Ligny, C. L. (196) 51, (198) 301
Lim, C. K., Rideout, J. M. and Samson, D. M.
Determination of 5-aminolaevulinic acid and porphobilinogen by HPLC (185) 605
Lim, K. L., Young, R. W. and Driskell, J. A.
Separation of vitamin B_6 components by HPLC (188) 285
Lin, E. T., Baughman, R. A. Jr. and Benet, L. Z.
HPLC determination of prazosin in human plasma, whole blood and urine (183) 367
Lin, J.-T., Heftmann, E. and Hunter, I. R.
HPLC of the reduction products of progesterone (190) 169
Lin, K. D., Hwang, D. L. and Foard, D. E.
Affinity chromatographic interactions of proteases with low-molecular-weight soybean protease inhibitors (195) 385
Lin, P. T., see Hirata, Y. (181) 287
Lin, S.-N., Smith, L. A. and Caprioli, R. M.
Analysis of dipeptide mixtures by the combination of ion-pair reversed-phase HPLC and GC-MS techniques (197) 31
Lindberg, M., see Eriksson, B.-M. (185) 575
Linde, H. C. van der, see Van der Linde, H. C. (182) 110
Lindeberg, G.
Separation of vasopressin analogues by reversed-phase HPLC (193) 427
Lindgren, R., see Jansson, L. (181) 242
Lindner, K. R. and Mannschreck, A.
Separation of enantiomers by HPLC on triacetylcellulose (193) 308
Lindner, W.
N-Chloromethyl-4-nitrophthalimide as derivatization reagent in HPLC (in German) (198) 367
Lindner, W., LePage, J. N., Davies, G., Seitz, D. E. and Karger, B. L.
Reversed-phase separation of optical isomers of Dns-amino acids and peptides using chiral metal chelate additives (185) 323
Lindström, B., see Dahlström, H. (183) 511
Lindström, B., see Ekman, L. (182) 478
Lindström, B., see Floberg, S. (182) 63
Lindström, B., see Lanbeck, K. (182) 262
Linnett, L. M. and Atkinson, D. J.
Electronic singal differentiation as an aid to quantitation in GC (197) 1
Linssen, H. A. J., see Deelder, R. S. (185) 241
Lipecky, R., see Jost, W. (185) 403
Lisboa, B. P., see Halket, J. M. (189) 267, (192) 434
Little, J. A., see Kuksis, A. (182) 1
Litvinov, I. A., see Berezkin, V. G. (193) 132
Litwiller, R. D., see Mattox, V. R. (189) 33
Liu, S.-F., see Coutts, R. T. (190) 53
Ljungcrantz, M., see Melin, A. T. (185) 225
Lloyd, J. B. F.
Phenanthrimidazoles as fluorescent derivatives in the analysis of fatty acids by HPLC (189) 359
Lloyd, R. J., see Uden, P. C. (196) 403
Lloyd-Jones, G., see Sugden, K. (192) 228
Lo, L. Y. and Bye, A.
Specific and sensitive method for the determination of aspirin and salicylic acid in plasma using reversed-phase HPLC (181) 473
Lo, M. and Riegelman, S.
Determination of propranolol and its major metabolites in plasma and urine by HPLC without solvent extraction (183) 213
Lodge, B. A., see Gagné, D. (193) 160
Loeherlein, C., see Bosisio, A. B. (189) 317
Loenen, A. C. van, see Van Loenen, A. C. (181) 363
Loenhout, J. W. A. van, see Van Loenhout, J. W. A. (182) 487
Lohninger, A. and Nikiforov, A.
Quantitative determination of natural dipalmitoyl lecithin with dimyristoyl lecithin as internal standard by capillary GLC (192) 185
Lohse, K., see Krstulovic, A. M. (186) 733
Løkke, H.
Determination of amitrole by ion-pair HPLC (200) 234
Lombard, A., see Buffa, M. (200) 309
Lombrozo, L., Anderson, T. J., Kanaske, K. and Hollister, L. E.
Modification of assays for the routine analysis of 3-methoxy-4-hydroxyphenylglycol in urine by electron-capture GC (181) 1
Lomecky-Janousek, M., see De Vries, C. P. (183) 87
Longerich, L., see Gault, M. H. (182) 465
Lönroth, U., see Eksborg, S. (185) 583
Loo, T. L., see Rosenblum, M. G. (183) 363
Loo, T. L., see Schweinsberg, P. D. (181) 103
Lores, E. M., Meekins, F. C. and Moseman, R. F.
Determination of halogenated anilines in urine by HPLC with an electrochemical detector (188) 412
Low, G. K. C. and Haddad, P. R.
Influence of detector time constant variations on efficiency calculations in the standardisation of HPLC columns (198) 235
Low, G. K.-C., see Lea, A. R. (198) 41
Low, R. I., see Gallaher, C. (181) 490

Lu Peichang and Li Haochun
The Third National Symposium on Chromatography in China (Review) (184) 215
Lu Peichang, Zhou Liangmo, Wang Chinghai, Wang Guanghua, Xia Aizu and Xu Fangbao
High-performance GC columns packed with micro-particles (186) 25
Lu Peichang, see Zhang Yukuei (197) 97
Lucchesi, B.R., see Patterson, E. (181) 33
Ludemann, W.D., see Sass, S. (187) 447
Ludwig, G., see Voelter, W. (199) 345
Lukáš, J.
Modification of the Rohrschneider method for the classification of the polarity of polymeric sorbents based on methacrylic acid esters (190) 13
Lukás, J., Votavová, E., Švec, F., Kálal, J. and Popl, M.
Reactive polymers. XXXI. Thermally treated polymeric sorbents based on glycidylmethacrylate copolymers (194) 297
Lukevics, E., see Shatz, V.D. (200) 105
Lumper, G., see Jurenitsch, J. (189) 43
Luporsi, C., see Akiki, C. (188) 435
Lustenberger, P., Formstecher, P. and Dautrevaux, M.
Quantitative determination of alkylamino side-chains coupled to agarose beads. Comparison of methods (193) 451

M

Ma, C.-Y., Elsohly, M.A. and Baker, J.K.
HPLC separation of urushiol congeners in poison ivy and poison oak (200) 163
Ma, J.K.H., see Zia, H. (181) 77
Maass, K., see Bode, J. (190) 437
McBrayer, G., see Sosa, J.M. (189) 169
McCormick, R.M. and Karger, B.L.
Role of organic modifier sorption on retention phenomena in reversed-phase LC (199) 259
McDonough, P.S., see Cardwell, T.J. (193) 53
Macek, K., Deyl, Z. and Smrž, M.
Two-dimensional TLC of Dns-amino acids on reversed-phase silica gel (193) 421
Macek, K., see Svojtková, E. (183) 221
MacGee, J., see Mendenhall, C.L. (190) 197
McGinnis, G.D., see Ervin, H.E. (190) 203
McGuire, J.K., Miller, T.Y., Tipps, R.W., Snyder, R.S. and Righetti, P.G.
New experimental approaches to the isoelectric fractionation of cells (194) 323
MacKay, C., see Burns, D.T. (190) 141
Mackay, C., see Burns, D.T. (200) 300
MacKenzie, S.L. and Finlayson, A.J.
Analysis of cysteic acid and cysteine sulfinic acid by GLC (187) 239
MacKichan, J.J.
Simultaneous LC analysis for carbamazepine and carbamazepine 10,11-epoxide in plasma and saliva by use of double internal standardization (181) 373
McLaren, J.V., see Taylor, R. (195) 154
McLaughlin, J.L., see Doetsch, P.W. (189) 79
McLean, A.E.M., see Wilson, I.D. (194) 343
McLemore, T.L., see Marshall, M.V. (197) 217
McLennan, F., see Knox, J.H. (185) 289
McLeod, H.A., see Lawrence, J.F. (195) 113
McLimont, M.J., see Joy, K.W. (194) 76
McNair, A., see Angelo, H.R. (183) 159
McNay, J.L., see Haegele, K.D. (187) 171
McVey, J.P., see Middleditch, B.S. (195) 359
McWeeny, D.J., see Gilbert, J. (197) 71
Madajová, V., see Kaniansky, D. (194) 11
Maderspach, A., see Tomori, É. (191) 261
Maessen, F.J.M.J., see Gast, C.H. (185) 549
Maestas, R.R., Prieto, J.R., Kuehn, G.D. and Hageman, J.H.
Polyacrylamide-boronate beads saturated with biomolecules: a new general support for affinity chromatography of enzymes (189) 225
Magnuson, E., see Sjöquist, B. (183) 17
Mahapatra, G.N., Nath, J.P., Pattnaik, B.K. and Rout, D.N.
Separation of isomeric ureas and oxazoles by TLC (193) 338
Mahendroo, P.P., see Sharma, S.K. (184) 471
Mahoney, W.C. and Duksin, D.
Separation of tunicamycin homologues by reversed-phase HPLC (198) 506
Maire, C.E., see Vanderslice, J.T. (196) 176
Makin, H.L.J., see Seamark, D.A. (197) 271
Makita, M., Ohkaru, Y. and Yamamoto, S.
Resolution of histidine diastereomers by GC (188) 408
Makita, M., see Yamamoto, S. (194) 399
Makkar, H.P.S., see Sharma, O.P. (196) 515
Makowski, A., see Peterson, R.G. (188) 420
Makuch, B., see Gazda, K. (191) 9
Malcolme-Lawes, D.J. and Warwick, P.
Beta-induced fluorescence as a detection technique for LC. II. Prototype detector for normal-phase operation (200) 47
Malinar, M.J., see Ćelap, M.B. (196) 59
Malingré, T.M., see Huizing, H.J. (195) 407
Malingré, T.M., see Jellema, R. (189) 406
Maliński, E., see Błotny, G. (193) 61
Mallet, V.N., see Lévesque, D. (200) 228
Malvoisin, E., Evrard, E., Roberfroid, M. and Mercier, M.
Determination of Kováts retention indices with a capillary column and electron-capture detection: application to the assay of the enzymatic conversion of 3,4-epoxy-1-butene into diepoxybutane (186) 81
Mamer, O.A., Montgomery, J.A. and Taguchi, V.Y.
Origin of the two peaks for 2-keto-3-

methylvaleric acid produced by the oximation of the keto acids occurring in maple syrup urine disease (182) 221
Manchester, J.E. and Manchester, K.L. Separation of hypoglycin A from leucine and other amino acids on Sephadex G-10 (193) 148
Manchester, K.L., see Manchester, J.E. (193) 148
Mandal, S.B., Biswas, S.K. and Chakrabarti, P. Assay of phospholipase D on a TLC plate (188) 292
Maniara, G. and Pietr, S.J. GC determination of Lenacil and Pyrazon in soil and bacterial cultures (189) 89
Manna, C., see Zappia, V. (189) 399
Manning, W.B., see Janini, G.M. (193) 444
Manning, W.B., see Shaikh, B. (195) 392
Mannschreck, A., see Lindner, K.R. (193) 308
Månsson, M.-O., see Glad, M. (200) 254
Marcantonio, L.A., Auld, W.H.R. and Skellern, G.G. Determination of the diuretic bumetanide in biological fluids by HPLC (183) 118
Marcelin, G., Traynor, S.G., Goins, W. Jr. and Hirschy, L.M. Polydentate phosphonium salts for the deactivation of wall-coated open-tubular columns. Comparison of temperature stability with other deactivation methods (187) 57
Marcell, P.D., see Gal, J. (181) 123
Marchant, B., see Crampton, E.L. (183) 141
Marcucci, F., Garbagna, L., Monti, F., Bonazzi, P., Canobbio, L., Zuccato, E. and Mussini, E. GC determination of two fluorinated benzodiazepines in rats and mice (198) 180
Marcucci, F., see Cotellessa, L. (192) 441
Mardens, Y., see Kumps, A. (182) 116
Mårdh, P.-A., see Larsson, L. (182) 402
Marekov, N.L., see Milkova, T.S. (194) 429
Marfil, F., see Sioufi, A. (182) 241
Marion, A., see Lesko, L.J. (182) 226
Markham, G., Lichty, D.G. and Wightman, F. Comparative study of derivatisation procedures for the quantitative determination of the auxin, phenylacetic acid, by GC (192) 429
Marks, G.S., see Zelt, D.T. (189) 209
Marlow, W., see Heyes, W.F. (194) 416
Marnela, K.-M. Automated ion-exchange chromatography in the detection of aspartylglucosaminuria (182) 409
Marriott, P.J., see Cardwell, T.J. (193) 53
Marsh, K.C., Repta, A.J. and Sternson, L.A. HPLC assay for methylglyoxal bis(guanylhydrazone) (methyl GAG) in plasma and urine (187) 101
Marshall, M.A., see Robinson, J.L. (189) 145
Marshall, M.V., Gonzalez, M.A., McLemore, T.L., Busbee, D.L., Wray, N.P. and Griffin, A.C. Reversed-phase separation of benzo[*a*]pyrene metabolites by TLC (197) 217
Martin, A.J.P., Halász, I., Engelhardt, H. and Sewell, P. "Flip-flop" chromatography (186) 15
Martin, M., see Colin, H. (185) 79
Martin, T.W., see Griest, W.H. (192) 173
Martinelli, E.M., see Bombardelli, E. (196) 121
Martínez, E. and Gelpí, E. Trimethylsilylation of biogenic indoleamines. Practical kinetic and analytical aspects related to tryptamine and 5-hydroxytryptamine (186) 619
Marunaka, T., Shibata, T., Minami, Y. and Umeno, Y. Simultaneous determination of phenylbutazone and its metabolites in plasma and urine by HPLC (183) 331
Marunaka, T., Umeno, Y. and Minami, Y. Quantitative determination of 1,3-bis(tetrahydro-2-furanyl)-5-fluoro-2,4-pyrimidinedione and its metabolites in visceral tissues by HPLC and GC mass fragmentography (188) 270
Marunaka, T., Umeno, Y. and Minami, Y. Analysis of pyrimidine bases in biological materials by GC-MS (190) 107
Masahara, R., see Tsukida, K. (192) 395
Mason, D.T., see Gallaher, C. (181) 490
Massarani, G., see Zio, C.A. (200) 245
Masson, P. and Anguille, J. Determination of the free electrophoretic mobility of proteins by polyacrylamide gradient gel electrophoresis: A new approach (192) 402
Masuda, Y., see Ono, A. (197) 251
Mathes, D., see Engelhardt, H. (185) 305
Mathew, C.G.P., Goodwin, G.H. and Johns, E.W. Quantitative analysis of non-histone chromosomal proteins HMG 14 and HMG 17 by polyacrylamide gel electrophoresis (198) 80
Mathews, R.G., Torres, J. and Schwartz, R.D. Glass capillary adsorption columns for GC (186) 183
Mathews, R.G., Torres, J. and Schwartz, R.D. Applications of surface-modified porous silicas to glass capillary column preparation (199) 97
Matsuki, Y., Fukuhara, K., Ito, T., Ono, H., Ohara, N., Yui, T. and Nambara, T. Determination of captopril in biological fluids by GLC (188) 177
Matsuki, Y., see Hamano, T. (190) 462
Matsumoto, G. and Hanya, T. GC-MS identification of phenolic acids in recent sediments (193) 89
Matsumoto, G. and Hanya, T. Presence of squalane in urban aquatic environments (194) 199
Matsumoto, U. and Shibusawa, Y. Surface affinity chromatographic separation of blood cells. I. Separation of human and rabbit peripheral granulocytes, lymphocytes and erythrocytes using

polyethylene glycol-bonded column packings (187) 351
Matsumura, S., see Kakita, K. (181) 248
Matsuoka, A., see Kimura, M. (183) 467
Matsuoka, K., see Takeuchi, T. (192) 127
Matsushima, Y., see Kono, A. (182) 125
Matsushima, Y., see Sadanaga, T. (183) 246
Matthews, J.B., see Kennedy, J.F. (196) 379
Mattox, V.R. and Litwiller, R.D.
Liquid ion exchanges in PC of steroidal glucosiduronic esters (189) 33
Matz, B.
Separation of halogenated uracil derivatives from nucleobases and nucleosides by TLC on silica gel (187) 453
Maume, B.F., Millot, C., Mesnier, D., Patouraux, D., Doumas, J. and Tomori, E.
Quantitative analysis of corticosteroids in adrenal cell cultures by capillary column GC combined with MS (186) 581
Maurizis, J.C., see Michelot, J. (188) 431
Mauro, D., Wetzel, D., Lee, C.H. and Seib, P.A.
HPLC determination of L-ascorbate-2-phosphate in phosphorylation reactions (187) 421
Maya, L. and Danis, P.O.
Bis(octadecyl phosphate)zirconium(IV). Novel support for reversed-phase chromatography (190) 145
Maybaum, J., Klein, F.K. and Sadee, W.
Determination of pyrimidine ribotide and deoxyribotide pools in cultured cells and mouse liver by HPLC (188) 149
Maybaum, J., see Cohen, M.B. (198) 435
Mazrimas, J.A., see Pallavicini, M.G. (183) 449
Mazur, M., see Smith, R.M. (181) 319
Meekins, F.C., see Lores, E.M. (188) 412
Meeles, M.T.H.A., see Tjaden, U.R. (181) 227
Meer, C. van der, see Van der Meer, C. (182) 121
Meer, M.J. van der, see Van der Meer, M.J. (181) 282
Meffin, P.J. and Smith, K.J.
GC analysis of naloxone in biological fluids (183) 352
Meffin, P.J., see Smith, K.J. (181) 469
Mefford, I.N. and Barchas, J.D.
Determination of tryptophan and metabolites in rat brain and pineal tissue by reversed-phase HPLC with electrochemical detection (181) 187
Meijers, C., see Mikkers, F. (182) 496
Meinecke, B., see Gips, H. (193) 322
Meisen, A., see Choy, E.T. (187) 145
Melander, W., Stoveken, J. and Horváth, C.
Stationary phase effects in reversed-phase chromatography. I. Comparison of energetics of retention on alkyl-silica bonded phases (199) 35
Melander, W., see Horváth, C. (186) 371
Melander, W.R., Chen, B.-K. and Horváth, C.
Mobile phase effects in reversed-phase chromatography. I. Concomitant dependence of retention on column temperature and eluent composition (185) 99
Melander, W.R., Nahum, A. and Horváth, C.
Mobile phase effects in reversed-phase chromatography. III. Changes in conformation and retention of oligo(ethylene glycol) derivatives with temperature and eluent composition (185) 129
Melander, W.R., Stoveken, J. and Horváth, C.
Mobile phase effects in reversed-phase chromatography. II. Acidic amine phosphate buffers as eluents (185) 111
Melin, A.T., Ljungcrantz, M. and Schill, G.
Reversed-phase ion-pair chromatography with an adsorbing stationary phase and a hydrophobic quaternary ammonium ion in the mobile phase. I. Retention studies with tetrabutylammonium as cationic component (185) 225
Mellstrand, T., see Moberg, U. (181) 147
Mendenhall, C.L., MacGee, J. and Green, E.S.
Simple rapid and sensitive method for the simultaneous quantitation of ethanol and acetaldehyde in biological materials using head-space GC (190) 197
Mendez, J., see Vielma, H. (196) 166
Meo, M. de, see De Meo, M. (183) 357
Mercier, M., see Malvoisin, E. (186) 81
Mercik, S., see Antal, E. (183) 149
Merkus, F.W.H.M., see Zuidema, J. (182) 130
Merli, F., Novotný, M. and Lee, M.L.
Fractionation and GC analysis of aza-arenes in complex mixtures (199) 371
Merrill, Y.N., see Branfman, A.R. (182) 163
Mesnier, D., see Maume, B.F. (186) 581
Meyer, C.J., see Fenimore, D.C. (186) 555
Meyer, H.D., Jacob, K., Vogt, W. and Knedel, M.
Diagnosis of porphyrias by ion-pair HPLC (199) 339
Meyer, L.J., see Bishop, C.A. (192) 222
Meyniel, G., see Michelot, J. (188) 431
Micali, G., Curro, P. and Calabro, G.
Reversed-phase HPLC for the determination of β-asarone (194) 245
Miceli, J.N., Ryan, D.M. and Done, A.K.
High-performance liquid column chromatography of fenoprofen in serum (183) 250
Michael, G.
Gas chromatography of trifluoroacetyl amino acid trimethylsilyl esters. II. Problems in preparing specified derivatives of hydroxyamino acids (in German) (188) 251
Michael, G.
GC of trimethylsilylesters of trifluoroacetylamino acids. III. Simple method for the preparation of the trimethylsilylesters of N-trifluoroacetyl amino acids (in German) (196) 160
Michalec, Č., Ledvinová-Reinišová, J. and Kolman, Z.
Biochemistry of sphingolipids. XXVII. Thin-layer microchromatography of DNP

derivatives of long-chain bases on Silufol $_{UV\ 254}$ sheets (190) 498
Michelot, J., Godeneche, D., Maurizis, J.C. and Meyniel, G.
Separation of iodinated compounds of L-tyrosyl-I-tyrosine from iodothyronines by Biogel P-2 CC (188) 431
Middleditch, B.S. and Basile, B.
Discharge of elemental sulfur and its distribution in surficial sediments in the region of the Buccaneer oil field (199) 161
Middleditch, B.S., Missler, S.R., Hines, H.B., McVey, J.P., Brown, A., Ward, D.G. and Lawrence, A.L.
Metabolic profiles of penaeid shrimp: dietary lipids and ovarian maturation (195) 359
Midha, K.K., see Coutts, R.T. (190) 53
Mikasa, H., Sasaki, K. and Kodama, H.
Simple estimation of urinary methylmalonic acid by isotachophoretic analysis (190) 501
Mikeš, F., see Brynda, E. (196) 39
Mikeš, O., Štrop, P., Smrž, M. and Čoupek, J.
Ion-exchange derivatives of Spheron. III. Carboxylic cation exchangers (192) 159
Mikhaylenko, V.P., Sereda, I.P. and Korol, A.N.
Minimizing the retention of chromium chelates with some fluorinated β-diketones (191) 231
Mikkers, F., Verheggen, T., Everaerts, F., Hulsman, J. and Meijers, C.
Direct determination of valproate in serum by isotachophoresis. Comparison with a GC method (182) 496
Mikkers, F., see Verheggen, T. (182) 317
Miková, Z., see Julák, J. (190) 183
Miles, J.M., see Haymond, M.W. (183) 403
Milkova, T.S., Popov, S.S., Marekov, N.L. and Kovachev, G.L.
Separation of sterols using digitonin-impregnated TLC plates (194) 429
Miller, D.A. and Grimsrud, E.P.
Analysis errors following hydrogen cleaning of an electron capture detector (190) 133
Miller, D.A., see Grimsrud, E.P. (192) 117, (197) 51
Miller, R.H. Jr., see Echols, R.E. (193) 470
Miller, R.J., see Jennings, W. (186) 189
Miller, T.Y., see McGuire, J.K. (194) 323
Millot, C., see Maume, B.F. (186) 581
Mills, D.J., see Bowker, M.J. (200) 261
Mills, J.A., see Frahn, J.L. (195) 379
Mills, S.C., see Ashes, J.R. (187) 297
Mimaki, Y., see Shinozawa, S. (190) 489, (196) 463
Min, B.H., Pao, J., Garland, W.A., De Silva, J.A.F. and Parsonnet, M.
Determination of an antisecretory trimethyl prostaglandin E_2 analog in human plasma by combined capillary column GC-negative chemical ionization MS (183) 411
Minami, Y., see Marunaka, T. (183) 331, (188) 270, (190) 107
Mincsovics, E., Tyihák, E. and Kalász, H.
Resolution and retention behaviour of some dyes in overpressured TLC (191) 293
Miners, J.O., see Grgurinovich, N. (182) 237
Mingrone, G., Greco, A.V. and Passi, S.
Reversed-phase HPLC separation and quantification of individual human bile acids (183) 277
Minnikin, D.E., Hutchinson, I.G., Caldicott, A.B and Goodfellow, M.
TLC of methanolysates of mycolic acid-containing bacteria (188) 221
Minor, J.M., see Glajch, J.L. (199) 57
Miquelis, R., Penel, C. and Simon, C.
Separation of rat thyroid lysosome subpopulations on Sepharose 2B (189) 101
Misharina, T.A., see Golovnya, R.V. (190) 1
Misiti, D., see Gasparrini, F. (194) 239
Mislovičová, D., Gemeiner, P., Kuniak, Ľ. and Zemek, J.
Affinity chromatography of rat liver lactate dehydrogenase on the Remazol derivative of bead cellulose (194) 95
Missler, S.R., see Middleditch, B.S. (195) 359
Mita, H., Yasueda, H. and Shida, T.
Quantitative analysis of histamine in biological samples by GC-MS (181) 153
Mitsuhashi, Y., see Hamano, T. (190) 462
Mix, M.C., see Riley, R.T. (189) 286
Miyai, K., see Yamaguchi, Y. (182) 430
Miyamoto, T., see Kawano, Y. (198) 317
Miyazaki, H., see Fukunaga, A. (190) 339
Miyazaki, H., see Katoh, K. (188) 383
Miyazaki, H., see Someno, T. (188) 185
Miyazaki, S., Tanaka, H., Horikawa, R., Tsuchiya, H. and Imai, K.
Rapid determination of glycine- and taurine-conjugated bile acids in human bile by HPLC (181) 177
Mizobuchi, Y., Tanaka, M. and Shono, T.
Preparation and sorption behaviour of cyclodextrin polyurethane resins (194) 153
Mizutani, T.
Use of silicone-coated porous glass for exclusion chromatography in an aqueous medium (196) 485
Mladenov, I.T., see Shopova, B.I. (187) 232
Mlejnek, O.
Application of pyrolysis-capillary GC to the characterization of polyethylene (191) 181
Moberg, U., Hjalmarsson, S.-G. and Mellstrand, T.
New rapid assay of theophylline in plasma by isotachophoresis (181) 147
Moczar, E. and Waldron-Edward, D.
Differentiation of the polysaccharide side-chains of glycoproteins by a fingerprinting technique: heterogeneity of human gastric mucin (181) 108
Modderman, E.S.M., see Zuidema, J. (182) 130
Moerloose, P. de, see De Moerloose, P. (193) 255
Moffat, A.C., see Ramsey, J.D. (184) 185

Moffat, A.C., see Williams, P.L. (186) 595
Molander, M., see Dahlström, H. (183) 511
Møller Jensen, K.
Determination of amitriptyline-N-oxide, amitriptyline and nortriptyline in serum and plasma by HPLC (183) 321
Molyneux, R.J. and Roitman, J.N.
Specific detection of pyrrolizidine alkaloids on thin-layer chromatograms (195) 412
Montgomery, J.A., see Mamer, O.A. (182) 221
Monti, F., see Marcucci, F. (198) 180
Moody, R.R., Selkirk, A.B. and Taylor, R.B.
HPLC of proguanil, cycloguanil and 4-chlorophenylbiguanide using hydrophobic pairing ion and its application to serum assay (182) 359
Moore, E.D.W. and Powis, G.
Plasma and tissue concentrations of hycanthone in the mouse determined by reversed-phase HPLC (183) 372
Moore, R.G., see Nichol, H. (182) 191
Moore, R.G., see Tsuei, S.E. (181) 135
Moree-Testa, P., see Saint-Jalm, Y. (198) 188
Morel, D. and Serpinet, J.
GC evidence for phase transitions in very compact octadecyl bonded silicas (200) 95
Morgan, E.D., Evershed, R.P. and Tyler, R.C.
GC detection and structure analysis of volatile pheromones in insects (186) 605
Morgan, E.D., see Bielby, C.R. (194) 43
Morgan, E.D., see Wilson, I.D. (194) 343
Mori, S.
Empirical approach to the universal calibration of isomers of phthalate esters and similar compounds in size exclusion chromatography (192) 295
Mori, S.
Determination of the composition of copolymers as a function of molecular weight by pyrolysis GC size-exclusion chromatography (194) 163
Mori, Y.
Simple method for the purification of C1q, a subcomponent of the first component of complement by affinity chromatography using IgG-Sepharose (189) 428
Mori, Y., see Agoh, T. (182) 171
Moriyasu, M., see Endo, M. (196) 334
Moros, S.A., see Rucki, R.J. (190) 359
Morris, C.E.M. and Grabovac, I.
Performance of an evaporative analyser detector for gel permeation chromatography (189) 259
Morris, R.J., see Ballantine, J.A. (198) 111
Morris, R.O., see Crozier, A. (198) 57
Morrison, H., Avnir, D. and Zarrella, T.
Analysis of *Z* and *E* isomers of urocanic acid by HPLC (183) 83
Mortensen, H.B.
Quantitative determination of hemoglobin A_1c by thin-layer isoelectric focusing (182) 325
Mosbach, K., see Glad, M. (200) 254
Moseman, R.F., see Lores, E.M. (188) 412
Mott, S.D., see Grushka, E. (186) 117
Mourot, D., Delépine, B., Boisseau, J. and Gayot, G.
Reversed-phase ion pair chromatography of oxytetracycline, epioxytetracycline and anhydrooxytetracycline (190) 486
Mrozik, H., see Tolan, J.W. (190) 367
Mucha, I., see Tóth, G. (189) 433
Mulchandani, N.B., see Sen, G. (198) 203
Mulder-Krieger, T., see Verpoorte, R. (184) 79
Mulé, S.J., see Jukofsky, D. (198) 534
Müller, J., see Stahl, E. (189) 293
Müller-Oerlinghausen, B., see Kresse, M. (183) 475
Munroe, M., see Scott, R.P.W. (186) 475
Murakami, F.
Quantitative structure-retention relationships in cation-exchange chromatography (198) 241
Murphy, G.M., see Podesta, M.T. (182) 293
Muschik, G.M., see Janini, G.M. (193) 444
Musha, S., see Abe, I. (200) 195
Mussche, P., see Verzele, M. (190) 331
Mussini, E., see Cotellessa, L. (192) 441
Mussini, E., see Marcucci, F. (198) 180
Musson, D.G. and Sternson, L.A.
Conversion of arylhydroxylamines to electrochemically-active derivatives suitable for HPLC analysis with amperometric detection (188) 159
Mutschler, E., see Schumacher, A. (181) 512
Mutschler, E., see Sobel, M. (183) 124
Myers, M.N., see Giddings, J.C. (185) 261, (186) 37
Myher, J.J., see Kuksis, A. (182) 1

N

Nachtmann, F., see Lankmayr, E.P. (198) 471
Nader, H.B., see Bianchini, P. (196) 455
Nagase, Y., see Shimada, K. (181) 51
Nagatsu, T., see Kojima, K. (189) 233
Nagels, L., Debeuf, C. and Esmans, E.
Quantitative determination of quinic acid and derivatives by HPLC after derivatization with *p*-bromophenacyl bromide (190) 411
Nagels, L., Van Dongen, W., De Brucker, J. and De Pooter, H.
HPLC separation of naturally occurring esters of phenolic acids (187) 181
Nagumo, T., see Iizima, N. (193) 464
Nahum, A. and Horváth, C.
Evaluation of octanol-water partition coefficients by using HPLC (192) 315
Nahum, A., see Gautam, S.R. (182) 482
Nahum, A., see Horváth, C. (186) 371
Nahum, A., see Melander, W.R. (185) 129

Naikwadi, K.P., Panse, D.G., Bapat, B.V. and Ghatge, B.B.
Liquid crystals. I. Synthesis and application as stationary phases in GLC (195) 309
Nair, B.R. and Francis, J.D.
TLC method for the separation and determination of the products of the reaction of amides with formaldehyde (195) 158
Naito, K. and Takei, S.
Adsorption effects in gas-liquid partition chromatography. Quantitative evaluation (190) 21
Nakagaki, M., see Nakayama, F. (183) 287
Nakagawa, G., see Tsuda, T. (199) 249
Nakamori, H., see Janado, M. (193) 345
Nakamura, A., Kojima, S. and Kaniwa, M.-A.
Quantitative GC determination of tris(2,3-dibromopropyl)phosphate in the 10-ng range by using a 0.8-mm I.D. column packed with a high liquid loaded support (196) 133
Nakamura, H., Tsuzuki, S., Tamura, Z., Yoda, R. and Yamamoto, Y.
Fluorescence detection of secondary amines on thin-layer plates using a fluorogenic reaction with fluorescamine (200) 324
Nakamura, K., Kuwahara, A., Ogata, H. and Takeo, K.
Study of the interaction between L-lactate dehydrogenase isoenzymes and immobilized 8-substituted adenosine 5'-monophosphate by means of affinity electrophoresis (192) 351
Nakamura, K., Kuwahara, A. and Takeo, K.
Study of the interaction between NADP-dependent dehydrogenase and immobilized adenosine 2'-monophosphate by means of affinity electrophoresis (196) 85
Nakanishi, M., see Sadanaga, T. (183) 246
Nakano, N.I., Shimamori, Y. and Yamaguchi, S.
Mutual displacement interactions in the binding of two drugs of human serum albumin by frontal affinity chromatography (188) 347
Nakayama, F. and Nakagaki, M.
Quantitative determination of bile acids in bile with reversed-phase HPLC (183) 287
Namba, K., see Nobuhara, Y. (190) 251
Nambara, T., see Matsuki, Y. (188) 177
Nasal, A., see Grzybowski, J. (196) 217
Nath, J.P., see Mahapatra, G.N. (193) 338
Nation, R.L., see Petersen, M.C. (183) 131
Nation, R.L., see Tsuei, S.E. (182) 55
Nau, H., Jesdinsky, D. and Wittfoht, W.
Microassay for primidone and its metabolites phenylethylmalondiamide, phenobarbital, and *p*-hydroxyphenobarbital in human serum, saliva, breast milk and tissues by GC-MS using selected ion monitoring (182) 71
Nawrocki, J., Szczepaniak, W. and Wasiak, W.
Coordination polymers as adsorbents and stationary phases in GC. VI. Separation properties of alkaline-earth metal di-*n*-hexylphosphinates (188) 323
Nayar, M.S.B., see Cohen, J.L. (181) 478
Neijt, J.P., see Hulshoff, A. (181) 363
Neil, J.M., see Fell, A.F. (186) 691
Neill, D.B., see Holdiness, M.R. (198) 329
Nelis, H.J.C.F. and De Leenheer, A.P.
Retention mechanisms of tetracyclines on a C_8 reversed-phase material (195) 35
Nelson, L.M. and Carruthers, M.
Semi-automated fluorimetric method for the estimation of urinary catecholamines using HPLC (183) 295
Neukom, H., see Waibel, R. (197) 86
Neukom, H.P., see Grob, K. Jr. (189) 109, (198) 64
Newman, R.A., see Allen, B.A. (190) 241
Ng, D.S., see Blass, K.G. (182) 311
Nguyen-Hoang-Nam, see Scherrmann, J.M. (196) 342
Nice, E.C., Capp, M. and O'Hare, M.J.
Use of hydrophobic interaction methods in the isolation of proteins from endocrine and paraendocrine tissues and cells by HPLC (185) 413
Nice, E.C., see O'Hare, M.J. (198) 23
Nichol, H., Vine, J., Thomas, J. and Moore, R.G.
Quantitation of doxapram in blood, plasma and urine (182) 191
Nichols, B.L., see Huang, C.T.L. (196) 150
Nickless, G., see Cooke, M. (187) 47
Nickless, G., see Ober, A.G. (196) 237
Nicolas, A., see Sarhan, F. (183) 505
Nieass, C.S., Wainwright, M.S. and Chaplin, R.P.
Sampling and HPLC analysis of organic compounds in liquified carbon dioxide (194) 335
Nieder, M., see Bayer, E. (186) 497
Niederberger, W., Schaub, P. and Beveridge, T.
HPLC determination of cyclosporin A in human plasma and urine (182) 454
Niekerk, P.J. van, see Van Niekerk, P.J. (187) 436
Nieman, T.A., see Veazey, R.L. (200) 153
Nieminen, A.-L., Kangas, L., Anttila, M. and Hautoniemi, L.
Determination of methenamine in biological samples by GLC (181) 11
Nierenberg, D.W., Pogolotti, A.L. Jr. and Santi, D.V.
Detection and quantitation of diadenosine tetraphosphate by HPLC (190) 212
Niijima, K., see Someno, T. (188) 185
Nijkerk, A.J., see Guelen, P.J.M. (181) 497
Nikiforov, A., see Lohninger, A. (192) 185
Nilsson, B., see Jansson, L. (181) 242
Nilsson, B.F. and Samuelson, O.
Chromatographic behaviour of aromatic compounds on anion-exchange resins in various carboxylate forms (198) 267
Nilsson, G., see Ekman, L. (182) 478
Nishida, K., see Uobe, K. (193) 83
Nishida, S., see Kakita, K. (181) 248
Nishida, S., see Kimura, M. (188) 417

Nishida, T., see Janado, M. (193) 345
Noach, E.L., see Broxterman, H.J. (183) 185
Nobuhara, Y., Hirano, S., Namba, K. and Hashimoto, M.
Separation and determination of opium alkaloids by HPLC (190) 251
Noda, K., see Suzuki, A. (182) 448
Noggle, F.T. Jr., see Clark, C.R. (188) 426
Noguchi, H., see Suzuki, A. (182) 448
Nooitgedacht, F., see De Jong, A.W.J. (193) 181
Nordlöv, H., Jordan, P.M., Burton, G. and Scott, A.I.
Improved separation of uroporphyrin isomers by HPLC (190) 221
Nose, N., see Tanaka, A. (194) 21
Novák, J., Vejrosta, J., Roth, M. and Janák, J.
Correlation of GC specific retention volumes of homologous compounds with temperature and methylene number (199) 209
Novák, J., see Rijks, J.A. (186) 167
Novotný, M., see Hirata, Y. (181) 287, (186) 521
Novotný, M., see Merli, F. (199) 371
Novotný, M., see Smith, S.L. (187) 111
Nozaki, H., see Hiramatsu, K. (182) 301
Nozawa, A. and Ohnuma, T.
Improved HPLC analysis of ethylene oxide condensates by their esterification with 3,5-dinitrobenzoyl chloride (187) 261
Nurit, Y., see Pernes, J.-F. (181) 254
Nyheim, A., see Österlöf, G. (183) 487

O

Ober, A.G., Cooke, M. and Nickless, G.
Adsorption chromatography capillary columns. I. Rubidium chloride (196) 237
Ober, A.G., see Cooke, M. (195) 265
Obwexer, H.W., see Vogt, W. (186) 197
Ochiai, M.
Determination of dissolved carbohydrates in natural water by GLC (194) 224
Ochoa, J.L., Porath, J., Kempf, J. and Egly, J.-M.
Electron donor-acceptor properties of urea and its role in charge-transfer chromatography (188) 257
Oda, T., see Shinozawa, S. (190) 489, (196) 463
Odham, G., see Larsson, L. (182) 402
O'Donnell, J.P., see Zia, H. (181) 77
Oerlemans, F., see Verheggen, T. (182) 317
Öfverstedt, L.-G., see Hjertén, S. (194) 1
Ogan, K. and Katz, E.
Retention characteristics of several bonded-phase LC columns for some polycyclic aromatic hydrocarbons (188) 115
Ogata, H., see Nakamura, K. (192) 351
Ogata, J.N., Okun, J.D., Hylin, J.W. and Bevenue, A.
GC method for the analysis of polychlorinated biphenyls in transformer oil (189) 425
Ogierman, L., Czech, B. and Piórko, A.
TLC separation of some ferrocene alcohols (198) 536
Ogilvie, K.K., see Quilliam, M.A. (194) 379, (196) 367
Ohara, N., see Matsuki, Y. (188) 177
O'Hare, M.J., Nice, E.C. and Capp, M.
Reversed- and normal-phase HPLC of 18-hydroxylated steroids and their derivatives. Comparison of selectivity, efficiency and recovery from biological samples (198) 23
O'Hare, M.J., see Nice, E.C. (185) 413
Ohashi, S., see Yoza, N. (196) 471
Ohkaru, Y., see Makita, M. (188) 408
Ohkura, Y., see Sadanaga, T. (183) 246
Ohlson, S., see Glad, M. (200) 254
Ohmae, M., see Tamegai, T. (193) 483
Ohmori, S., see Ikeda, M. (183) 41
Ohnesorge, A.-B., see Vogt, W. (186) 197, (199) 191
Ohnuma, T., see Nozawa, A. (187) 261
Ohsawa, A., see Hara, S. (200) 85
Okahara, S., see Yamamoto, S. (194) 399
O'Kelly, D.A., see Brodie, R.R. (182) 379
Okumura, T.
Sintered TLC (Review) (184) 37
Okun, J.D., see Ogata, J.N. (189) 425
Okuyama, T., see Seta, K. (181) 311
Olieman, C., see Schaaper, W.M.M. (195) 181
Oliveros, L., see Foucault, A. (185) 345
Olsen, O., see Helboe, P. (197) 199
Olson, E.B. Jr., see Ghias-Ud-Din, M. (192) 463
Onishi, S., Itoh, S., Kawade, N., Isobe, K. and Sugiyama, S.
Accurate and sensitive analysis of ethyl anthranilate azopigments from bile by reversed-phase HPLC (182) 105
Ono, A.
Separation of xylenol isomers on nitrofluorenes and nitro-9-fluorenones by GLC. I. Mixed columns and effect of temperature on retention behaviour (193) 300
Ono, A. and Masuda, Y.
Separation behaviour of some isomeric organic compounds on sugars, sugar alcohols and their mixed phases by GLC (197) 251
Ono, H., see Matsuki, Y. (188) 177
Onodera, S., Tabata, M., Suzuki, S. and Ishikura, S.
GC identification and determination of chlorinated quinones formed during chlorination of dihydric phenols with hypochlorite in dilute aqueous solution (200) 137
Onuska, F.I.
Poly(α-olefins): novel non-polar stationary phases for GC (186) 259
Opheim, K.E., see Raisys, V.A. (183) 441
Orbán, E., see Tomori, É. (191) 261

O'Reilly, W.J., Guelen, P.J.M., Hoes, M.J.A. and Van der Kleyn, E.
HPLC determination of pyridoxine and congeners in biological fluids of man after high-dose therapy (183) 492
Oreste, P. and Torri, G.
Fingerprinting of heparins by low-amperage electrophoresis in barium acetate (195) 398
Orme, M.L'E., see Walker, R.W. (181) 85
Oró, J., see Holzer, G. (186) 795, (194) 410
Ościk-Mendyk, B., see Jaroniec, M. (188) 27
Oseekey, K.B., Rowse, K.L. and Kostenbauder, H.B.
HPLC determination of chloramphenicol and its monosuccinate ester in plasma (182) 459
Oshima, R., Takai, N. and Kumanotani, J.
Improved separation of anomers of saccharides by HPLC on macroreticular anion-exchange resin in the sulphate form (192) 452
Oshima, R., see Yamauchi, Y. (198) 49
Osima, B., see Bianchini, P. (196) 455
Osman, M., Hill, H.H. Jr., Holdren, M. and Westberg, H.
Vapor-phase silylation of alcohols for air analysis (186) 273
Osman, S.F., see Heisler, E.G. (194) 89
Osselton, M.D., see Ramsey, J.D. (184) 185
Ossicini, L., see Cardaci, V. (198) 76
Östelius, J., see Arbin, A. (193) 405
Österlöf, G. and Nyheim, A.
The estimation of α-tocopherol in biological material by GC (183) 487
Ostheimer, G., see Lesko, L.J. (182) 226
Ostrovský, I., see Soják, L. (191) 187
Ottrey, A.L., see Perkal, M. (196) 180
Ouedraogo, M.A., see Bezard, J.A. (196) 279
Ouhrabková, J., see Vrátný, P. (191) 313
Ovodov, Y.S., see Evtushenko, E.V. (196) 331
Owen, J.A., see Zelt, D.T. (189) 209
Owens, C., see Vasiliades, J. (182) 439
Oyama, H., see Kakita, K. (181) 248
Ozawa, Y., see Inayama, S. (194) 85
Özcimder, M. and Hammers, W.E.
Fractionation of fish oil fatty acid methyl esters by means of argentation and reversed-phase HPLC, and its utility in total fatty acid analysis (187) 307

P

Paanakker, J.E. and Van de Laar, G.L.M.
Determination of Org NC 45 (a myoneural blocking agent) in human plasma using normal-phase HPLC (183) 459
Paasivirta, J., see Knuutinen, J. (194) 55
Paasivirta, J., see Sattar, M.A. (189) 73
Pacáková, V., see Dufek, P. (187) 341, (191) 115
Pacáková, V., see Štulík, K. (192) 135
Packer, N.H., see Batley, M. (198) 520
Pagé, M., see Emond, J.P. (200) 57
Pajurek, J., see Krejčí, M. (191) 17
Pallavicini, M.G. and Mazrimas, J.A.
HPLC analysis of cytosine arabinoside and metabolites in biological samples (183) 449
Pandey, R.C. and Toussaint, M.W.
HPLC and TLC of anthracycline antibiotics. Separation and identification of components of the daunorubicin complex from fermentation broth (198) 407
Panina, L.I., see Sakodynskii, K.I. (186) 227
Panopio, L.G., see Lawrence, J.F. (195) 113
Panse, D.G., see Naikwadi, K.P. (195) 309
Pantarotto, C., see Bidoli, F. (196) 314
Pao, J., see Min, B.H. (183) 411
Papazova, D., Dimov, N. and Bonchev, D.
Calculation of GC retention indices of isoalkanes based on a topological approach (188) 297
Papirer, E., see Kessaissia, Z. (196) 481
Papp, O., Szász, G. and Valkó, K.
Relation of GC behaviour to the chemical structure of pyrido[1,2α]pyrimidine derivatives (191) 245
Parihar, D.B., see Vasundhara, T.S. (194) 254
Pařízková, H. and Blattná, J.
Preparative TLC of the oxidation products of retinyl acetate (191) 301
Parker, C.E., see Albro, P.W. (197) 155
Parker, G.A., see Sass, S. (189) 331
Parker, S.R., see Lewin, A.H. (193) 371
Parks, E.J., Brinckman, F.E. and Blair, W.R.
Application of a graphite furnace atomic absorption detector automatically coupled to a high-performance liquid chromatograph for speciation of metal-containing macromolecules (185) 563
Parks, O.W., see Doerr, R.C. (196) 498
Parris, N., see Doerr, R.C. (196) 498
Parsonnet, M., see Min, B.H. (183) 411
Parsons, D.L.
Determination of ligand-macromolecule binding parameters and the method of Hummel and Dreyer (193) 520
Parsons, M.L., see Clement, R.E. (189) 53, (190) 136
Paryjczak, T., Rynkowski, J. and Karski, S.
Thermoprogrammed reduction of cobalt oxide catalysts (188) 254
Passi, S., see Mingrone, G. (183) 277
Pasutto, F.M., see Coutts, R.T. (195) 105
Patankar, A.V., see Sen, G. (198) 203
Paterson, A.R., see Hanrahan, J.M. (193) 265
Patouraux, D., see Maume, B.F. (186) 581
Patterson, E., Stetson, P. and Lucchesi, B.R.
Sensitive GC assay for the quantitation of bretylium in plasma, urine and myocardial tissue (181) 33
Patterson, J.I. and Brown, R.R.
Determination of urinary quinolinic acid by HPLC (182) 425

Patthy, M. and Tomori, É.
HPLC and GLC of some norgestrel intermediates. Physical properties of the isolated *syn*- and *anti*-isomers of oximes (191) 145
Pattinson, N., Collins, D. and Campbell, B.
Covalent coupling of cholic acid to aminohexylamino-Sepharose 4B and its use in affinity chromatography of serum albumin (187) 409
Pattnaik, B.K., see Mahapatra, G.N. (193) 338
Patton, J.F., see Gay, M.L. (187) 153
Patzelová, V.
Stabilized Y-type zeolites for GC separation (191) 175
Patzelová, V., Wurst, M. and Čatská, V.
JADO carbon as an adsorbent in sampling columns (193) 137
Pauncz, J.K. and Harsányi, I.
Aminoglycoside antibiotics: TLC bioautographic detection and quantitative assay (195) 251
Pav, J.W., see Giddings, J.C. (185) 261
Pearsall, K.A. and Bonner, F.T.
Analysis of dinitrogen-nitrogen oxide mixtures employing direct vacuum line/gas chromatograph injection (200) 224
Peikert, M., see Reece, P.A. (181) 207
Pellizzari, E.D., see Bunch, J.E. (186) 811
Pelster, E., see Rohrschneider, L. (186) 249
Penel, C., see Miquelis, R. (189) 101
Pennock, C.A., see Randell, J.A.J. (195) 257
Peralta, E., Yang, H.-Y.T., Hong, J. and Costa, E.
Combination of enzymatic and mass fragmentographic assays for the identification and measurement of [met^5]-enkephalin (190) 43
Percheron, F., see Akiki, C. (188) 435
Percheron, F., see Weber, M. (188) 377
Perier, C., Ronzière, M.C., Rattner, A. and Frey, J.
Employment of GLC for the analysis of collagen amino acids in biopsy tissue (182) 155
Perkal, M., Blackman, G.L., Ottrey, A.L. and Turner, L.K.
Determination of hallucinogenic components of *Psilocybe* mushrooms using HPLC (196) 180
Perkins, E.G., see Reingold, R.N. (190) 237
Perlman, S., see Kirschbaum, J. (200) 282
Pernes, J.-F., Nurit, Y. and De Heaulme, M.
Lipids: TLC separation in twelve fractions by three successive unidirectional developments on the same plate (181) 254
Personnaz, L., see Gareil, P. (192) 53
Persson, B.-A., see Eriksson, B.-M. (185) 575
Petersen, M.C., Nation, R.L. and Ashley, J.J.
Simultaneous determination of betamethasone, betamethasone acetate and hydrocortisone in biological fluids using HPLC (183) 131
Peterson, R.G., Rumack, B.H., Sullivan, J.B. Jr. and Makowski, A.
Amperometric HPLC method for narcotic alkaloids (188) 420
Petrie, G., see Kirschbaum, J. (190) 481
Petró-Turcza, M., see Vigh, G. (193) 432
Petroff, N., see Durand, J.P. (190) 85
Petronio, B.M., see Laganà, A. (198) 143
Petrusek, M., see Bidlingmeyer, B.A. (186) 419
Pettitt, B.C. Jr., see Damon, C.E. (195) 243
Pfeifer, P., see Ende, M. (183) 1
Pfrepper, G.
Stability of ion exchangers against treatment with nitric acid and radiation. II. Change in sorption properties of Dowex A-1, Chelex 100 and Wof MC 50 after treatment with nitric acid (in German) (198) 257
Philip, J. and Chafetz, L.
TLC estimation of 2,4-xylenol in 2,5-xylenol (192) 250
Phillips, C.S.G., see Horrocks, J.P. (197) 109
Phillips, L.V., see Wright, B.W. (199) 355
Phillips, M.P., see Sievers, R.E. (186) 3
Phillips, R.D., Hayes, A.W. and Berndt, W.O.
HPLC analysis of the mycotoxin citrinin and its application to biological fluids (190) 419
Phillips, S.F., see Bouska, J.B. (183) 72
Picciano, M.F., see Reingold, R.N. (190) 237
Picer, M., see Picer, N. (193) 357
Picer, N. and Picer, M.
Evaluation of macroreticular resins for the determination of low concentrations of chlorinated hydrocarbons in sea water and tap water (193) 357
Pickert, A., see Liebich, H.M. (199) 181
Pickett, E.E., see Aigner-Held, R. (189) 139
Pickett, E.E., see Aue, W.A. (189) 13
Picman, A.K., Ranieri, R.L., Towers, G.H.N. and Lam, J.
Visualization reagents for sesquiterpene lactones and polyacetylenes on thin-layer chromatograms (189) 187
Pierini, N., see Quercia, V. (193) 163
Pietr, S.J., see Maniara, G. (189) 89
Pietta, P., see Zio, C.A. (200) 245
Pike, R.K., see Barry, C. (195) 151
Pike, R.K., see Lanouette, M. (190) 208
Pile, J., see Walker, R.W. (181) 85
Pileire, B., Beaune, P., Laudat, M.H. and Cartier, P.
The use of formic acid in carrier gas. Rapid method for identification and determination of phytanic acid by GC-MS-chemical ionization (182) 269
Pillai, D., see Dilli, S. (190) 113
Pilon, J.C., see Ryan, J.J. (197) 171
Pinedo, H.M., see Hulshoff, A. (181) 363
Pinnock, A., see Shaikh, B. (195) 392
Piórko, A., see Ogierman, L. (198) 536
Pirkle, W.H., House, D.W. and Finn, J.M.
Broad spectrum resolution of optical isomers using chiral HPLC bonded phases (192) 143
Pitkänen, E., see Lajunen, K. (187) 455
Pitts, A.F., see Pitts, F.N. Jr. (193) 157

Pitts, F.N. Jr., Yago, L.S., Aniline, O. and Pitts, A.F.
Capillary GC with a nitrogen detector for measurement of phencyclidine, ketamine and other arylcycloalkylamines in the picogram range (193) 157
Pizzie, R.A., see Howard, A.G. (189) 436
Plag, S.M., see Fell, A.F. (186) 691
Plessis, L.M. du, see Du Plessis, L.M. (187) 436
Plzák, Z., see Smolková, E. (191) 61
Podesta, M.T., Murphy, G.M. and Dowling, R.H.
Measurement of faecal bile acid sulphates (182) 293
Poehland, B.L., Chan, J.A. and Chiu, C.C.
Rapid quantitative assay for 3,6-bis-(5-chloro-2-piperidyl)-2,5-piperazinedione in fermentation broth using HPLC (190) 246
Poet, R., see Kirschbaum, J. (190) 481
Pogolotti, A.L. Jr., see Nierenberg, D.W. (190) 212
Poh, T.T., see Dutt, M.C. (195) 133
Poirier, C.A., see Estes, S.A. (196) 265
Poll, J.M. van der, see Van der Poll, J.M. (187) 244, (197) 263
Polster, M., see Rittich, B. (197) 43
Pommier, F., see Sioufi, A. (181) 161, (183) 33
Poole, C.F., Johansson, L. and Vessman, J.
Formation of electron-capturing derivatives of alprenolol by transboronation. Application to the determination of alprenolol in plasma (194) 365
Poole, C.F., Singhawangcha, S., Chen Hu, L.-E., Sye, W.-F., Brazell, R. and Zlatkis, A.
tert.-Butylpentafluorophenylmethylchlorosilane as a reagent for the formation of hydrolytically stable alkylsilyl derivatives with electron-capturing properties (187) 331
Poole, C.F., Singhawangcha, S. and Zlatkis, A.
Comparison of electron-capturing boronic acids for the selective analysis of bifunctional compounds (186) 307
Poole, C.F., Sye, W.-F., Singhawangcha, S., Hsu, F., Zlatkis, A., Arfwidsson, A. and Vessman, J.
New electron-capturing pentafluorophenyldialkylchlorosilanes as versatile derivatizing reagents for GC (199) 123
Poole, C.F. and Zlatkis, A.
Cyclic derivatives for the selective chromatographic analysis of bifunctional compounds (Review) (184) 99
Poole, C.F., see Singhawangcha, S. (183) 433
Poole, C.F., see Zlatkis, A. (182) 137
Pooter, H. de, see De Pooter, H. (187) 181
Popl, M., see Lukás, J. (194) 297
Popl, M., see Šmejkal, F. (197) 147
Popov, S.S., see Milkova, T.S. (194) 429
Popp-Snijders, C., see De Vries, C.P. (183) 87
Poppe, H., see De Jong, A.W.J. (193) 181
Poppe, H., see Gast, C.H. (185) 549
Poppe, H., see Jonker, R.J. (186) 285
Poppe, H., see Snyder, L.R. (184) 363
Porath, J., see Hubert, P. (198) 247
Porath, J., see Ochoa, J.L. (188) 257
Porcelli, M., see Zappia, V. (189) 399
Porsch, B., see Vozka, S. (193) 128
Porskamp, P.A.T.W., see Hekman, P. (182) 252
Poshkus, D.P. and Grumadas, A.J.
Determination of some molecular structure parameters for hydrocarbons from the retention volumes on graphitized carbon black (191) 169
Poulin, G., see Szymanowicz, A.G. (190) 457
Powis, G. and Ames, M.M.
Determination of 6-diazo-5-oxo-L-norleucine in plasma and urine by reversed-phase HPLC of the dansyl derivative (181) 95
Powis, G., see Moore, E.D.W. (183) 372
Pragnell, R.J., see Taylor, R. (195) 154
Prasad, V.K., see Rapaka, R.S. (196) 512
Preskorn, S.H., Leonard, K. and Hignite, C.
LC of amitriptyline and related tricyclic compounds (197) 246
Prey, T., see Wurst, F. (188) 452
Price, W.P. Jr., see Bidlingmeyer, B.A. (186) 419
Prieto, J.R., see Maestas, R.R. (189) 225
Přikryl, Z., see Wurst, M. (191) 129
Přistoupil, T.I., Kramlová, M. and Kraml, J.
Electrofocusing of stroma-free haemoglobin on a thin layer of agarose gel (196) 142
Přistoupil, T.I., see Kramlová, M. (193) 515
Pritchard, D.G., see Barnes, S. (183) 269
Procházka, M. and Smolková-Keulemansová, E.
Silica gel modified with phthalocyanines as a stationary phase in gas-solid chromatography (189) 25
Proveaux, W.J., see Zia, H. (181) 77
Prusiewicz, K., see Gazda, K. (191) 9
Průšová, D., see Punčochářová, J. (191) 81
Průšová, D., see Vodička, L. (198) 457, (200) 238
Puchinger, L., see Wurst, F. (188) 452
Pudova, O.A., see Shatz, V.D. (200) 105
Punčochářová, J., Kříž, J., Vodička, L. and Průšová, D.
Influence of mobile phase composition and nature of sample on retention data from HPLC (191) 81
Purnell, C.J., see Bagon, D.A. (190) 175
Purnell, J.H., see Al-Thamir, W.K. (188) 79
Purokoski, S., see Lajunen, K. (187) 455
Putterman, G.J., see Ito, Y. (193) 37
Pyysalo, H., Seppä, E.-L. and Widén, K.-G.
Application of GC to the analysis of sesquiterpene lactones from *Lactarius (Russulaceae)* mushrooms (190) 466
Pyysalo, H., see Widén, C.-J. (188) 213

Q

Quarterman, C.P., Kendall, M.J. and Jack, D.B.
Determination of metoprolol metabolites in plasma and urine by electron-capture GLC (183) 92
Quercia, V., Battaglino, G., Pierini, N. and Turchetto, L.
Determination of the bitter constituents of the Gentiana root by HPLC (193) 163
Quick, M.P., see Ting, H.-H. (195) 441
Quilliam, M.A., Ogilvie, K.K., Sadana, K.L. and Westmore, J.B.
Study of rearrangement reactions occurring during GC of *tert.* -butyldimethylsilyl ether derivatives of uridine (194) 379
Quilliam, M.A., Ogilvie, K.K., Sadana, K.L. and Westmore, J.B.
GC retention data for silyl and acyl derivatives of nucleosides (196) 367
Qureshi, S., see Verzele, M. (199) 105

R

Raabe, O.G., see Jennings, W. (186) 189
Radecki, A. and Halkiewicz, J.
Quantitative analysis of zinc, copper and nickel diethyldithiocarbamates by GLC (187) 363
Radecki, A., see Grzybowski, J. (196) 217
Radhakrishnamurthy, B., Dalferes, E.R. Jr., Vijayagopal, P. and Berenson, G.S.
Determination of molecular-weight distribution of aorta glycosaminoglycans by automated gel filtration (192) 307
Radivojša, P.N., see Ćelap, M.B. (196) 59
Radjai, M.K. and Hatch, R.T.
Fast determination of free amino acids by ion-pair HPLC using on-line post-column derivatization (196) 319
Raffin, J.P. and Leray, C.
Control of 8-bromo-AMP and 8-(6-aminohexyl)amino-AMP synthesis by HPLC (196) 323
Ragione, F. della, see Della Ragione, F. (189) 399
Raisys, V.A., Friel, P.N., Graaff, P.R., Opheim, K.E. and Wilensky, A.J.
HPLC and GLC determination of diazepam and nordiazepam in plasma (183) 441
Rakhit, A., see Guentert, T.W. (183) 514
Ramsdell, H.S. and Tanaka, K.
GC retention indices of twenty metabolically important acylglycines as trimethylsilyl derivatives (181) 90
Ramsey, J.D., Lee, T.D., Osselton, M.D. and Moffat, A.C.
GLC retention indices of 296 non-drug substances on SE-30 or OV-1 likely to be encountered in toxicological analyses (Review) (184) 185
Randell, J.A.J. and Pennock, C.A.
Brain gangliosides: an improved simple method for their extraction and identification (195) 257
Ranieri, R.L., see Picman, A.K. (189) 187
Rankin, J., see Ghias-Ud-Din, M. (192) 463
Rao, G.H.R., Krick, T.P. and White, J.G.
Preparation, separation and characterization of vitamin E quinone (196) 506
Rapaka, R.S., Knight, P.W. and Prasad, V.K.
Quantitative elution of thyroidal amino acids from a cation-exchange column: analysis of tablet formulations containing dyes (196) 512
Rasmussen, K.E., see Christophersen, A.S. (192) 363
Ratech, H., Thorbecke, G.J. and Hirschhorn, R.
Identification of adenosine and eight modified adenine nucleosides using reversed-phase HPLC and enzymatic peak shift with adenosine deaminase (183) 499
Rattner, A., see Perier, C. (182) 155
Ravenscroft, P.J., see Ahokas, J.T. (183) 65
Rayss, J., Witkiewicz, Z., Waksmundzki, A. and Dąbrowski, R.
Effect of the support surface on the structure of the film of liquid crystalline stationary phase (188) 107
Redant, G., see Verzele, M. (199) 105
Reddrop, C.J., Riess, W. and Slater, T.F.
Rapid, sensitive determination of unchanged promethazine in biological material using a nitrogen-selective flame ionization detector. Identification of metabolites by GC-MS (192) 375
Reddrop, C.J., Riess, W. and Slater, T.F.
Two rapid methods for the simultaneous GLC determination of carbon tetrachloride and chloroform in biological material and expired air (193) 71
Reding, R., see Sorrell, R.K. (185) 655
Redmond, J.W., see Batley, M. (198) 520
Reece, P.A., Cozamanis, I. and Zacest, R.
Selective HPLC assays for hydralazine and its metabolites in plasma of man (181) 427
Reece, P.A. and Peikert, M.
Simple and selective HPLC method for estimating plasma quinidine levels (181) 207
Reece, P.A. and Stanley, P.E.
HPLC assay for tocainide in human plasma: comparison with GLC assay (183) 109
Rees, J.A., see Crampton, E.L. (183) 141
Reeve, J., Kuhlenkamp, J. and Kaplowitz, N.
Estimation of glutathione in rat liver by reversed-phase HPLC: separation from cysteine and γ-glutamylcysteine (194) 424
Regnier, F.E., see Alpert, A.J. (185) 375

Regnier, F.E., see Barth, H.G. (192) 275
Řehák, V. and Smolková, E.
Comparison of linear and branched non-polar chemically bonded stationary phases in HPLC (191) 71
Reingold, R.N., Picciano, M.F. and Perkins, E.G.
Separation of folate derivatives by *in situ* paired-ion HPLC (190) 237
Rejholec, V., see Kuchař, M. (195) 329
Renck, B. and Einarsson, R.
Application of high-speed aqueous gel permeation chromatography to insect venoms (197) 278
Renema, J., see Hulshoff, A. (186) 535
Rennert, O.M., see Chan, W.Y. (181) 259
Repta, A.J., see Marsh, K.C. (187) 101
Rey, F., see Curtius, H.-C. (199) 171
Rice, R.H., see Agris, P.F. (194) 205
Richard, J.J., Chriswell, C.D. and Fritz, J.S.
Concentration and determination of organic acids in complex aqueous samples (199) 143
Richardson, D.G., see Akhavan, I. (190) 452
Richardson, J.P., see Tarbit, I.F. (181) 337
Rideout, J.M., see Lim, C.K. (185) 605
Riedo, F. and Kováts, E. sz.
Effect of adsorption on solute retention in GLC (186) 47
Riegelman, S., see Guentert, T.W. (183) 514
Riegelman, S., see Lo, M. (183) 213
Riegelman, S., see Upton, R.A. (190) 119
Riess, W., see Reddrop, C.J. (192) 375, (193) 71
Rietvelde, D., see Bredael, P. (192) 216
Righetti, P.G.
Molarity and ionic strength of focused carrier ampholytes in isoelectric focusing (190) 275
Righetti, P.G., Gianazza, E. and Ek, K.
New developments in isoelectric focusing (Review) (184) 415
Righetti, P.G., see Bosisio, A.B. (189) 317
Righetti, P.G., see Gianazza, E. (193) 1
Righetti, P.G., see McGuire, J.K. (194) 323
Rijks, J.A., Drozd, J. and Novák, J.
Versatile all-glass splitless sample-introduction system for trace analysis by capillary GC (186) 167
Rijks, J.A., see Dubois, J.-E. (194) 121
Rijks, J.A., see Soják, L. (191) 187
Rijks, J.A., see Uralets, V.P. (194) 135
Riley, C.M., Tomlinson, E. and Jefferies, T.M.
Functional group behaviour in ion-pair reversed-phase HPLC using surface-active pairing ions (185) 197
Riley, R.T. and Mix, M.C.
TLC of citric acid cycle intermediates, lactic acid, and the amino acid taurine (189) 286
Rinehart, R., see James, G.T. (195) 287
Rittich, B., Polster, M. and Králik, O.
Reversed-phase HPLC. I. Relationship between R_M values and Hansch's π parameters for a series of phenols (197) 43
Riva, R., see Cotellessa, L. (192) 441
Roberfroid, M., see Malvoisin, E. (186) 81
Roberts, D.J., see Cooke, M. (187) 47, (193) 437
Robinson, J.L., Robinson, W.J., Marshall, M.A., Barnes, A.D., Johnson, K.J. and Salas, D.S.
LSC on Amberlite XAD-2 and other styrene-divinylbenzene adsorbents. I. Development of a solvent eluotropic scale (189) 145
Robinson, J.R. and Chapman, R.A.
Determination of carbofuran residues in potato, onion and turnip: comparative analyses using GC-selected ion monitoring or HPLC (193) 213
Robinson, W.J., see Robinson, J.L. (189) 145
Roboz, J., Suzuki, R. and Rose, E.
MS technique for the determination of N-phosphonoacetyl-L-aspartic acid in serum (181) 195
Rocca, J.L., see Coq, B. (186) 457
Roediger, W.E.W., see Darke, D.J. (181) 449
Roelenbosch, M. van, see Van Roelenbosch, M. (186) 435
Roets, E. and Hoogmartens, J.
TLC of the acid hydrolysis products of nineteen benzodiazepine derivatives (194) 262
Rogiers, V.
GC determination of the fatty acid pattern of red cell membrane plasmalogens in healthy children (182) 27
Rohrschneider, L. and Pelster, E.
Influence of different carrier gases and temperatures on the overall performance of a packed column (186) 249
Roitman, J.N., see Molyneux, R.J. (195) 412
Ronzière, M.C., see Perier, C. (182) 155
Roos, B.A., see Lambert, P.W. (198) 293
Roozen, A.M.P., see Gelsema, W.J. (196) 51
Röper, H. and Heyns, K.
Analysis of N-nitroso compounds. V. Trace analysis of N-nitrosoamino acids: Gas chromatography and spectroscopy of N-nitrosoamino acid methyl esters (in German) (193) 381
Rose, E., see Roboz, J. (181) 195
Roseboom, H., see Hulshoff, A. (186) 535
Rosegay, A., see Walker, R.W. (181) 85
Rosen, M.T., see Holdiness, M.R. (198) 329
Rosenbaum, M., Hančil, V. and Komers, R.
Resolution of overlapping chromatographic peaks by parameter estimation of their model (191) 157
Rosenberg, A., see Leskawa, K.C. (193) 316
Rosenblum, M.G. and Loo, T.L.
HPLC analysis of the anticancer agent methylglyoxal bis(guanylhydrazone) (MGBG, NSC-32946) in biological fluids (183) 363
Rosenmund, H., see Wad, N. (183) 387
Roshchina, T.M., see Gavrilova, T.B. (192) 323
Rosing, D.R., see Jaouni, T.M. (182) 473
Ross, A., see Rucki, R.J. (190) 359
Ross-Lee, L.M., Eadie, M.J., Bochner, F., Hooper, W.D. and Tyrer, J.H.
Electron-capture GC assay for metoclopramide in plasma (183) 175

Rossum, J.M. van, see Van Rossum, J.M. (182) 179
Rotatori, M., see Laganà, A. (198) 143
Roth, M., see Novák, J. (199) 209
Rothbart, H.L., see Barford, R.A. (185) 393
Roumeliotis, P. and Unger, K.K.
Preparative separation of proteins and enzymes in the mean molecular-weight range of 10,000-100,000 on LiChrosorb Diol packing by high-performance size-exclusion chromatography (185) 445
Rout, D.N., see Mahapatra, G.N. (193) 338
Routhier, R., see Seidah, N.G. (193) 291
Routledge, P.A., see Bjornsson, T.D. (181) 417
Rovei, V., Sanjuan, M. and Hrdina, P.D.
Analysis of tricyclic antidepressant drugs by GC using nitrogen-selective detection with packed and capillary columns (182) 349
Rowe, E., see Kullberg, M.P. (187) 264
Rowland, S.P., see Stanonis, D.J. (190) 448
Rowse, K.L., see Oseekey, K.B. (182) 459
Roy, C., Bellemare, G.R. and Chornet, E.
Effect of the sensitivity setting of a katharometer on response factors at low concentrations (197) 121
Różyło, J.K., see Jaroniec, M. (188) 27
Rucki, R.J., Ross, A. and Moros, S.A.
Application of an electrochemical detector to the determination of procarbazine hydrochloride by HPLC (190) 359
Rudolph, J. and Bächmann, K.
Use of isothermal GC for the determination of the adsorption enthalpies and entropies of inorganic halides at high temperatures (187) 319
Ruff, F., see Szókán, G. (198) 207
Ruff, P., see Gracza, L. (193) 486
Rumack, B.H., see Peterson, R.G. (188) 420
Ruyter, M.G.M. de, see De Ruyter, M.G.M. (183) 193
Ryan, D.M., see Miceli, J.N. (183) 250
Ryan, E.A. and Kropinski, A.M.
Separation of amino sugars and related compounds by two-dimensional TLC (195) 127
Ryan, J.J. and Pilon, J.C.
HPLC in the analysis of chorinated dibenzodioxins and dibenzofurans in chicken liver and wood shaving samples (197) 171
Rynkowski, J., see Paryjczak, T. (188) 254
Ryšlavý, Z., see Boček, P. (191) 271

S

Sabbatini, J.Z., see Desiderio, D.M. (195) 369
Sachok, B., Kong, R.C. and Deming, S.N.
Multifactor optimization of reversed-phase LC separations (199) 317
Sachok, B., see Bidlingmeyer, B.A. (186) 419
Sachok, B., see Kong, R.C. (199) 307
Sachse, J.
Determination of perloline in grass species (in German) (192) 199
Sadana, K.L., see Quilliam, M.A. (194) 379, (196) 367
Sadanaga, T., Hikida, K., Tameto, K., Nakanishi, M., Matsushima, Y. and Ohkura, Y.
Determination of carpipramine in plasma by HPLC (183) 246
Sadée, W., see Cohen, M.B. (198) 435
Sadee, W., see Maybaum, J. (188) 149
Sadowski, J., see Witkiewicz, Z. (200) 65
Saeed, T., Sandra, P. and Verzele, M.
Synthesis and properties of a novel chiral stationary phase for the resolution of amino acid enantiomers (186) 611
Sahil, A., see Fritz, D.F. (186) 63
Saint-Jalm, Y. and Moree-Testa, P.
Study of nitrogen-containing compounds in cigarette smoke by GC-MS (198) 188
Saito, M., see Leskawa, K.C. (193) 316
Sakagami, K., see Tanaka, N. (199) 327
Sakodynskii, K.I., Boeva, V.I., Panina, L.I. and Glazunova, L.D.
Polyfunctional polymer sorbents (186) 227
Sakodynskii, K.I., see Yeroshenkova, G.V. (198) 377
Salas, D.S., see Robinson, J.L. (189) 145
Salemink, C.A., see Huibregtse-Minderhout, L. (196) 425
Salmon, D.G., see Adlard, E.R. (186) 207
Salmon, J.R., see Heyes, W.F. (194) 416
Salmona, M., see Gazzotti, G. (188) 400
Salovaara, P., see Widén, C.-J. (188) 213
Salschutz, M., see Hui, K.-S. (192) 341
Salvà, P., see Cotellessa, L. (192) 441
Samperi, R., see Di Corcia, A. (198) 347
Sampson, B. and Barlow, G.B.
Separation of peptides and amino acids by ion-exchange chromatography of their copper complexes (183) 9
Samson, D.M., see Lim, C.K. (185) 605
Samuelson, O., see Jahangir, L.M. (193) 197
Samuelson, O., see Nilsson, B.F. (198) 267
Sánchez, M.G., see Albet, C. (181) 504
Sandra, P., see Saeed, T. (186) 611
Sandra, P., see Verzele, M. (190) 331, (199) 105
Sandt, J.S.F. van der, see Van der Sandt, J.S.F (182) 201
Sanjuan, M., see Rovei, V. (182) 349
Sano, M., see Hakusui, H. (182) 47
Sansare, S.D., see Choudhary, V.R. (192) 420
Santi, D.V., see Nierenberg, D.W. (190) 212
Sarhan, F., Ziegler, J.M., Nicolas, A. and Siest, G.
Rapid method for simultaneous determination of phenobarbital, diphenylhydantoin and their main

hydroxylated metabolites by nitrogen-selective GC (183) 505
Sasaki, H., see Kato, Y. (190) 297, (190) 305, (193) 29, (193) 311, (193) 458
Sasaki, K., see Kodama, H. (183) 226
Sasaki, K., see Mikasa, H. (190) 501
Sass, S. and Ludemann, W.D.
TLC of phosphonic acids (187) 447
Sass, S. and Parker, G.A.
Structure-response relationship of GC-flame photometric detection to some organophosphorus compounds (189) 331
Sassard, J., see Bernard, N. (183) 99
Sassard, J., see Cuisinaud, G. (181) 399
Sasse, F., Hammer, J. and Berlin, J.
Fluorimetric and HPLC determination of harmane alkaloids in *Peganum harmala* cell cultures (194) 234
Satoh, Y., see Suzuki, T. (195) 277
Sattar, M.A. and Paasivirta, J.
TLC of 2-methyl-4-chlorophenoxyacetic acid and its soil metabolites (189) 73
Savelsberg, M., see Vierkorn-Rudolph, B. (186) 219
Sawada, Y., see Kato, Y. (190) 305
Schaaper, W.M.M., Voskamp, D. and Olieman, C.
Perfluoroalkanoic acids as lipophilic ion-pairing reagents in reversed-phase LC of peptides including secretin (195) 181
Schallies, M., see Heller, W. (186) 843
Schaub, P., see Niederberger, W. (182) 454
Scherrmann, J.M., Hoellinger, H., Sonnier, M., Hoffelt, J. and Nguyen-Hoang-Nam
Simple, rapid method for extraction of urinary cannabinoids by LSC (196) 342
Schier, G.M. and Eng Tho Gan, I.
Micro estimation of plasma theophylline by GLC with on-column butylation and nitrogen-specific detection (182) 232
Schill, G., see Melin, A.T. (185) 225
Schley, J., see Kresse, M. (183) 475
Schmidt, K., see Heller, W. (186) 843
Schmuckler, G., see Gjerde, D.T. (186) 509, (187) 35
Schmujlovich, S.M., see Lazaris, A.Y. (198) 337
Schoenmakers, P.J., Billiet, H.A.H. and De Galan, L.
Influence of organic modifiers on the retention behaviour in reversed-phase LC and its consequences for gradient elution (185) 179
Scholten, A.H.M.T., Welling, P.L.M., Brinkman, U.A.T. and Frei, R.W.
Use of PTFE coils in post-column photochemical reactors for LC: application to pharmaceuticals (199) 239
Schott, H.
Preparative isolation of deoxyriboadenylic acids from hydrolysates of oxidized herringsperm DNA using template-chromatography (in German) (187) 119
Schott, H. and Watzlawick, H.
Preparation of oligomers of deoxyriboguanylic acids by partial chemical hydrolysis of DNA (in German) (196) 435
Schott, H., see Dizdaroglu, M. (188) 273
Schröter, H., see Bode, J. (190) 437
Schulz, J.M. and Herrmann, K.
Analysis of hydroxybenzoic and hydroxycinnamic acids in plant material. I. Sample preparation and TLC (195) 85
Schulz, J.M. and Herrmann, K.
Analysis of hydroxybenzoic and hydroxycinnamic acids in plant material. II. Determination by GLC (195) 95
Schumacher, A., Geissler, H.E. and Mutschler, E.
Quantitative determination of diclofenac sodium salt in plasma by absorption spectroscopy with direct quantitation of thin-layer chromatograms (in German) (181) 512
Schüsler-van Hees, M.T.I.W. and Beijersbergen van Henegouwen, G.M.J.
Determination of catecholamines and O-methylated metabolites by reversed-phase HPLC with fluorimetric detection and its application to enzyme kinetics (196) 101
Schwartz, D.P., see White, J.D. (196) 303
Schwartz, R.D., see Mathews, R.G. (186) 183, (199) 97
Schweinsberg, P.D. and Loo, T.L.
Simultaneous analysis of ATP, ADP, AMP, and other purines in human erythrocytes by HPLC (181) 103
Schweizer, A., Willis, J.V., Jack, D.B. and Kendall, M.J.
Determination of total monohydroxylated metabolites of diclofenac in urine by electron-capture GLC (195) 421
Schwenn, J.D. and Jender, H.G.
Reversed-phase HPLC of adenine nucleotides: application to the kinetics of an adenosine 3'-phosphate 5'-sulphatophosphate sulphotransferase from plants (193) 285
Schwertfeger, G., see Vogt, W. (199) 191
Scott, A.I., see Nordlöv, H. (190) 221
Scott, K.F., see Horrocks, J.P. (197) 109
Scott, R.P.W. and Kucera, P.
Use of microbore columns for the separation of substances of biological origin (185) 27
Scott, R.P.W., Kucera, P. and Munroe, M.
Use of microbore columns for rapid LC separations (186) 475
Scott, R.P.W. and Simpson, C.F.
Solute-solvent interactions on the surface of reversed phases. I. Stationary phase interactions and their dependence on bonding characteristics (197) 11
Scott, R.P.W. and Traiman, S.
Solute-solvent interactions on the surface of silica gel. III. Multilayer adsorption of water on the surface of silica gel (196) 193
Scovill, J.P., see Brown, N.D. (195) 146, (200) 267
Seale, T., see Chan, W.Y. (181) 259

Seamark, D.A., Trafford, D.J.H., Hiscocks, P.G. and Makin, H.L.J.
HPLC of vitamin D: enhanced UV absorbance by prior conversion to isotachysterol derivatives (197) 271
Sebille, B., Thuaud, N. and Tillement, J.P.
Usefulness of chromatographic methods for the determination of drug-protein binding parameters (193) 522
Seccia, M., see Bernard, N. (183) 99
Seccia, M., see Cuisinaud, G. (181) 399
Segall, H.J., see Dimenna, G.P. (192) 474
Seib, P.A., see Mauro, D. (187) 421
Seidah, N.G., Routhier, R., Benjannet, S., Larivière, N., Gossard, F. and Chrétien, M.
Reversed-phase HPLC purification and characterization of the adrenocorticotropin/lipotropin precursor and its fragments (193) 291
Seifert, K. and Kraus, G.
Solution behaviour of liquid crystal phases in capillary GC. II. Solution properties of smectic A-, B- and C-modifications (in German) (192) 97
Seitz, D.E., see Lindner, W. (185) 323
Seiwell, R. and Brater, C.
Separation and analysis of azosemide in urine and in serum by HPLC (182) 257
Selkirk, A.B., see Moody, R.R. (182) 359
Semanský, M., see Tichá, M. (194) 183
Sen, G., Mulchandani, N.B. and Patankar, A.V.
Reversed-phase HPLC separation and analysis of some physalins (13,14-*seco*-16,24-cyclo-steroids) (198) 203
Sepaniak, M.J. and Yeung, E.S.
Determination of adriamycin and daunorubicin in urine by HPLC with laser fluorometric detection (190) 377
Seppä, E.-L., see Pyysalo, H. (190) 466
Sereda, I.P., see Mikhaylenko, V.P. (191) 231
Serpinet, J., see Morel, D. (200) 95
Seta, K., Washitake, M., Anmo, T., Takai, N. and Okuyama, T.
High-performance anion-exchange chromatography of UV-absorbing constituents of human urine (181) 311
Settine, R.L., see Barnes, S. (183) 269
Settlage, J.A., see Jennings, W. (186) 189
Ševčík, J.
Information content of multi-dimensional switching systems in GC (186) 129
Severini, C., see Di Corcia, A. (198) 347
Sewell, P., see Martin, A.J.P. (186) 15
Shabtai, J., see Klemm, L.H. (198) 1
Shaikh, B., Hallmark, M.R., Hallmark, R.K., Manning, W.B., Pinnock, A. and Kawalek, J.C.
Separation and detection of 2-aminoanthracene and its metabolites by HPLC (195) 392
Shaikh, N.A., see Kuksis, A. (182) 1
Shaligin, G.F., see Berezkin, V.G. (193) 132
Shand, D.G., see Bjornsson, T.D. (181) 417
Sharifi-Sandjani, N., see Tirgan, M.R. (193) 397
Sharma, O.P. and Makkar, H.P.S.
TLC of lantadene A and some related triterpenoids (196) 515
Sharma, S.K. and Mahendroo, P.P.
Affinity chromatography of cells and cell membranes (Review) (184) 471
Shatkay, A.
Comparison of different methods of quantitative analysis employing GLC, illustrated by the determination of water in organic solvents (198) 7
Shatz, V.D., Belikov, V.A., Zelchan, G.I., Solomennikova, I.I., Yerchak, N.P., Pudova, O.A. and Lukevics, E.
Chromatography of organometallic and organometalloidal derivatives of amino alcohols. II. GC evidence for transannular bonds in solutes (200) 105
Shaw, D.C. and West, C.E.
The isolation of methionine and ethionine by silver ligand chromatography and application to methionine-containing peptides (200) 185
Shay, C., see Joy, K.W. (194) 76
Shaywitz, B.A., see Anderson, G.M. (181) 453
Shearer, M.J., see Haroon, Y. (200) 293
Shepherd, A.M.M., see Haegele, K.D. (187) 171
Shepherd, M.J., see Gilbert, J. (193) 235, (197) 71
Shibamoto, T., Harada, K., Yamaguchi, K. and Aitoku, A.
GC retention characteristics of different coating thicknesses on a glass capillary column (194) 277
Shibata, T., see Inayama, S. (194) 85
Shibata, T., see Marunaka, T. (183) 331
Shibusawa, Y., see Matsumoto, U. (187) 351
Shida, T., see Mita, H. (181) 153
Shikama, K., see Suzuki, T. (195) 277
Shimada, K. and Nagase, Y.
Quantitative HPLC determinations of aminopyrine and its metabolites in man (181) 51
Shimada, K., see Iwasaki, H. (198) 429
Shimamori, Y., see Nakano, N.I. (188) 347
Shimono, T., Takagi, H., Isobe, T. and Tarutani, T.
Gel permeation chromatographic separation and identification of trimethylsilyl derivatives of silicate anions (197) 59
Shinozawa, S., Mimaki, Y., Araki, Y. and Oda, T.
Determination of the concentration of adriamycin and its metabolites in the serum and tissues of Ehrlich carcinoma-bearing mice by HPLC (196) 463
Shinozawa, S., Mimaki, Y., Tomano, H., Araki, Y. and Oda, T.
Determination of adriamycin in liposomes by HPLC using a fluorescence detector (190) 489
Shiono, S., Karino, I., Ishimura, A. and Enomoto, J.
Separation and characterization of by-

product oligomers in epoxy resins by reversed-phase HPLC (193) 243
Shiu, G.K. and Goehl, T.J.
High-performance paired-ion LC determination of bleomycin A_2 in urine (181) 127
Shmidel, Y.B., see Berezkin, V.G. (191) 95
Shono, T., see Mizobuchi, Y. (194) 153
Shopova, B.I. and Mladenov, I.T.
Separation of some azo-pigments from their diazo- and coupling components by chromatography on Sephadex LH-20 with N,N-dimethylformamide as eluent (187) 232
Siciliano, J., see Heisler, E.G. (194) 89
Siest, G., see Sarhan, F. (183) 505
Sievers, R.E., Phillips, M.P., Barkley, R.M., Wizner, M.A., Bollinger, M.J., Hutte, R.S. and Fehsenfeld, F.C.
Selective electron-capture sensitization (186) 3
Sievers, S., see König, W.A. (200) 189
Silva, J.A.F. de, see De Silva, J.A.F. (181) 272, (183) 411
Silverman, R.B. and Dolphin, D.
TLC of acid-labile cobalamins (194) 273
Simard, J.-L., see Garneau, F.-X. (188) 445
Simic, M.G., see Dizdaroglu, M. (188) 273, (195) 119
Simmonds, P.G. and Kerns, E.
Direct aqueous injection GC for the analysis of trace organics in water (186) 785
Simon, Á., see Bartik, T. (197) 92
Simon, C., see Miquelis, R. (189) 101
Šímová, A., see Krupčík, J. (191) 207
Simpson, C.F., see Scott, R.P.W. (197) 11
Singh, U.S. and Tripathi, R.K.
Estimation of the systemic fungicide Ridomil by TLC (200) 317
Singhawangcha, S., Poole, C.F. and Zlatkis, A.
The determination of bifunctional compounds. IX. A selective reaction for the determination of guaifenesin in plasma by GC (183) 433
Singhawangcha, S., see Poole, C.F. (186) 307, (187) 331, (199) 123
Sinibaldi, M., see Corradini, D. (187) 458
Siouffi, A.M., Wawrzynowicz, T., Bressolle, F. and Guiochon, G.
Problems and applications of reversed-phase TLC (186) 563
Sioufi, A., Colussi, D. and Marfil, F.
GC determination of valproic acid in human plasma (182) 241
Sioufi, A. and Pommier, F.
GC determination of low concentrations of benzoic acid in human plasma and urine (181) 161
Sioufi, A. and Pommier, F.
GC determination of amantadine hydrochloride (Symmetrel) in human plasma and urine (183) 33
Siraj, M.Y., see Chan, P.K. (194) 387
Sirimanne, P., see Cheetham, N.W.H. (196) 171
Sisco, W.R., see Gilpin, R.K. (194) 285
Siu, K.W.M. and Aue, W.A.
Binary liquid phase system for GLC with *in situ* variable polarity (189) 255
Siu, K.W.M. and Aue, W.A.
Effect of dopants on the linearity of a d.c. electron-capture detector (192) 413
Sjöquist, B. and Magnuson, E.
Analysis of salsolinol and salsoline in biological samples using deuterium-labelled internal standards and GC-MS (183) 17
Sjövall, J., see Axelson, M. (186) 725
Skellern, G.G., see Knight, B.I. (192) 247
Skellern, G.G., see Marcantonio, L.A. (183) 118
Skews, H., see Jackman, G.P. (182) 277
Skrdlant, H.B., see Haegele, K.D. (187) 171
Šlais, K. and Šubert, J.
Determination of *cis*- and *trans*-isomers of dosulepine and dithiadene by HPLC (191) 137
Šlais, K., see Krejčí, M. (191) 3
Slater, T.F., see Reddrop, C.J. (192) 375, (193) 7
Śledzińska, B., see Gazda, K. (191) 9
Sleeman, H.K., see Brown, N.D. (195) 146, (200) 267
Sliwinski, B.J., see Barford, R.A. (185) 393
Smalley, A.W., see Thompson, D.P. (187) 399
Šmejkal, F., Popl, M., Číhová, A. and Zázvorková, M.
Reversed-phase LC of aromatic nitro compounds on a macroporous polystyrene gel (197) 147
Smith, B.J., Toogood, C.I.A. and Johns, E.W.
Quantitative staining of submicrogram amounts of histone and high-mobility group proteins on sodium dodecylsulphate-polyacrylamide gels (200) 200
Smith, E.C., Jones, A.D. and Hammond, E.W.
Investigation of the use of argentation HPLC for the analysis of triglycerides (188) 205
Smith, G., see Clements, J.A. (189) 272
Smith, H.W., see Friedman, Z. (182) 414
Smith, J.L., see Stocklinski, A.W. (195) 281
Smith, J.S. and Hanrahan, J.M.
Adsorption-derivatization GC-MS determination of aqueous 1,6-hexanediamine at the low and sub-ppm level (193) 271
Smith, K.J. and Meffin, P.J.
Mexiletine analysis in blood and plasma using GC and nitrogen-selective detection (181) 469
Smith, K.J., see Meffin, P.J. (183) 352
Smith, L.A., see Lin, S.-N. (197) 31
Smith, M.D.
Rapid method for determination of riboflavin in urine by HPLC (182) 285
Smith, R.A., see Stea, B. (188) 363
Smith, R.M., Doran, D., Mazur, M. and Bush, B.
HPLC determination of protoporphyrin and zinc protoporphyrin in blood (181) 319
Smith, S.J., see Holzer, G. (194) 410
Smith, S.L., Jorgenson, J.W. and Novotný, M.
Applications of reversed-phase chromatography and nephelometric

detection to analysis of non-polar mixtures at microgram level (187) 111
Smith, W.J., see Johnston, M.A. (189) 241
Smolková, E., Zima, J., Dousek, F.P., Jansta, J. and Plzák, Z.
A PTFE-based carbon adsorbent in HPLC (191) 61
Smolková, E., see Řehák, V. (191) 71
Smolková-Keulemansová, E. and Krýsl, S.
Inclusion compounds in chromatography (Review) (184) 347
Smolková-Keulemansová, E., see Procházka, M. (189) 25
Smrž, M., see Macek, K. (193) 421
Smrž, M., see Mikeš, O. (192) 159
Smulders, C.F.A., see Hulshoff, A. (181) 363
Snyder, L.R. and Dolan, J.W.
Coated-open-tubular chromatography with flow segmentation. I. Theory (185) 43
Snyder, L.R. and Poppe, H.
Mechanism of solute retention in liquid-solid chromatography and the role of the mobile phase in affecting separation. Competition *versus* "sorption" (Review) (184) 363
Snyder, L.R., see Dolan, J.W. (185) 57
Snyder, L.R., see Gant, J.R. (185) 153
Snyder, R.S., see Bosisio, A.B. (189) 317
Snyder, R.S., see McGuire, J.K. (194) 323
Sobel, B.E., see Gross, R.W. (197) 79
Sobel, M. and Mutschler, E.
Fluorimetric determination of xipamide in body fluids with a new fluorescent reagent (in German) (183) 124
Soegtrop, H.H., see Brodie, R.R. (182) 379
Sohda, S., see Tamegai, T. (193) 483
Soják, L. and Krupčík, J.
Use of Carbowax 20M glass capillary columns below their lower temperature limit (190) 283
Soják, L., Krupčík, J. and Janák, J.
Comparison of capillary columns coated with C_{87} hydrocarbon and squalane in the analysis of *n*-pentadecene isomers (191) 199
Soják, L., Krupčík, J. and Janák, J.
GC of all C_{15}-C_{18} linear alkenes on capillary columns with very high resolution power (195) 43
Soják, L., Ostrovský, I., Leclercq, P.A. and Rijks, J.A.
Identification of *n*-hepta- and *n*-octadienes by high-resolution GC using structure-retention correlations and MS (191) 187
Soják, L., see Dubois, J.-E. (194) 121
Sokolowski, A. and Wahlund, K.-G.
Peak tailing and retention behaviour of tricyclic antidepressant amines and related hydrophobic ammonium compounds in reversed-phase ion-pair LC on alkyl-bonded phases (189) 299
Solomennikova, I.I., see Shatz, V.D. (200) 105
Someno, T., Katoh, K., Niijima, K. and Miyazaki, H.
Rapid method for the determination of urokinase activity by HPLC (188) 185
Sonnier, M., see Scherrmann, J.M. (196) 342
Sörbo, B., see Hannestad, U. (200) 171
Sørensen, H., see Helboe, P. (197) 199
Sorokina, E.J., see Berezkin, V.G. (193) 132
Sorrell, R.K. and Reding, R.
Analysis of polynuclear aromatic hydrocarbons in environmental waters by HPLC (185) 655
Sosa, J.M., Hernández, R. and McBrayer, G.
Crosslinked polymethylphenylsiloxanes as adsorbents for GC (189) 169
Souter, R.W.
GC quantitation of residual impurities in raw material matrixes (193) 207
Špaček, P., see Vozka, S. (193) 128
Speck, M. and Dirr, E.
GC determination of Metalaxyl (Ridomil) residues in tobacco (200) 313
Spencer, P.S., see Zlatkis, A. (182) 137
Spiker, S.
Slab gel designed for enzymatic digestion of proteins in polyacrylamide gel slices and direct resolution of peptides (198) 169
Spiteller, G., see Ende, M. (183) 1
Spitz, H.D.
Determination of 2,6-di-(*tert.* -butyl)-4-methylphenol in rubber-base materials and identification of an oxidation product from 2,6-di-(*tert.* -butyl)-4-methylphenol (190) 193
Squire, K.M., see Glajch, J.L. (199) 57
Srisukh, D., see Haken, J.K. (190) 291, (199) 199
Srisukh, D., see Wainwright, M.S. (188) 246, (198) 156
Srivastava, S.P., Dua, V.K. and Chauhan, L.S.
Chromatographic behaviour of aliphatic amines on phenol-impregnated thin layers (196) 225
Stahl, E. and Müller, J.
New spreader for the preparation of gradient layers in TLC (189) 293
Stan, H.-J. and Abraham, B.
Determination of residues of anabolic drugs in meat by GC-MS (195) 231
Stankoviansky, S., see Kaniansky, D. (194) 11
Stanley, P.E., see Reece, P.A. (183) 109
Stanonis, D.J. and Rowland, S.P.
Distribution ratios and elution behavior of carbamates and polyhydroxy compounds from Sephadex G-15 (190) 448
Starratt, A.N. and Stevens, M.E.
Ion-pair HPLC of the insect neuropeptide proctolin and some analogs (194) 421
Startin, J.R., see Gilbert, J. (189) 86, (197) 71
Stea, B., Halpern, R.M., Halpern, B.C. and Smith, R.A.
Quantitative determination of pterins in biological fluids by HPLC (188) 363
Stebbins, R.G., see Grimsrud, E.P. (197) 51
Stein, J.L., see Desiderio, D.M. (195) 369
Sternson, L.A., see Golander, Y. (181) 41
Sternson, L.A., see Marsh, K.C. (187) 101
Sternson, L.A., see Musson, D.G. (188) 159
Stetson, P., see Patterson, E. (181) 33
Stevens, M.E., see Starratt, A.N. (194) 421

Stewart, J.G. and Williams, P.A.
Formulation of multicomponent mobile solvents for LC (198) 489
Stewart, J.T., see Honigberg, I.L. (181) 266
Sticher, O., see Hostettmann, K. (186) 529
Stierle, U., see Liebich, H.M. (199) 181
Stinson, E.E., see Heisler, E.G. (194) 89
Stocklinski, A.W., Williams, H., Hale, W.B. and Smith, J.L.
UV-induced silica gel GF photoluminescence studied by liquid scintillation spectrophotometry (195) 281
Stockmeyer, M., see Gfeller, J.C. (198) 162
Stockton, R.A., see Brinckman, F.E. (191) 31
Storey, G.C.A., see Flanagan, R.J. (187) 391
Stoveken, J., see Melander, W. (199) 35
Stoveken, J., see Melander, W.R. (185) 111
Stranieri, A.M., see Downing, D.T. (192) 208
Straus, A.H., see Bianchini, P. (196) 455
Streeter, A.J., see Facchini, V. (187) 218
Strife, R.J., Jardine, I. and Colvin, M.
Analysis of the anticancer drugs VP 16-213 and VM 26 and their metabolites by HPLC (182) 211
Strojny, N. and De Silva, J.A.F.
Determination of mecillinam in urine by reversed-phase HPLC (181) 272
Strong, J.M., see Branfman, A.R. (182) 163
Štrop, P., see Brynda, E. (196) 39
Štrop, P., see Mikeš, O. (192) 159
Stubbles, J.A., see Bowker, M.J. (200) 261
Studebaker, J.F.
Solid-phase reagents for LC detection (185) 497
Štulík, K. and Pacáková, V.
Electrochemical detector for HPLC (192) 135
Šubert, J., see Šlais, K. (191) 137
Sugawara, Y., see Suzuki, T. (195) 277
Sugden, D., see Gault, M.H. (182) 465
Sugden, K., Hunter, C. and Lloyd-Jones, G.
Ligand-exchange chromatography. I. Resolution of L- and D-proline on a copper(II)-proline complex bound to microparticulate silica gel (192) 228
Sugiyama, S., see Onishi, S. (182) 105
Sullivan, J.B. Jr., see Peterson, R.G. (188) 420
Sundaram, K.M.S., Szeto, S.Y. and Hindle, R.
Detection of aminocarb and its major metabolites by TLC (194) 100
Sundaram, K.M.S., see Szeto, S.Y. (200) 179
Suyama, K., see Ishida, M. (189) 421
Suzuki, A., Noda, K. and Noguchi, H.
HPLC determination of ceftizoxime, a new cephalosporin antibiotic, in rat serum, bile and urine (182) 448
Suzuki, I., see Iwakawa, J. (193) 333
Suzuki, K., see Ujimoto, K. (190) 161
Suzuki, R., see Roboz, J. (181) 195
Suzuki, S., see Onodera, S. (200) 137
Suzuki, S., see Watabe, K. (192) 89
Suzuki, T., Kawada, T. and Iwai, K.
Effective separation of capsaicin and its analogues by reversed-phase HPLC (198) 217
Suzuki, T., Sugawara, Y., Satoh, Y. and Shikama, K.
Human oxymyoglobin: isolation and characterization (195) 277
Suzuki, W., see Hakusui, H. (182) 47
Švec, F., see Lukás, J. (194) 297
Svenneby, G., see Lenda, K. (198) 516
Svoboda, L., see Jandera, P. (192) 37
Svojtková, E., Macek, K., Deyl, Z. and Adam, M.
Determination of specific urinary hydroxyproline-containing peptides in patients with lung cancer (183) 221
Swoboda, W., see Herkner, K. (197) 232
Sye, W.-F., see Poole, C.F. (187) 331, (199) 123
Szabó, A. and Karácsony, E.M.
Detection and assay of dihydroergot alkaloids by TLC using *o*-phthalaldehyde-sulphuric acid reagent (193) 500
Szafranek, J. and Wiśniewski, A.
GLC and HPLC analyses of the acid-catalyzed dehydration reaction of xylitol (187) 131
Szafranek, J., see Błotny, G. (193) 61
Szárföldi-Szalma, I., see Vigh, G. (193) 432
Szász, G., see Papp, O. (191) 245
Szathmáry, C., Galácz, J., Vida, L. and Alexander, G.
Capillary GC-MS determination of some mycotoxins causing fusariotoxicoses in animals (191) 327
Szczepaniak, W., see Nawrocki, J. (188) 323
Szczepanik-van Leeuwen, P.A., see Huang, C.T.L. (196) 150
Székely, G. and Baumgartner, P.
TLC of slightly soluble substances at elevated temperature (186) 575
Szepesi, G., Gazdag, M. and Terdy, L.
Separation of ergotoxine alkaloids by HPLC on silica (191) 101
Szeto, S.Y. and Sundaram, K.M.S.
Simplified method for the analysis of some carbamate insecticides in foliage, forest soil and fish tissue by direct GLC (200) 179
Szeto, S.Y., see Sundaram, K.M.S. (194) 100
Szókán, G., Ruff, F. and Kucsman, Á.
HPLC of diastereomeric sulphoxides and sulphilimines (198) 207
Szokolay, A.M.
HPLC determination of impurities in commercial saccharin (187) 249
Szulc, J., see Witkiewicz, Z. (200) 65
Szymanowicz, A.G., Poulin, G., Fontaine, N., Werquin, J.P. and Borel, J.P.
Improved method for the preparation of 3-hydroxyproline, and some of its chromatographic properties (190) 457

T

Tabata, M., see Onodera, S. (200) 137
Taguchi, V.Y., see Mamer, O.A. (182) 221
Takács, J.M., see Egri, L. (198) 85
Takagi, H., see Shimono, T. (197) 59
Takahashi, H.K., see Bianchini, P. (196) 455
Takahashi, N., see Barendse, G.W.M. (198) 449
Takai, N., see Oshima, R. (192) 452
Takai, N., see Seta, K. (181) 311
Takayama, Y. and Kawai, S.
Improved GC of amygdalin and its diastereomer (197) 240
Takei, S., see Naito, K. (190) 21
Takeo, K., see Nakamura, K. (192) 351, (196) 85
Takeuchi, T. and Ishii, D.
Ultra-micro HPLC (190) 150
Takeuchi, T., Matsuoka, K., Watanabe, Y. and Ishii, D.
Studies on open-tubular microcapillary LC. VI. Styrene-divinylbenzene copolymer stationary phase (192) 127
Takeuchi, T., see Ishii, D. (185) 73
Talseth, T., see Haegele, K.D. (187) 171
Tamegai, T., Sohda, S., Hirose, N., Ohmae, M. and Kawabe, K.
Separation of the diastereoisomers of 1-(4-hydroxy-3-methoxymethylphenyl)-2-N-(*p*-methoxyphenylisopropyl)aminoethanol by HPLC (193) 483
Tameto, K., see Sadanaga, T. (183) 246
Tamura, Z., see Nakamura, H. (200) 324
Tan, Y., see Van Loenhout, J.W.A. (182) 487
Tanabe, H., see Ishibashi, T. (195) 416
Tanács, B., see Tóth, G. (189) 433
Tanaka, A., Nose, N. and Watanabe, A.
GC determination of nitrite in foods as trimethylsilyl derivative of 1*H*-benzotriazole (194) 21
Tanaka, H., see Miyazaki, S. (181) 177
Tanaka, K. and Ishizuka, T.
Ion-exclusion chromatography of condensed phosphates on a cation-exchange resin (190) 77
Tanaka, K., see Ramsdell, H.S. (181) 90
Tanaka, M., see Kobayashi, K. (187) 413
Tanaka, M., see Kono, A. (182) 125
Tanaka, M., see Mizobuchi, Y. (194) 153
Tanaka, N., Sakagami, K. and Araki, M.
Effect of alkyl chain length of the stationary phase on retention and selectivity in reversed-phase LC. Participation of solvent molecules in the stationary phase (199) 327
Tangendjaja, B., Buckle, K.A. and Wootton, M.
Analysis of phytic acid by HPLC (197) 274
Tangerman, A., see Jansen, P.L.M. (182) 100
Tanimitsu, N., see Honda, S. (194) 191
Tanzer, M.L., see Black, C. (190) 393
Tarascio, C.M., see Gal, J. (181) 123
Tarbit, I.F., Richardson, J.P. and Dale, G.
Rapid automated ion-exchange analysis of plasma tyrosine and phenylalanine with data print-out (181) 337
Tarutani, T., see Iwasaki, H. (198) 429
Tarutani, T., see Shimono, T. (197) 59
Taub, D., see Walker, R.W. (181) 85
Taylor, A.A., see Barrow, S.E. (181) 219
Taylor, G.A., Dady, P.J. and Harrap, K.R.
Quantitative HPLC of nucleosides and bases in human plasma (183) 421
Taylor, R., Pragnell, R.J. and McLaren, J.V.
Separation of urea-formaldehyde addition products by gel permeation chromatography (195) 154
Taylor, R.B., see Moody, R.R. (182) 359
Taylor, R.F. and Gaudio, L.A.
HPLC of cancer chemotherapeutic agents: bis(substituted aminoalkylamino)anthraquinones (187) 212
Tenku, A., see Kakita, K. (181) 248
Teplý, J. and Dressler, M.
Direct determination of organic compounds in water using steam-solid chromatography (191) 221
Terdy, L., see Szepesi, G. (191) 101
Terweij-Groen, C.P., Heemstra, S. and Kraak, J.C.
Rapid determination of indomethacin and salicylic acid in serum by means of reversed-phase LC (181) 385
Tesařík, K.
Preparation of glass capillary columns for LC (191) 25
Tesařík, K., see Kahle, V. (191) 121
Tesařík, K., see Krejčí, M. (191) 17
Tesařová, E., see Dufek, P. (191) 115
Tetu, T., see Uden, P.C. (196) 403
Thach, A.B., see James, G.T. (195) 287
Thibault, J.-F.
Separation of α-D-galacturonic acid oligomers by chromatography on polyacrylamide gel (194) 315
Thijssen, H.H.W.
Analysis of isoxazolyl penicillins and their metabolites in body fluids by HPLC (183) 339
Thomas, A.H. and Thomas, J.M.
Use of the image analyser Optomax for the quantitative evaluation of antibiotics separated by gel electrophoresis and by TLC (195) 297
Thomas, J., see Nichol, H. (182) 191
Thomas, J., see Tsuei, S.E. (181) 135, (182) 55
Thomas, J.M., see Thomas, A.H. (195) 297
Thompson, D.P. and Smalley, A.W.
Separation of fluorescent compounds in members of the genus *Rhizopus* by TLC (187) 399
Thomsen, M., see Helboe, P. (189) 249
Thomson, I. and Anderson, R.A.
Determination of cyanide and thiocyanate in biological fluids by GC-MS (188) 357
Thorbecke, G.J., see Ratech, H. (183) 499
Thuaud, N., see Sebille, B. (193) 522

Thurman, E.M.
Conductimetric detection of organic acids in reversed-phase chromatography (185) 625
Thys, C.P., see Tjaden, U.R. (181) 227
Tibbo, N., see Gault, M.H. (182) 465
Tichá, M., Barthová, J., Labský, J. and Semanský, M.
Determination of the interaction of lactate dehydrogenase with high-molecular-weight derivatives of AMP by affinity electrophoresis (194) 183
Tichá, M., see Čeřovský, V. (194) 175
Tichá, M., see Turková, R. (192) 408
Tickell, S.M., see Howard, A.G. (189) 436
Tieckelmann, H., see Lee, G.J.-L. (195) 402
Tiedje, J.M., see Kaspar, H.F. (193) 142
Till, A., see Garzó, G. (191) 253
Tillement, J.P., see Sebille, B. (193) 522
Tillman, J., see Burns, D.T. (190) 141
Timbrell, J.A.
Stability of hydralazine pyruvate hydrazone (198) 150
Timbrell, J.A., see Facchini, V. (183) 167, (187) 218
Tindall, G.W. and Wininger, P.E.
GC–MS method for identifying and determining polychlorinated biphenyls (196) 109
Ting, H.-H. and Quick, M.P.
Simple TLC method for detection of pentachlorophenol in sawdust and woodshavings (195) 441
Tipps, R.W., see McGuire, J.K. (194) 323
Tirgan, M.R. and Sharifi-Sandjani, N.
GC study of methylstyrenes in the light-oil fraction of coal distillate using selective hydrogenation (193) 397
Tishbee, A. and Kirson, I.
HPLC separation and analysis of steroidal constituents of two solanaceous plants (195) 425
Tjaden, U.R., Meeles, M.T.H.A., Thys, C.P. and Van der Kaay, M.
Determination of some benzodiazepines and metabolites in serum, urine and saliva by HPLC (181) 227
Tolan, J.W., Eskola, P., Fink, D.W., Mrozik, H. and Zimmerman, L.A.
Determination of avermectins in plasma at nanogram levels using HPLC with fluorescence detection (190) 367
Tomano, H., see Shinozawa, S. (190) 489
Tomlinson, E., see Riley, C.M. (185) 197
Tomori, É., Orbán, E. and Maderspach, A.
Capillary GC studies of cholesterol biosynthesis in rats treated with EGYT-1299 (191) 261
Tomori, E., see Maume, B.F. (186) 581
Tomori, É., see Patthy, M. (191) 145
Tompson, J.G., see Agris, P.F. (194) 205
Tomsová, Z., Gregorová, I. and Horký, K.
Separation of urinary steroids by GLC on packed and mixed-bed columns (200) 221
Tonelli, D., see Gattavecchia, E. (193) 340
Tonnaer, J.A.D.M., Verhoef, J., Wiegant, V.M. and De Jong, W.
Separation and quantification of angiotensins and some related peptides by HPLC (183) 303
Toogood, C.I.A., see Smith, B.J. (200) 200
Torchia, M.G. and Danzinger, R.G.
Reversed-phase HPLC assay for cefoxitin in proteinaceous biological samples (181) 120
Tornabene, T.G., see Holzer, G. (186) 795
Torres, J., see Mathews, R.G. (186) 183, (199) 97
Torri, G., see Oreste, P. (195) 398
Toste, A.P.
HPLC of myosin's subfragment 1 and light chains on Spherogel TSK-type columns (197) 207
Tóth, G., Mucha, I. and Tanács, B.
Adsorption chromatographic separation of ^{125}I-labelled prostaglandin $F_{2\alpha}$ and prostaglandin E_2 tyrosine methyl ester (189) 433
Tourn, M.L., see Buffa, M. (200) 309
Toussaint, M.W., see Pandey, R.C. (198) 407
Towers, G.H.N., see Picman, A.K. (189) 187
Trafford, D.J.H., see Seamark, D.A. (197) 271
Traiman, S., see Scott, R.P.W. (196) 193
Traynor, S.G., see Marcelin, G. (187) 57
Treppendahl, S. and Jakobsen, P.
Isolation of α- and β-peltatin and podophyllotoxin by LC and analysis by HPLC (189) 276
Tripathi, R.K., see Singh, U.S. (200) 317
Tříska, J., see Kuraš, M. (191) 319
Troost, J.J., see Verpoorte, R. (184) 79
Tsibul'skaya, I.A., see Ioffe, B.V. (186) 851
Tsuchiya, H., see Miyazaki, S. (181) 177
Tsuda, T. and Nakagawa, G.
Open-tubular microcapillary LC with 30-40 μm I.D. columns (199) 249
Tsuda, T., see Hibi, K. (189) 179
Tsuda, T., see Ishii, D. (185) 73
Tsuei, S.E., Thomas, J. and Moore, R.G.
Quantification of oxprenolol in biological fluids using HPLC (181) 135
Tsuei, S.E., Thomas, J. and Nation, R.L.
Simultaneous quantitation of chlormethiazole and two of its metabolites in blood and plasma by GLC (182) 55
Tsukida, K., Masahara, R. and Ito, M.
HPLC analysis of *cis-trans* stereoisomeric 3-dehydroretinals in the presence of retinal isomers (192) 395
Tsutsui, M., see Uobe, K. (193) 83
Tsuzuki, S., see Nakamura, H. (200) 324
Tumlinson, J.H., see Heath, R.R. (189) 199
Turchetto, L., see Quercia, V. (193) 163
Tureček, F., see Julák, J. (190) 183
Turková, J., see Čeřovský, V. (194) 175
Turková, R., Tichá, M. and Kocourek, J.
Studies on lectins. XLV. Effect of ionic strength on the interaction of lectins with carbohydrates as determined by affinity electrophoresis (192) 408
Turner, L.K., see Perkal, M. (196) 180

Tusell, J.M. and Gelpi, E.
Prostaglandins E and F, and 19-hydroxylated E and F (series I and II) in semen of fertile men. GC and LC separation with selected ion detection (181) 295
Twomey, T.M., Bartolucci, S.R. and Hobbs, D.C.
Analysis of piroxicam in plasma by HPLC (183) 104
Tyihák, E., see János, É. (191) 239
Tyihák, E., see Mincsovics, E. (191) 293
Tyler, R.C., see Morgan, E.D. (186) 605
Tyrer, J.H., see Ross-Lee, L.M. (183) 175
Tyson, R.W. and Wickstrom, E.
Rapid purification of radiolabeled trinucleoside diphosphates by reversed-phase HPLC (192) 485

U

Ubuka, T., Kinuta, M., Akagi, R. and Kiguchi, S.
Separation of S-sulphocysteine and related compounds by anion-exchange chromatography and electrophoresis (188) 442
Uden, P.C., Henderson, D.E., DiSanzo, F.P., Lloyd, R.J. and Tetu, T.
Applications of Dexsil and other GC porous-layer open tubular columns to metal compound separations and pyrolysis (196) 403
Uden, P.C., see Estes, S.A. (196) 265
Ujimoto, K., Ando, I., Yoshimura, T., Suzuki, K. and Kurihara, H.
Gel chromatographic separations of monomeric oxo anions of phosphorus on Sephadex G-10 and Bio-Gel P-2 (190) 161
Ulrych, S., see Kramlová, M. (193) 515
Umeno, Y., see Marunaka, T. (183) 331, (188) 270, (190) 107
Unger, K.K., see Jost, W. (185) 403
Unger, K.K., see Roumeliotis, P. (185) 445
Unger, P.D. and Friedman, M.A.
HPLC of caprolactam and its metabolites in urine and plasma (187) 429
Untz, G., see Bauer, M. (192) 479
Uobe, K., Nishida, K., Inoue, H. and Tsutsui, M.
Thermal degradation of sugars in a GC injection port (193) 83
Upadhyaya, J.S. and Upadhyaya, S.K.
TLC separation of potential antineoplastic agents: 1-ethoxycarbonyl-2-arylazo-2-nitroethanes (198) 224
Upadhyaya, S.K., see Upadhyaya, J.S. (198) 224
Uppal, G.S., see Brown, P.M. (190) 201
Upton, R.A., Buskin, J.N., Guentert, T.W., Williams, R.L. and Riegelman, S.
Convenient and sensitive HPLC assay for ketoprofen, naproxen and other allied drugs in plasma or urine (190) 119
Upton, R.A., see Guentert, T.W. (183) 514
Uralets, V.P., Rijks, J.A. and Leclercq, P.A.
syn-anti Isomerisation of 2,4-dinitrophenylhydrazones of volatile carbonyl compounds in capillary GC-MS analyses (194) 135

V

Vacík, J., see Gaš, B. (192) 253
Vakhrusheva, N.M., see Evtushenko, E.V. (196) 331
Valkó, K., see Papp, O. (191) 245
Valles, E.M., see Damiani, D.E. (196) 355
Van Boven, M., Daenens, P. and Vandereycken, G.
Determination of nitromethaqualone in blood by electron-capture-GC (182) 435
Van Damme, J.-C. and Galoux, M.
HPLC analysis of formulations of phenoxycarboxylic acids, dicamba, ioxynil and bromoxynil mixtures (in French) (190) 401
Van de Laar, G.L.M., see Paanakker, J.E. (183) 459
Van de Lagemaat, D., see Van der Poll, J.M. (197) 263
Van de Venne, J.L.M., see Deelder, R.S. (185) 241
Van de Werken, P.H., see Barendse, G.W.M. (198) 449
Van den Bosch, N. and De Vos, D.
Some aspects of the GLC analysis of cyclophosphamide in plasma (183) 49
Van den Eeckhout, E., Bens, G.A. and De Moerloose, P.
Separation and quantitation of bromperidol in pharmaceutical preparations by HPLC and HPTLC (193) 255
Van der Hoeven, T.A.
Separation of metabolites of testosterone hydroxylation by HPLC (196) 494
Van der Kaaden, A., see Ernst, G.F. (198) 526
Van der Kaay, M., see Tjaden, U.R. (181) 227
Van der Kerk-Van Hoof, A.C., see Huibregtse-Minderhout, L. (196) 425
Van der Kleijn, E., see Lankelma, J. (182) 35
Van der Kleyn, E., see O'Reilly, W.J. (183) 492
Van der Linde, H.C., see Hundt, H.K.L. (182) 110
Van der Meer, C. and Haas, R.E.
Determination of caffeine in serum by straight-phase HPLC (182) 121
Van der Meer, M.J. and Hundt, H.K.L.
Estimation of serum indomethacin at

therapeutic levels by means of TLC and spectrophotometry (181) 282
Van der Poll, J.M. and De Vos, R.H. Improved determination of dalapon residues in plant tissues and natural water by derivatization and electron capture GC (187) 244
Van der Poll, J.M. and Van de Lagemaat, D. Direct determination of propoxur in plant tissues by GC with an alkali flame-ionization detector (197) 263
Van der Sandt, J.S.F., see Barends, D.M. (182) 201
Van Dongen, W., see Nagels, L. (187) 181
Van Ginneken, C.A.M., see Hekman, P. (182) 252, (182) 492
Van Ginneken, C.A.M., see Van Loenhout, J.W.A. (182) 487
Van Hal, H.J.M., see Vink, J. (181) 25, (181) 115
Van Loenen, A.C., see Hulshoff, A. (181) 363
Van Loenhout, J.W.A., Ketelaars, H.C.J., Gribnau, F.W.J., Van Ginneken, C.A.M. and Tan, Y. Rapid HPLC method for the quantitative determination of diflunisal in plasma (182) 487
Van Niekerk, P.J. and Du Plessis, L.M. HPLC determination of *tert.*-butylhydroquinone in vegetable oils (187) 436
Van Roelenbosch, M., see Verzele, M. (186) 435
Van Rossum, J.M., see Fleuren, H.L.J. (182) 179
Van Vliet, H.P.M., Bootsman, T.C., Frei, R.W. and Brinkman, U.A.T. On-line trace enrichment in HPLC using a pre-column (185) 483
Vančura, V., see Wurst, M. (191) 129
VandenHeuvel, W.J.A., see Walker, R.W. (181) 85
Vanderborgh, N.E., see Jones, C.E.R. (186) 831
Vandereycken, G., see Van Boven, M. (182) 435
Vanderhaeghe, H. and Kerremans, L. TLC of macrolide antibiotics (193) 119
Vanderslice, J.T. and Maire, C.E. LC separation and quantification of B_6 vitamers at plasma concentration levels (196) 176
Vanhaelen, M. and Vanhaelen-Fastré, R. HPLC, GLC and TLC of naturally occurring flavonoids, phenolic and related compounds (187) 255
Vanhaelen-Fastré, R., see Vanhaelen, M. (187) 255
Varga-Puchony, Z., Vigh, G., Hlavay, J., Balogh, S. and Welsch, T. Separation of tetrazolium derivatives by reversed-phase HPLC (196) 185
Varga-Puchony, Z., see Vigh, G. (193) 432, (196) 1
Vasiliades, J. and Owens, C. Determination of midazolam in serum by GC with a nitrogen-selective detector (182) 439
Vasundhara, T.S. and Parihar, D.B. Studies of pyrazines as their n-π charge-transfer complexes with some nitro aromatic compounds (194) 254
Veazey, R.L. and Nieman, T.A. Chemiluminescence HPLC detector applied to ascorbic acid determination (200) 153
Vejrosta, J., see Novák, J. (199) 209
Venne, J.L.M. van de, see Van de Venne, J.L.M. (185) 241
Vercruysse, A., see Violon, C. (189) 94
Verebey, K., see Jukofsky, D. (198) 534
Vergin, H. and Bishop, G. HPLC determination of benzbromarone and the main metabolite benzarone in serum (183) 383
Verheggen, T., Mikkers, F., Everaerts, F., Oerlemans, F. and De Bruyn, C. Determination of uric acid in serum using isotachophoresis (182) 317
Verheggen, T., see Mikkers, F. (182) 496
Verhoef, J., see Tonnaer, J.A.D.M. (183) 303
Vermeer, J.M., see Guelen, P.J.M. (181) 497
Vernon, F., see Haken, J.K. (186) 89, (190) 291
Verpoorte, R., Mulder-Krieger, T., Troost, J.J. and Baerheim Svendsen, A. TLC separation of *Cinchona* alkaloids (Review) (184) 79
Verwey-van Wissen, C.P.W., see Fleuren, H.L.J. (182) 179
Verzele, M., Lammens, J. and Van Roelenbosch, M. Simplified HPLC phase characterization: particle size, surface area, mean pore diameter, amount of bonded phase and activity (186) 435
Verzele, M., Mussche, P. and Sandra, P. Characterization of HPLC stationary phases: the nature of silica gel bonded material (190) 331
Verzele, M., Redant, G., Qureshi, S. and Sandra, P. High-temperature quantitative glass capillary GC. Analysis of piperine and of quinine-quinidine (199) 105
Verzele, M., see Dewaele, C. (197) 189
Verzele, M., see Saeed, T. (186) 611
Vespalec, R., see Krejčí, M. (191) 3
Vessman, J. Quantitation of non-halogenated compounds by electron-capture GC (Review) (184) 313
Vessman, J., see Arbin, A. (196) 255
Vessman, J., see Gyllenhaal, O. (190) 347
Vessman, J., see Poole, C.F. (194) 365, (199) 123
Vida, L., see Szathmáry, C. (191) 327
Vielma, H. and Mendez, J. GC separation of 3-methylhistidine (196) 166
Vierkorn-Rudolph, B., Savelsberg, M. and Bächmann, K. Determination of trace amounts of hydrogen chloride by derivatization with epoxides and GC separation (186) 219
Vigh, G. and Varga-Puchony, Z. Influence of temperature on the retention

behaviour of members of homologous series in reversed-phase HPLC (196) 1
Vigh, G., Varga-Puchony, Z., Hlavay, J., Petró-Turcza, M. and Szárföldi-Szalma, I.
Separation of saturated and unsaturated aldehyde and ketone 2,4-dinitrophenylhydrazone derivatives by reversed-phase HPLC (193) 432
Vigh, G., see Varga-Puchony, Z. (196) 185
Vijayagopal, P., see Radhakrishnamurthy, B. (192) 307
Villanueva, V.R., see Adlakha, R.C. (187) 442
Vincent, J., see Engström, N. (189) 284
Vine, J.
Analysis of fatty acid methyl esters by high-resolution GC-chemical ionisation MS (196) 415
Vine, J., see Nichol, H. (182) 191
Vine, J.H., see Whitlam, J.B. (181) 463
Vink, J. and Van Hal, H.J.M.
Simplified method for determination of the tetracyclic antidepressant mianserin in human plasma using GC with nitrogen detection (181) 25
Vink, J., Van Hal, H.J.M. and Delver, B.
Comparative statistical study of assay methods using mass fragmentography and GC with nitrogen detection for determination of the tetracyclic antidepressant mianserin in human plasma (181) 115
Viola, J.L., see Williams, R.C. (185) 505
Violon, C. and Vercruysse, A.
Screening procedure for therapeutic benzodiazepines by HPLC of their benzophenones (189) 94
Vitenberg, A.G., see Ioffe, B.V. (186) 851
Vlasáková, V., see Holan, Z. (190) 67
Vliet, H.P.M. van, see Van Vliet, H.P.M. (185) 483
Vodička, L., Kříž, J., Průšová, D. and Burkhard, J.
HPLC of adamantanols and other cyclic alcohols (198) 457
Vodička, L., Průšová, D., Kříž, J. and Burkhard, J.
HPLC of adamantanones and other cyclic ketones (200) 238
Vodička, L., see Kuraš, M. (191) 319
Vodička, L., see Punčochářová, J. (191) 81
Voelter, W., Zech, K., Arnold, P. and Ludwig, G.
Determination of selected pyrimidines, purines and their metabolites in serum and urine by reversed-phase ion-pair chromatography (199) 345
Vogt, C.A., see Kapila, S. (196) 397
Vogt, C.R., see Kapila, S. (195) 17
Vogt, W., Jacob, K., Ohnesorge, A.-B. and Obwexer, H.W.
Capillary GC injection system for large sample volumes (186) 197
Vogt, W., Jacob, K., Ohnesorge, A.-B. and Schwertfeger, G.
Highly sensitive method for the quantitation of homovanillic acid in cerebrospinal fluid (199) 191
Vogt, W., see Meyer, H.D. (199) 339
Voituriez, L., see Cadet, J. (195) 139
Volkov, S.A., see Yeroshenkova, G.V. (198) 377
Volkov, S.M., Goryayev, V.M., Anikeyev, V.I. and Khripach, V.A.
Coating of glass capillary chromatographic columns: a new variant of the static method (190) 445
Von Arx, E., see Faupel, M. (193) 511
Vos, D. de, see De Vos, D. (183) 49, (183) 185
Vos, R.H. de, see De Vos, R.H. (187) 244
Voskamp, D., see Schaaper, W.M.M. (195) 181
Votavová, E., see Lukás, J. (194) 297
Votruba, J., see Holan, Z. (190) 67
Vozka, S., Porsch, B., Špaček, P. and Kubín, M.
Glass columns with high resistance to internal pressure for HPLC (193) 128
Vrátný, P., Ouhrabková, J. and Čopíková, J.
LC of non-reducing oligosaccharides: a new detection principle (191) 313
Vree, T.B., see Guelen, P.J.M. (181) 497
Vries, C.P. de, see De Vries, C.P. (183) 87
Vries, G. de, see De Vries, G. (192) 331, (198) 421
Vučković, G., see Ćelap, M.B. (196) 59

W

Wad, N., Weidkuhn, E. and Rosenmund, H.
A quantitative method for the simultaneous determination of carbamazepine, mephenytoin, phenylethylmalonamide, phenobarbital, phenytoin and primidone in serum by thin-layer chromatography. Additional comments to the analysis of carbamazepine-10,11-epoxide (183) 387
Wahlund, K.-G., see Sokolowski, A. (189) 299
Waibel, R., Amadò, R. and Neukom, H.
Purification of an α-L-arabinofuranosidase from *Aspergillus niger* by substrate affinity chromatography (197) 86
Wainwright, M.S. and Haken, J.K.
Evaluation of procedures for the estimation of dead time (Review) (184) 1
Wainwright, M.S., Haken, J.K. and Srisukh, D.
Linearity of the retention time *versus* carbon number plot for *n*-alkanes (188) 246
Wainwright, M.S., Haken, J.K. and Srisukh, D.
Linearity of retention data plots for *n*-alkanes on porous polymers in GC (198) 156
Wainwright, M.S., see Nieass, C.S. (194) 335
Wajcman, H., see Bachner, L. (193) 491
Waksmundzki, A., see Rayss, J. (188) 107
Walden, M.K., see Hunter, I.R. (198) 363
Waldron-Edward, D., see Moczar, E. (181) 108

Walker, E.A. and Castegnaro, M.
Response of nitramines in the thermal analyzer (187) 229
Walker, J.M., Hastings, J.R.B. and Johns, E.W.
Presence of sodium azide during acid hydrolysis of protein samples causes the destruction of tyrosine, phenylalanine and histidine (189) 106
Walker, R.W., Gruber, V.F., Pile, J., Yabumoto, K., Rosegay, A., Taub, D., Orme, M.L'E., Wolf, F.J. and VandenHeuvel, W.J.A.
Capillary column GLC-MS assay for 7α-hydroxy-5,11-diketotetranorprostane-1,16-dioic acid, the major human urinary metabolite of prostaglandins E_1 and E_2 (181) 85
Wall, R.A.
Hydrophobic chromatography with dynamically coated stationary phases. II. Dynamic cation-exchange separations of tyrosinyl peptides (194) 353
Wall, R.A., see Ghaemi, Y. (198) 397
Wallwork, M.A., see Gilbert, J. (193) 235
Walmsley, L.M., see Brodie, R.R. (182) 379
Walters, D.G., Lake, B.G. and Cottrell, R.C.
HPLC of coumarin and its metabolites (196) 501
Walters, R.R. and Buck, R.P.
Non-bonded affinity chromatography: a biospecific oil-soluble coated support for concanavalin A (188) 61
Wang Chinghai, see Lu Peichang (186) 25
Wang Guanghua, see Lu Peichang (186) 25
Wang, S.Y., see Cadet, J. (195) 139
Wännman, T., see Blomberg, L. (186) 159
Ward, D.G., see Middleditch, B.S. (195) 359
Warsh, J.J., Chiu, A., Li, P.P. and Godse, D.D.
Comparison of LC-electrochemical and GC-MS methods for brain dopamine and serotonin (183) 483
Warwick, P., see Malcolme-Lawes, D.J. (200) 47
Washitake, M., see Seta, K. (181) 311
Wasiak, W., see Nawrocki, J. (188) 323
Watabe, K., Suzuki, S. and Araki, S.
Liquid crystals as the stationary phase in GC. Adsorption phenomena caused by an electric field applied across the column (192) 89
Watanabe, A., see Tanaka, A. (194) 21
Watanabe, Y., see Takeuchi, T. (192) 127
Watson, T.R., see Allan, R.J. (183) 311
Watts, C.D., see Crathorne, B. (185) 671
Watzlawick, H., see Schott, H. (196) 435
Wawrzynowicz, T., see Siouffi, A.M. (186) 563
Weber, M., Foglietti, M.J. and Percheron, F.
Affinity chromatography of a cellulase complex from *Trichoderma viride* on cross-linked cellulose (in French) (188) 377
Weekley, A., see Dilli, S. (195) 349
Weerdt, G.A. de, see De Weerdt, G.A. (193) 504
Wehr, C.T. and Abbott, S.R.
High-speed steric exclusion chromatography of biopolymers (185) 453
Wei-Tung Wang, Xian-Dong Ding and Xin-Jie Wu
Determination of water at the parts per million level in buta-1,3-diene and associated solvents by GC (199) 149
Weidkuhn, E., see Wad, N. (183) 387
Weidner, M.M., see Jardine, I. (182) 395
Wejman, W., see Kaniewska, T. (182) 81
Welling, P.L.M., see Scholten, A.H.M.T. (199) 239
Welsch, T., see Varga-Puchony, Z. (196) 185
Wenk, G. and Greenland, R.
Investigation of the performance of a HPLC system with an electrochemical detector (183) 261
Wentworth, W.E. and Chen, E.C.M.
Analysis of kinetics and concentration dependence of electron-capture detector (186) 99
Wentworth, W.E., see Ayala, J.A. (195) 1
Wentworth, W.E., see Chen, E.C.M. (188) 89
Werken, P.H. van de, see Van de Werken, P.H. (198) 449
Werquin, J.P., see Szymanowicz, A.G. (190) 457
West, C.E., see Shaw, D.C. (200) 185
Westberg, H., see Osman, M. (186) 273
Westenberg, H.G.M., see Greving, J.E. (186) 683
Westerdahl, G., see Larsson, L. (182) 402
Westerlund, D. and Erixson, E.
Reversed-phase chromatography of zimelidine and similar dibasic amines. I. Analysis in biological material (185) 593
Westerlund, D., see Henke, B. (187) 189
Westmore, J.B., see Quilliam, M.A. (194) 379, (196) 367
Wetzel, D., see Mauro, D. (187) 421
Wheals, B.B.
Isocratic multi-column HPLC as a technique for qualitative analysis and its application to the characterisation of basic drugs using an aqueous methanol solvent (187) 65
Wheeler, L.A., De Meo, M., Kirby, B.D., Jerauld, R.S. and Finegold, S.M.
HPLC assay for measurement of cefoxitin in serum (183) 357
White, J.D. and Schwartz, D.P.
Evaluation of Amberlite XAD-4 for recovery of glucuronides and sulfates from aqueous solution (196) 303
White, J.G., see Rao, G.H.R. (196) 506
White, P.C.
Use of dual-wavelength UV detection in HPLC for the identification of barbiturates (200) 271
Whitehead, T.P., see Clark, P.M.S. (181) 347
Whitlam, J.B. and Vine, J.H.
Quantitation of ibuprofen in biological fluids by GC-MS (181) 463
Wibell, L., see Ekman, L. (182) 478
Wibell, L., see Lanbeck, K. (182) 262
Wickramanayake, P.P. and Aue, W.A.
Empirical correlation of layer thickness and support pore size in "chemically bonded" phases (195) 25

Wickramanayake, P.P., see Aue, W.A. (197) 21, (200) 3
Wickstrom, E., see Tyson, R.W. (192) 485
Wickström, G., see Hartvig, P. (183) 229
Widén, C.-J., Pyysalo, H. and Salovaara, P.
Separation of naturally occurring acylphloroglucinols by HPLC (188) 213
Widén, K.-G., see Pyysalo, H. (190) 466
Wiegant, V.M., see Tonnaer, J.A.D.M. (183) 303
Wightman, F., see Markham, G. (192) 429
Wightman, R.M., see Hirata, Y. (181) 287
Wijkens, P., see Huibregtse-Minderhout, L. (196) 425
Wilensky, A.J., see Raisys, V.A. (183) 441
Wilhelm, J.C. and Bloch, J.M.
Separation of long-chain fatty acids extracted from peat by gel permeation chromatography (in French) (193) 329
Willems, G.H.W.
New results with a flow-impedance bridge detector (189) 3
Williams, H., see Stocklinski, A.W. (195) 281
Williams, K., see Ballantine, J.A. (198) 111
Williams, K.J., Li Wan Po, A. and Irwin, W.J.
Sample-solvent-induced peak broadening in the reversed-phase HPLC of Aspirin and related analgesics (194) 217
Williams, P.A., see Stewart, J.G. (198) 489
Williams, P.L., Moffat, A.C. and King, L.J.
Combined HPLC and radioimmunoassay method for the analysis of Δ^9-tetrahydrocannabinol metabolites in human urine (186) 595
Williams, R.C. and Viola, J.L.
Application of an automated extractor/concentrator to the analysis of anticonvulsant drugs in blood serum by HPLC (185) 505
Williams, R.L., see Upton, R.A. (190) 119
Willis, J.V., see Schweizer, A. (195) 421
Wilson, I.D., Bielby, C.R., Morgan, E.D. and McLean, A.E.M.
Comparison of HPLC and GC for the analysis of ecdysteroids (194) 343
Wilson, I.D., see Bielby, C.R. (194) 43
Wilson, J.B., see Henson, J.B. (198) 443
Wilson, N.Y., see Di Ferrante, D.T. (187) 271
Wininger, P.E., see Tindall, G.W. (196) 109
Wiśniewski, A., see Szafranek, J. (187) 131
Wissemann, K.W. and Lee, C.Y.
Purification of grape polyphenoloxidase with hydrophobic chromatography (192) 232
Witkiewicz, Z., Szulc, J., Dąbrowski, R. and Sadowski, J.
Properties of liquid crystalline cyanoazoxybenzene alkyl carbonates as stationary phases in GC (200) 65
Witkiewicz, Z., see Rayss, J. (188) 107
Wittfoht, W., see Nau, H. (182) 71
Witzgall, H. and Hassan-Ali, S.
PC systems for the separation of aldosterone, 18-hydroxycorticosterone, 18-hydroxydeoxycorticosterone, corticosterone and deoxycorticosterone (198) 70
Wizner, M.A., see Sievers, R.E. (186) 3
Wojaczyńska, M. and Kolarz, B.N.
Structure and sorption properties of porous copolymers of styrene, acrylonitrile and divinylbenzene (196) 75
Wolf, F.J., see Walker, R.W. (181) 85
Wolff, J.L., see Hoener, B.-A. (182) 246
Wolfram, K.M. and Bjornsson, T.D.
HPLC analysis of dipyridamole in plasma and whole blood (183) 57
Wolfram, K.M., see Bjornsson, T.D. (181) 417
Wöll, J., see Liebich, H.M. (199) 181
Wootton, M., see Tangendjaja, B. (197) 274
Wray, N.P., see Marshall, M.V. (197) 217
Wright, B.W., Lee, M.L., Graham, S.W., Phillips, L.V. and Hercules, D.M.
Glass surface analytical studies in the preparation of open tubular columns for GC (199) 355
Wright, B.W., see Lee, M.L. (184) 235
Wrolstad, R.E., see Akhavan, I. (190) 452
Wroński, M.
Counter-current distribution of hydrophilic thiols in the presence of tributyltin hydroxide (190) 156
Wu, H.-L., Hsu, L.-C. and Hsü, C.-Y.
GC determination of piromidic acid (193) 476
Wurst, F., Prey, T., Puchinger, L. and Bancher, E.
New method for the quantitative analysis of galanthamine in drug extracts from *Leucojum aestivum* (in German) (188) 452
Wurst, M., Přikryl, Z. and Vančura, V.
HPLC of plant hormones. I. Separation of plant hormones of the indole type (191) 129
Wurst, M., see Patzelová, V. (193) 137

X

Xia Aizu, see Lu Peichang (186) 25
Xian-Dong Ding, see Wei-Tung Wang (199) 149
Xin-Jie Wu, see Wei-Tung Wang (199) 149
Xu Fangbao, see Lu Peichang (186) 25

Y

Yabumoto, K., see Walker, R.W. (181) 85
Yago, L.S., see Pitts, F.N. Jr. (193) 157
Yamagami, K., see Inayama, S. (194) 85
Yamaguchi, K., see Shibamoto, T. (194) 277
Yamaguchi, S., see Nakano, N.I. (188) 347

Yamaguchi, Y., Hayashi, C. and Miyai, K.
Enzymatic color development of 3α-hydroxysteroids on thin-layer chromatograms for determination of excretion pattern of 3α-hydroxysteroids in patients with some adrenogenital syndrome (182) 430
Yamamoto, I., see Kawano, Y. (198) 317
Yamamoto, M., see Kodama, H. (183) 226
Yamamoto, S., Kakuno, K., Okahara, S., Kataoka, H. and Makita, M.
GC of phenolic amines, 3-methoxycatecholamines, indoleamines and related amines as their N,O-ethyloxycarbonyl derivatives (194) 399
Yamamoto, S., see Makita, M. (188) 408
Yamamoto, Y., see Nakamura, H. (200) 324
Yamauchi, Y., Oshima, R. and Kumanotani, J.
Separation of Japanese lac urushiol diacetate on silver nitrate-coated silica gel columns by HPLC (198) 49
Yanagihara, K., see Kawano, Y. (198) 317
Yang, F.C., see Budd, R.D. (190) 129
Yang, H.-Y.T., see Peralta, E. (190) 43
Yang, S.K., see Chou, M.W. (185) 635
Yano, Y. and Janado, M.
Hydrophobic interaction chromatography of aliphatic alcohols on unsubstituted Sephadex gels with high dextran concentrations (200) 125
Yano, Y., see Janado, M. (193) 345
Yashin, Y.I., see Kiselev, A.V. (195) 205
Yasueda, H., see Mita, H. (181) 153
Yeadon, G., see Dawkins, J.V. (188) 333
Yerchak, N.P., see Shatz, V.D. (200) 105
Yeroshenkova, G.V., Volkov, S.A. and Sakodynskii, K.I.
Influence of sorption on the shape of chromatographic elution curves (198) 377
Yeung, E.S., see Sepaniak, M.J. (190) 377
Yoda, R., see Nakamura, H. (200) 324
Yokoyama, T., see Furukawa, M. (198) 212
Yoshimura, T., see Ujimoto, K. (190) 161
Young, J.G., see Anderson, G.M. (181) 453
Young, R.W., see Lim, K.L. (188) 285
Yoza, N., Ito, K., Hirai, Y. and Ohashi, S.
Optimization of HPLC of inorganic polyphosphates for routine analyses (196) 471
Yui, T., see Matsuki, Y. (188) 177

Z

Zacest, R., see Reece, P.A. (181) 427
Zaerr, J.B., see Crozier, A. (198) 57
Zahn, R.K., see Breter, H.-J. (193) 175
Zakaria, M., see Krstulovic, A.M. (186) 733
Zanini, M.G., see Caccia, S. (190) 475
Zapata, R., see Hanna, S. (200) 277
Zappia, V., Galletti, P., Porcelli, M., Manna, C. and Della Ragione, F.
HPLC separation of natural adenosyl-sulphur compounds (189) 399
Zarrella, T., see Morrison, H. (183) 83
Zázvorková, M., see Šmejkal, F. (197) 147
Zech, K., see Voelter, W. (199) 345
Zeeuw, R.A. de, see De Zeeuw, R.A. (186) 683, (191) 109
Zegota, H.
Separation and quantitative determination of fructose as the O-methyloxime by GLC using glass capillary columns (192) 446
Zelchan, G.I., see Shatz, V.D. (200) 105
Zelenský, I., see Kaniansky, D. (194) 11
Zelt, D.T., Owen, J.A. and Marks, G.S.
Second derivative-HPLC-fluorometric detection of porphyrins in chick embryo liver cell culture medium (189) 209
Zemek, J., see Mislovičová, D. (194) 95
Zhang Yukuei, Bao Miansheng, Li Xiouzhen and Lu Peichang
HPLC columns of small diameter (197) 97
Zhou Liangmo, see Lu Peichang (186) 25
Zia, H., Proveaux, W.J., O'Donnell, J.P. and Ma, J.K.H.
Chromatographic analysis of griseofulvin and metabolites in biological fluids (181) 77
Ziegler, J.M., see Sarhan, F. (183) 505
Zielinski, W.L. Jr. and Janini, G.M.
Utility of high-temperature thermotropic liquid crystals as stationary phases for novel GLC separations (186) 237
Zielinski, W.L. Jr., see Janini, G.M. (193) 444
Zima, J., see Smolková, E. (191) 61
Zimmerman, L.A., see Tolan, J.W. (190) 367
Zio, C.A., Massarani, G. and Pietta, P.
Estimation of pipemidic acid and N-nitrosopipemidic acid by HPLC (200) 245
Zlatkis, A., Poole, C.F., Brazeli, R., Bafus, D.A. and Spencer, P.S.
Volatile metabolites in sera of normal and diabetic patients (182) 137
Zlatkis, A., see Poole, C.F. (184) 99, (186) 307, (187) 331, (199) 123
Zlatkis, A., see Singhawangcha, S. (183) 433
Zöllner, E.J., see Breter, H.-J. (193) 175
Zuccato, E., see Marcucci, F. (198) 180
Zuidema, J. and Hilbers, H.
GLC determination of acephylline in urine (182) 445
Zuidema, J., Modderman, E.S.M., Hilbers, H.W., Merkus, F.W.H.M. and Huikeshoven, H.
Rapid HPLC method for the determination of dapsone and monoacetyldapsone in biological fluids (182) 130
Zumwalt, R.W., see Gehrke, C.W. (188) 129

SUBJECT INDEX

VOLUMES 181—200

The volume number is placed in parentheses before the page number

A

Abscisic acid
High-performance steric exclusion chromatography of plant hormones (198) 57

Acebutolol metabolites
Semi-automated HPLC method for the determination of diacetolol hydrochloride in medicated animal diets (200) 261

Acephylline
GLC determination of acephylline in urine (182) 445

Acetaldehyde
Simple rapid and sensitive method for the simultaneous quantitation of ethanol and acetaldehyde in biological materials using head-space GC (190) 197

Acetaldehyde
GLC analysis of dimedone derivatives of formaldehyde and other aliphatic aldehydes on capillary columns (191) 239

Acetaldehyde
Determination of formaldehyde, acetaldehyde and associated components in solution and in vapours by GC (200) 206

Acetamide
TLC method for the separation and determination of the products of the reaction of amides with formaldehyde (195) 158

Acetates
Separation of a series of positional and geometrical isomers of olefinic aliphatic primary alcohols and acetates by capillary GC (189) 199

Acetophenones
Separation and identification of some hydroxyacetophenones and their derivatives by TLC (188) 289

Acetylhydrazinophthalazinone
Determination of hydralazine metabolites: 4-hydrazino-phthalazin-1-one and N-acetylhydrazinophthalazin-1-one by GC and *s*-triazolo[3,4-*a*]phthalazine and phthalazinone by HPLC (183) 167

Acetylprocainamide
Comparison of HPLC with an enzyme multiplied immunoassay technique for the analysis of serum procainamide and N-acetylprocainamide (181) 490

Acetylsalicylic acid
Sample-solvent-induced peak broadening in the reversed-phase HPLC of Aspirin and related analgesics (194) 217

Acids
Relative effectiveness of ion-exchange and lead acetate precipitation methods in isolating pear sugars and acids (190) 452

Acrylonitrile
Structure and sorption properties of porous copolymers of styrene, acrylonitrile and divinylbenzene (196) 75

Acrylonitrile-styrene copolymer
Determination of the composition of copolymers as a function of molecular weight by pyrolysis GC size-exclusion chromatography (194) 163

Acylglycines
GC retention indices of twenty metabolically important acylglycines as trimethylsilyl derivatives (181) 90

Acylon
GC determination of Metalaxyl (Ridomil) residues in tobacco (200) 313

Acylon
Estimation of the systemic fungicide Ridomil by TLC (200) 317

Acylphloroglucinols
Separation of naturally occurring acylphloroglucinols by HPLC (188) 213

Adamantanols
HPLC of adamantanols and other cyclic alcohols (198) 457

Adamantanones
HPLC of adamantanones and other cyclic ketones (200) 238

Adamexina
Determination of a new mucolytic drug Adamexina in biological liquids by photodensitometry (181) 504

Adenine nucleosides
Identification of adenosine and eight modified adenine nucleosides using reversed-phase HPLC and enzymatic peak shift with adenosine deaminase (183) 499

Adenine nucleotides
Simple anion-exchange chromatography for the determination of adenine nucleotides by using AG MP-1 resin (192) 193

Adenine nucleotides
Reversed-phase HPLC of adenine nucleotides: application to the kinetics of an adenosine 3'-phosphate 5'-sulphatophosphate sulphotransferase from plants (193) 285

Adenine nucleotides
Operational parameters of anion-exchange chromatography using AG MP-1 resin for rapid assay of adenine nucleotides (198) 500

Adenosine
Identification of adenosine and eight modified adenine nucleosides using reversed-phase HPLC and enzymatic peak shift with adenosine deaminase (183) 499

Adenosine monophosphate
Study of the interaction between L-lactate dehydrogenase isoenzymes and immobilized 8-substituted adenosine 5'-monophosphate by means of affinity electrophoresis (192) 351

Adenosine monophosphate
Study of the interaction between NADP-dependent dehydrogenase and immobilized adenosine 2'-monophosphate by means of affinity electrophoresis (196) 85

Adenosine monophosphate
Control of 8-bromo-AMP and 8-(6-aminohexyl)amino-AMP synthesis by HPLC (196) 323

Adenosine nucleotides
Simultaneous analysis of ATP, ADP, AMP, and other purines in human erythrocytes by HPLC (181) 103

Adenosyl-sulphur compounds
HPLC separation of natural adenosyl-sulphur compounds (189) 399

Adenylic acid
Application of a weakly basic dimethylamino-modified silica ion exchanger to the separation of oligonucleotides (185) 403

Adiphenine hydrochloride
Determination of adiphenine hydrochloride and diphenylacetic acid by ion-pair HPLC (200) 267

Adrenaline
Simple method for isolating free urinary catecholamines by boric acid gel chromatography (183) 78

Adrenaline
Semi-automated fluorimetric method for the estimation of urinary catecholamines using HPLC (183) 295

Adrenaline
Electrochemical detector for HPLC (192) 135

Adrenocortical steroids
Quantitative HPLC determination of Δ^4-3-ketosteroids in adrenocortical extracts (193) 413

Adrenocorticosteroidal agents
Analysis of dexamethasone sodium phosphate formulations by HPLC (193) 160

Adrenocorticotropin/lipotropin precursor
Reversed-phase HPLC purification and characterization of the adrenocorticotropin/lipotropin precursor and its fragments (193) 291

Adriamycin
Determination of adriamycin and daunorubicin in urine by HPLC with laser fluorometric detection (190) 377

Adriamycin
Determination of adriamycin in liposomes by HPLC using a fluorescence detector (190) 489

Adriamycin
Determination of the concentration of adriamycin and its metabolites in the serum and tissues of Ehrlich carcinoma-bearing mice by HPLC (196) 463

Adriamycin
HPLC and TLC of anthracycline antibiotics. Separation and identification of components of the daunorubicin complex from fermentation broth (198) 407

Adsorbents
Coordination polymers as adsorbents and stationary phases in GC. VI. Separation properties of alkaline-earth metal di-*n*-hexylphosphinates (188) 323

Adsorbents
LSC on Amberlite XAD-2 and other styrene-divinylbenzene adsorbents. I. Development of a solvent eluotropic scale (189) 145

Adsorbents
Crosslinked polymethylphenylsiloxanes as adsorbents for GC (189) 169

Adsorbents
A PTFE-based carbon adsorbent in HPLC (191) 61

Adsorbents
Stabilized Y-type zeolites for GC separation (191) 175

Adsorbents
Chromatography of hydrocarbons on molybdenum sulphide (192) 323

Adsorbents
JADO carbon as an adsorbent in sampling columns (193) 137

Adsorbents
Resolution of neutral complexes of transition metals by stereoselective adsorption on optically active complexes. I. Partial resolution of tris(acetylacetonato)cobalt(III) and tris(acetylacetonato)chromium(III) on cobalt(III) complexes (198) 172

Adsorption
Mechanism of solute retention in liquid-solid chromatography and the role of the mobile phase in affecting separation. Competition *versus* "sorption" (Review) (184) 363

Adsorption
GC technique for temperature-programmed desorption of reversibly and irreversibly adsorbed species (192) 420

Adsorption-desorption rates
GC determination of adsorption-desorption rates. Interaction of hydrogen on a nickel catalyst (196) 355

Adsorption effects
Effect of adsorption on solute retention in GLC (186) 47

Adsorption effects
Study of the adsorption effects at the surface of poly(ethylene glycol)-coated column packings (186) 63

Adsorption effects
Adsorption effects in gas-liquid partition chromatography. Quantitative evaluation (190) 21

Adsorption enthalpies and entropies
Use of isothermal GC for the determination of the adsorption enthalpies and entropies of inorganic halides at high temperatures (187) 319

Adsorption isotherms
GC adsorption studies on synthetic imogolite (188) 97

Adsorption phenomena
Liquid crystals as the stationary phase in GC. Adsorption phenomena caused by an electric field applied across the column (192) 89

Aerosols
Correction for artifacts in the analysis of atmospheric aerosols (190) 136

Affinity ligands
Affinity chromatography of cells and cell membranes (Review) (184) 471

Agarose
Model studies of the reactions and non-specific adsorptive properties of cyanogen bromide-activated macroporous agarose (196) 379

Agarose beads
Quantitative determination of alkylamino side-chains coupled to agarose beads. Comparison of methods (193) 451

Agarose isoelectric focusing
New developments in isoelectric focusing (Review) (184) 415

Aglycones
Chromogenic reagent for vicine and convicine on thin-layer plates (194) 251

Air analysis
Vapor-phase silylation of alcohols for air analysis (186) 273

Air analysis
Evaluation of chromatographic sorbents used in air pollution studies (186) 811

Alcohols
Vapor-phase silylation of alcohols for air analysis (186) 273

Alcohols
Separation of a series of positional and geometrical isomers of olefinic aliphatic primary alcohols and acetates by capillary GC (189) 199

Alcohols
HPLC of adamantanols and other cyclic alcohols (198) 457

Alcohols
Study of the interactions between substituted phenols and alcohols and dibenzyl sulphoxide by means of GC (198) 495

Alcohols
TLC separation of some ferrocene alcohols (198) 536

Alcohols
Hydrophobic interaction chromatography of aliphatic alcohols on unsubstituted Sephadex gels with high dextran concentrations (200) 125

Aldehydes
GC determination of low-molecular-weight carbonyl compounds in aqueous solution as their O-(2,3,4,5,6-pentafluorobenzyl) oximes (187) 413

Aldehydes
GLC analysis of dimedone derivatives of formaldehyde and other aliphatic aldehydes on capillary columns (191) 239

Aldehydes
Separation of saturated and unsaturated aldehyde and ketone 2,4-dinitrophenylhydrazone derivatives by reversed-phase HPLC (193) 432

Aldehydes
GC analysis of aliphatic and aromatic aldehydes as trimethylsilylated dithioacetals of 2-mercaptoethanol (194) 191

Aldosterone
PC systems for the separation of aldosterone, 18-hydroxycorticosterone, 18-hydroxydeoxycorticosterone, corticosterone and deoxycorticosterone (198) 70

Aldrin
Retention behavior of chlorinated benzenes, chlorinated phenols, Aldrin, and Aroclor 1242 on Tenax-GC, Chromosorb 101, Florisil, and Carbopack C HT (200) 115

Alkadienes
Identification of *n*-hepta- and *n*-octadienes by high-resolution GC using structure-retention correlations and MS (191) 187

Alkaline-earth metal dihexylphosphinates
Coordination polymers as adsorbents and stationary phases in GC. VI. Separation properties of alkaline-earth metal di-*n*-hexylphosphinates (188) 323

Alkaloids
Analysis of salsolinol and salsoline in biological samples using deuterium-labelled internal standards and GC-MS (183) 17

Alkaloids
TLC separation of *Cinchona* alkaloids (Review) (184) 79

Alkaloids
Rapid quantitative analysis of solasodine, solasodine glycosides and solasodiene by HPLC (187) 87

Alkaloids
Chromogenic reactions of *Rauwolfia* alkaloids after separation by TLC (187) 199

Alkaloids
Amperometric HPLC method for narcotic alkaloids (188) 420

Alkaloids
New method for the quantitative analysis of galanthamine in drug extracts from *Leucojum aestivum* (in German) (188) 452

Alkaloids
Cactus alkaloids. XL. Identification of mescaline and other β-phenethylamines in *Pereskia, Pereskiopsis* and *Islaya* by use of fluorescamine conjugates (189) 79

Alkaloids
Reversed-phase HPLC analysis of cinchona alkaloids in pharmaceuticals (189) 241

Alkaloids
Optical brighteners as TLC detection reagents for glycoalkaloids and steroid alkaloids in *Solanum* species. II. Blankophor BA 267%, BBU neu and KU, and Tinopal CBS-X and 5 BMS-X (189) 406

Alkaloids
Separation and determination of opium alkaloids by HPLC (190) 251

Alkaloids
Separation of ergotoxine alkaloids by HPLC on silica (191) 101

Alkaloids
Investigation of vinca alkaloid-plasma membrane interactions by detergent gel chromatography (191) 287

Alkaloids
Determination of perloline in grass species (in German) (192) 199

Alkaloids
Rapid HPLC isolation of monoesters, diesters and macrocyclic diester pyrrolizidine alkaloids from *Senecio jacobaea* and *Amsinckia intermedia* (192) 474

Alkaloids
Analysis of cinchona alkaloids by HPLC (in French) (192) 479

Alkaloids
Detection and assay of dihydroergot alkaloids by TLC using *o*-phthalaldehyde-sulphuric acid reagent (193) 500

Alkaloids
Fluorimetric and HPLC determination of harmane alkaloids in *Peganum harmala* cell cultures (194) 234

Alkaloids
Preparative separation of the pyrrolizidine alkaloids, intermedine and lycopsamine, as their borate complexes (195) 379

Alkaloids
Chloranil, a sensitive detection reagent for pyrrolizidine alkaloids on thin-layer chromatograms (195) 407

Alkaloids
Specific detection of pyrrolizidine alkaloids on thin-layer chromatograms (195) 412

Alkaloids
HPLC of *Solanum* and *Veratrum* alkaloids (198) 363

Alkanes
Linearity of the retention time *versus* carbon number plot for *n*-alkanes (183) 246

Alkanes
Dependence of the relative molar response of isomeric alkanes on molecular structure (in German) (194) 213

Alkanes
Adsorption chromatography capillary columns. I. Rubidium chloride (196) 237

Alkanes
Error in the determination of the specific retention volume. Specific retention volumes

of *n*-alkanes on Apolane-87 stationary phase as functions of column temperature (198) 85

Alkanes
Linearity of retention data plots for *n*-alkanes on porous polymers in GC (198) 156

Alkenes
Use of Carbowax 20M glass capillary columns below their lower temperature limit (190) 283

Alkenes
Topological analysis of the behaviour of linear alkenes up to tetradecenes in GLC on squalane (194) 121

Alkenes
GC of all C_{15}-C_{18} linear alkenes on capillary columns with very high resolution power (195) 43

Alkenyl catechols
HPLC separation of urushiol congeners in poison ivy and poison oak (200) 163

Alkenylnaphthalenes
GC of some alkenyl- and cycloalkenylnaphthalenes on Bentone 34, silicone fluid DC 550, and Apiezon L phases (198) 1

Alkylamino side-chains
Quantitative determination of alkylamino side-chains coupled to agarose beads. Comparison of methods (193) 451

Alkylammonium ions
Ion-pair chromatography in the low concentration range by use of highly absorbing counter ions. III. HPLC of quaternary alkylammonium ions as ion pairs with naphthalene-2-sulphonate, using a silica support of low surface area (193) 225

Alkylbenzenes
Preparative separation of C_9 and C_{10} monoaromatic hydrocarbon mixtures by HPLC on silica gel (191) 319

Alkylbenzyldimethylammonium compounds
GLC of straight chain homologues of alkylbenzyldimethylammonium compounds (200) 216

Alkyl chain length
Effect of alkyl chain length of the stationary phase on retention and selectivity in reversed-phase LC. Participation of solvent molecules in the stationary phase (199) 327

Alkyl chains
GC study of grafted alkyl chains used as stationary phases (196) 481

Alkyl group redistribution products
GC with plasma emission spectroscopic detection of Friedel-Crafts catalyzed alkyl group redistribution products among Si, Ge, Sn and Pb atoms (196) 265

Alkyl monochlorides
Non-linear responses of the electron-capture detector to alkyl monochlorides (192) 117

Alkyltin halides
Comparative assessment of GLC and HPLC for the separation of tin tetraalkyls and alkyltin halides (200) 305

Allethrin
Examination of the conversion products of pyrethrins and allethrin formulations exposed to sunlight by GC and MS (198) 317

Allopurinol
Improvements to a HPLC assay for allopurinol and oxipurinol in plasma (181) 286

Alprenolol
Formation of electron-capturing derivatives of alprenolol by transboronation. Application to the determination of alprenolol in plasma (194) 365

Alternariol
HPLC determination of major mycotoxins produced by *Alternaria* molds (194) 89

Amantadine
GC determination of amantadine hydrochloride (Symmetrel) in human plasma and urine (183) 33

Amarogentin
Determination of the bitter constituents of the Gentiana root by HPLC (193) 163

Amaropanin
Determination of the bitter constituents of the Gentiana root by HPLC (193) 163

Amaroswerin
Determination of the bitter constituents of the Gentiana root by HPLC (193) 163

Amberlite XAD-4
Evaluation of Amberlite XAD-4 for recovery of glucuronides and sulfates from aqueous solution (196) 303

Amide-formaldehyde addition products
TLC method for the separation and determination of the products of the reaction of amides with formaldehyde (195) 158

Amine phosphate buffers, acidic
Mobile phase effects in reversed-phase chromatography. II. Acidic amine phosphate buffers as eluents (185) 111

Amines
Retension mechanism in reversed-phase ion-pair chromatography of amines and amino acids on bonded phases (185) 241

Amines
Improved GC method for the quantitative determination of secondary amines as sulphonamides formed by reaction with benzenesulphonyl chloride (190) 462

Amines
GC and MS properties of ethyl esters of dithiocarbamic acids of primary and secondary amines (193) 61

Amines
Adsorption-derivatization GC-MS determination of aqueous 1,6-hexanediamine at the low and sub-ppm level (193) 271

Amines
GLC determination of toxic diamines in permanent hair dyes (193) 277

Amines
GC of phenolic amines, 3-methoxycatechol-amines, indoleamines and related amines as their N,O-ethyloxycarbonyl derivatives (194) 399

Amines
Quantitative structure-retention relationships in cation-exchange chromatography (198) 241

Amines
Fluorescence detection of secondary amines on thin-layer plates using a fluorogenic reaction with fluorescamine (200) 324

Amines, aliphatic
Chromatographic behaviour of aliphatic amines on phenol-impregnated thin layers (196) 225

Amino acid enantiomers
Synthesis and properties of a novel chiral stationary phase for the resolution of amino acid enantiomers (186) 611

Amino acids
Method for the determination of tryptophan in serum and cerebrospinal fluid (181) 100

Amino acids
Determination of tryptophan and metabolites in rat brain and pineal tissue by reversed-phase HPLC with electrochemical detection (181) 187

Amino acids
Rapid automated ion-exchange analysis of plasma tyrosine and phenylalanine with data print-out (181) 337

Amino acids
HPLC determination of 5-hydroxyindole-3-acetic acid in urine using Sephadex G-10 for isolation (181) 456

Amino acids
TLC method for the quantitative analysis of L-tryptophan in human plasma (182) 110

Amino acids
GLC of free amino acids in the hyaloplasm of rat cerebral, cerebellar and ocular tissues, and in skeletal and heart muscle (182) 147

Amino acids
Employment of GLC for the analysis of collagen amino acids in biopsy tissue (182) 155

Amino acids
Automated ion-exchange chromatography in the detection of aspartylglucosaminuria (182) 409

Amino acids
Rapid HPLC method for quantitation of 3-methylhistidine (182) 414

Amino acids
Separation of peptides and amino acids by ion-exchange chromatography of their copper complexes (183) 9

Amino acids
Isotachophoretic analysis of some sulfur-containing amino acids in human urine (183) 226

Amino acids
Determination of leucine flux in vivo by GC-MS utilizing stable isotopes for trace and internal standard (183) 403

Amino acids
Retension mechanism in reversed-phase ion-pair chromatography of amines and amino acids on bonded phases (185) 241

Amino acids
Reversed-phase separation of optical isomers of Dns-amino acids and peptides using chiral metal chelate additives (185) 323

Amino acids
Ligand-exchange chromatography of enantiomeric amino acids on copper-loaded chiral bonded silica gel and of amino acids on copper(II)-modified silica gel (185) 345

Amino acids
HPLC of amino acids, peptides and proteins. XX. Investigation of the effect of pH and ion-pair formation on the retention of peptides on chemically-bonded hydrocarbonaceous stationary phases (185) 429

Amino acids
LC resolution of enantiomers on normal-phase chiral amide-bonded silica gel. Retentions of optically active α-amino acid derivatives on N-acyl homologues of L-valylaminopropylsilanized silica phases (186) 543

Amino acids
Analysis of cysteic acid and cysteine sulfinic acid by GLC (187) 239

Amino acids
Chromatographic method for the measurement of hydroxylysine, hydroxylysine glycosides and 3-methylhistidine in human urine (187) 271

Amino acids
Gas chromatography of trifluoroacetyl amino acid trimethylsilyl esters. II. Problems in preparing specified derivatives of hydroxyamino acids (in German) (188) 251

Amino acids
Resolution of histidine diastereomers by GC (188) 408

Amino acids
Separation of iodinated compounds of L-tyrosyl-L-tyrosine from iodothyronines by Biogel P-2 CC (188) 431

Amino acids
Separation of S-sulphocysteine and related compounds by anion-exchange chromatography and electrophoresis (188) 442

Amino acids
Presence of sodium azide during acid hydrolysis of protein samples causes the destruction of tyrosine, phenylalanine and histidine (189) 106

Amino acids
TLC of S-methylcysteine sulphoxide (kale anaemia factor) in the presence of common amino acids (189) 436

Amino acids
Improved method for the preparation of 3-hydroxyproline, and some of its chromatographic properties (190) 457

Amino acids
HPLC of amino acids, peptides and proteins. XXI. The application of preparative reversed-phase HPLC for the purification of a synthetic underivatised peptide (192) 222

Amino acids
Ligand-exchange chromatography. I. Resolution of L- and D-proline on a copper(II)-proline complex bound to microparticulate silica gel (192) 228

Amino acids
Separation of alkylaminonaphthylenesulfonyl peptides and amino acids by HPLC. Methods for measuring melanotropin inhibiting factor breakdown (192) 341

Amino acids
New horizontal flow-through coil planet centrifuge for counter-current chromatography. III. Separation and purification of dinitrophenyl amino acids and peptides (193) 37

Amino acids
Simple and rapid mass fragmentographic method for the determination of glycine in brain tissue (193) 95

Amino acids
Separation of hypoglycin A from leucine and other amino acids on Sephadex G-10 (193) 148

Amino acids
Two-dimensional TLC of Dns-amino acids on reversed-phase silica gel (193) 421

Amino acids
Program for processing amino acid data with a programmable pocket calculator (194) 69

Amino acids
Effect of some organic buffers on the estimation of aspartic acid and resolution in amino acid analysis (194) 76

Amino acids
Reversed-phase and soap TLC of amino acids (195) 65

Amino acids
GC of trimethylsilylesters of trifluoroacetylamino acids. III. Simple method for the preparation of the trimethylsilylesters of N-trifluoroacetyl amino acids (in German) (196) 160

Amino acids
GC separation of 3-methylhistidine (196) 166

Amino acids
Fast determination of free amino acids by ion-pair HPLC using on-line post-column derivatization (196) 319

Amino acids
Model studies of the reactions and non-specific adsorptive properties of cyanogen bromide-activated macroporous agarose (196) 379

Amino acids
Quantitative elution of thyroidal amino acids from a cation-exchange column: analysis of tablet formulations containing dyes (196) 512

Amino acids
Effect of derivative structure on flame-ionization detector response of amino acid oxazolidinones (197) 226

Amino acids
Rapid HPLC determination of amino acids in synaptosomal extracts (198) 516

Amino acids
Reversed-phase HPLC resolution of D- and L-Dns-amino acids by mixed chelate complexation (199) 295

Amino acids
The enantiomer as internal standard for the quantitation of the alkylated amino acid S-methyl-L-cysteine in haemoglobin by GC-chemical ionisation MS with single ion detection (200) 145

Amino acids
The isolation of methionine and ethionine by silver ligand chromatography and application to methionine-containing peptides (200) 185

Amino acids
Resolution of amino acid enantiomers by glass capillary GC on easily prepared optically active stationary phases (200) 195

Amino alcohols
Chromatography of organometallic and organometalloidal derivatives of amino

alcohols. II. GC evidence for transannular bonds in solutes (200) 105

Aminoanthracene
Separation and detection of 2-aminoanthracene and its metabolites by HPLC (195) 392

Aminoanthraquinones
HPLC of cancer chemotherapeutic agents: bis(substituted aminoalkylamino)anthraquinones (187) 212

Aminobenzamidine
Interaction of trypsin with immobilized *p*-aminobenzamidine derivatives studied by means of affinity electrophoresis (194) 175

Aminocarb
Analysis of aminocarb formulations by HPLC (190) 208

Aminocarb
Detection of aminocarb and its major metabolites by TLC (194) 100

Aminocarb
Simplified method for the analysis of some carbamate insecticides in foliage, forest soil and fish tissue by direct GLC (200) 179

Aminocarb
GLC analysis of aminocarb and seven derivatives (200) 228

Aminoglycoside antibiotics
Aminoglycoside antibiotics: TLC bioautographic detection and quantitative assay (195) 251

(Aminohexyl)aminoadenosine monophosphate
Control of 8-bromo-AMP and 8-(6-aminohexyl)amino-AMP synthesis by HPLC (196) 323

Aminolaevulinic acid
Determination of 5-aminolaevulinic acid and porphobilinogen by HPLC (185) 605

Aminopyrine
Quantitative HPLC determinations of aminopyrine and its metabolites in man (181) 51

Aminosalicylic acid
Non-extractive fluorometric measurement of *p*-aminosalicylic acid in plasma by ion-pairing techniques and HPLC (181) 266

Amino sugars
Separation of amino sugars and related compounds by two-dimensional TLC (195) 127

Amiodarone
Rapid HPLC method for the measurement of amiodarone in blood plasma or serum at the concentrations attained during therapy (187) 391

Amitriptyline
Determination of amitriptyline-N-oxide, amitriptyline and nortriptyline in serum and plasma by HPLC (183) 321

Amitriptyline
LC of amitriptyline and related tricyclic compounds (197) 246

Amitrole
Determination of amitrole by ion-pair HPLC (200) 234

Ammonium compounds
Analysis of quaternary ammonium compounds and basic drugs based on ion-pair adsorption HPLC (186) 683

Amperometric detector
Design and selective application of a dropping mercury electrode amperometric detector in column LC (186) 489

Amphetamine
Microdetermination of stimulant drugs in urine by HPLC (196) 334

Ampholines
Gel chromatographic comparison of the molecular weight distributions of Ampholine, Servalyte and Pharmalyte carrier ampholytes used in isoelectric focusing (198) 301

Ampholyte-displacement chromatography
An approach to ampholyte-displacement chromatography (200) 57

Amrinone
HPLC analysis of amrinone and its N-acetyl derivative in plasma. Pharmacokinetics of amrinone in the dog (187) 264

Amygdalin
Improved GC of amygdalin and its diastereomer (197) 240

Anabolic drugs
Determination of residues of anabolic drugs in meat by GC-MS (195) 231

Anaestetic drugs
Capillary GC with a nitrogen detector for measurement of phencyclidine, ketamine and other arylcycloalkylamines in the picogram range (193) 157

Anaesthetic drugs
Positive identification and quantitation of isomeric cocaines by HPLC (193) 371

Anaesthetics
GC determination of bromo and fluoro derivatives of benzodiazepine in human body fluids (182) 81

Anaesthetics
Simultaneous determination of bupivacaine and 2,6-pipecoloxylidide in serum by GLC (182) 226

Analgesics
Quantitative HPLC determinations of aminopyrine and its metabolites in man (181) 51

Analgesics
HPLC separation and isolation of quinidine and quinine metabolites in rat urine (181) 219

Analgesics
Improvements to a HPLC assay for allopurinol and oxipurinol in plasma (181) 286

Analgesics
Rapid HPLC method for the determination of probenecid in biological fluids (182) 252

Analgesics
HPLC determination of indoprofen in plasma and urine (182) 262

Analgesics
HPLC separation of colchicine and its phenolic and N-desacetylated derivatives (188) 280

Analgesics
Reversed-phase HPLC analysis of cinchona alkaloids in pharmaceuticals (189) 241

Analgesics
Purity determination of zomepirac sodium dihydrate by HPLC (190) 429

Analgesics
Measurement of the formation of paracetamol and *p*-nitrophenol glucuronides *in vitro*, by ion-pair HPLC (192) 247

Analgesics
Sample-solvent-induced peak broadening in the reversed-phase HPLC of Aspirin and related analgesics (194) 217

Androstenes
Intermediate- and large-scale reversed-phase preparative HPLC on an axially compressed column: a facile, quantitative separation of 7α- and 7β-methyl-17β-acetoxy-3-oxoandrost-4-enes (194) 239

Angiotensins
Separation and quantification of angiotensins and some related peptides by HPLC (183) 303

Anhydroglucitol
Qualitative TLC separation of 1,5-anhydroglucitol in the presence of other carbohydrates on silica gel impregnated with borate buffer (187) 455

Anhydrooxytetracycline
Reversed-phase ion pair chromatography of oxytetracycline, epioxytetracycline and anhydrooxytetracycline (190) 486

Anilines
Determination of halogenated anilines in urine by HPLC with an electrochemical detector (188) 412

Anion chromatography
Anion chromatography with low-conductivity eluents (186) 509

Anion-exchange resins
Chromatographic behaviour of aromatic compounds on anion-exchange resins in various carboxylate forms (198) 267

Anions
Anion chromatography with low-conductivity eluents. II (187) 35

Anions
Determination of some anions separated by TLC (187) 462

Anions
Role of the charge number of the counter-ionic constituent in the separation of anions by isotachophoresis (194) 11

Anomer separation
Improved separation of anomers of saccharides by HPLC on macroreticular anion-exchange resin in the sulphate form (192) 452

Anthelmintic agents
Determination of avermectins in plasma at nanogram levels using HPLC with fluorescence detection (190) 367

Anthelmintic drugs
Two HPLC determinations for mebendazole and its metabolites in human plasma using a rapid Sep Pak C18 extraction (183) 311

Anthelmintic drugs
Extractive alkylation of 5,2'-dichloro-4-nitrosalicylanilide (niclosamide) for GLC analysis (189) 375

Anthracyclines
HPLC and TLC of anthracycline antibiotics. Separation and identification of components of the daunorubicin complex from fermentation broth (198) 407

Anti-anginal drugs
Analysis of verapamil in plasma by LC (182) 473

Anti-anginal drugs
HPLC in the elucidation of the metabolic fate of butoprozine. Screening for unknown metabolites in the rat (191) 109

Anti-arrhythmic drugs
Sensitive GC assay for the quantitation of bretylium in plasma, urine and myocardial tissue (181) 33

Anti-arrhythmic drugs
Simple and selective HPLC method for estimating plasma quinidine levels (181) 207

Anti-arrhythmic drugs
HPLC separation and isolation of quinidine and quinine metabolites in rat urine (181) 219

Anti-arrhythmic drugs
Mexiletine analysis in blood and plasma using GC and nitrogen-selective detection (181) 469

Anti-arrhythmic drugs
Comparison of HPLC with an enzyme multiplied immunoassay technique for the

analysis of serum procainamide and N-acetylprocainamide (181) 490

Anti-arrhythmic drugs
Analysis of verapamil in plasma by LC (182) 473

Anti-arrhythmic drugs
Simultaneous analysis of disopyramide and quinidine in plasma by HPLC (183) 65

Anti-arrhythmic drugs
HPLC assay for tocainide in human plasma: comparison with GLC assay (183) 109

Anti-arrhythmic drugs
Rapid HPLC method for the measurement of amiodarone in blood plasma or serum at the concentrations attained during therapy (187) 391

Anti-asthmatic drugs
New rapid assay of theophylline in plasma by isotachophoresis (181) 147

Anti-asthmatic drugs
HPLC of the oral anti-asthmatic agent pirquinozol and its active metabolite (200) 282

Antibacterial agents
GC determination of piromidic acid (193) 476

Antibacterial agents
HPLC analysis of furazolidone in liver and kidney (198) 526

Antibiotics
Reversed-phase HPLC assay for cefoxitin in proteinaceous biological samples (181) 120

Antibiotics
HPLC micro-assay for chloramphenicol in human blood plasma and cerebrospinal fluid (181) 123

Antibiotics
High-performance paired-ion LC determination of bleomycin A_2 in urine (181) 127

Antibiotics
Determination of mecillinam in urine by reversed-phase HPLC (181) 272

Antibiotics
Micro determination of gentamicin in serum by HPLC with UV detection (182) 201

Antibiotics
HPLC determination of ceftizoxime, a new cephalosporin antibiotic, in rat serum, bile and urine (182) 448

Antibiotics
HPLC determination of chloramphenicol and its monosuccinate ester in plasma (182) 459

Antibiotics
Analysis of isoxazolyl penicillins and their metabolites in body fluids by HPLC (183) 339

Antibiotics
HPLC assay for measurement of cefoxitin in serum (183) 357

Antibiotics
LC determination of tetracycline in plasma and urine (185) 583

Antibiotics
Mechanism of reversed-phase separation of tetracyclines by HPLC (186) 763

Antibiotics
Rapid analysis of doxycycline from biological samples by HPLC (187) 439

Antibiotics
Improved HPLC method for the comparison of heptaene macrolide antibiotics (189) 249

Antibiotics
Determination of adriamycin and daunorubicin in urine by HPLC with laser fluorometric detection (190) 377

Antibiotics
Reversed-phase ion pair chromatography of oxytetracycline, epioxytetracycline and anhydrooxytetracycline (190) 486

Antibiotics
Determination of adriamycin in liposomes by HPLC using a fluorescence detector (190) 489

Antibiotics
Analytical studies of maridomycin. III. Spectrophotometric determination of maridomycins and 9-propionylmaridomycins as iodine complexes after TLC (190) 493

Antibiotics
Confirmatory identification of Susceptibility Card antibiotics by ion-pair and reversed-phase HPLC (192) 467

Antibiotics
TLC of macrolide antibiotics (193) 119

Antibiotics
Fast and simple method for the separation of intermediates and cofactors involved in the biosynthesis of cephalosporin C using chemically bonded C_{12} reversed-phase TLC (193) 511

Antibiotics
Aminoglycoside antibiotics: TLC bioautographic detection and quantitative assay (195) 251

Antibiotics
Use of the image analyser Optomax for the quantitative evaluation of antibiotics separated by gel electrophoresis and by TLC (195) 297

Antibiotics
Determination of the concentration of adriamycin and its metabolites in the serum and tissues of Ehrlich carcinoma-bearing mice by HPLC (196) 463

Antibiotics
HPLC and TLC of anthracycline antibiotics. Separation and identification of components of the daunorubicin complex from fermentation broth (198) 407

Antibiotics
Separation of tunicamycin homologues by reversed-phase HPLC (198) 506

Antibiotics
HPLC analysis of furazolidone in liver and kidney (198) 526

Anticholinergic drugs
Quantitative determination of benzilonium bromide in plasma by GC-MS after oxidation to benzophenone (183) 511

Anticholinergic drugs
Determination of aprophen and its hydrolytic by-products by ion-pair HPLC (195) 146

Anticholinergic drugs
Determination of adiphenine hydrochloride and diphenylacetic acid by ion-pair HPLC (200) 267

Anticonvulsant drugs, see also anti-epileptic drugs

Anticonvulsant drugs
Simultaneous LC analysis for carbamazepine and carbamazepine 10,11-epoxide in plasma and saliva by use of double internal standardization (181) 373

Anticonvulsant drugs
Simultaneous quantitation of chlormethiazole and two of its metabolites in blood and plasma by GLC (182) 55

Anticonvulsant drugs
Application of an automated extractor/concentrator to the analysis of anticonvulsant drugs in blood serum by HPLC (185) 505

Antidepressant drugs
Simplified method for determination of the tetracyclic antidepressant mianserin in human plasma using GC with nitrogen detection (181) 25

Antidepressant drugs
Comparative statistical study of assay methods using mass fragmentography and GC with nitrogen detection for determination of the tetracyclic antidepressant mianserin in human plasma (181) 115

Antidepressant drugs
Analysis of tricyclic antidepressant drugs by GC using nitrogen-selective detection with packed and capillary columns (182) 349

Antidepressant drugs
Determination of tranylcypromine in human plasma and urine using high-resolution GLC with nitrogen-sensitive detection (183) 25

Antidepressant drugs
Chemical ionisation mass fragmentographic measurement of dothiepin plasma concentrations following a single oral dose in man (183) 141

Antidepressant drugs
Technical considerations in the GC analysis of desipramine (183) 149

Antidepressant drugs
The use of a pre-column for the direct HPLC determination of the antidepressants clovoxamine and fluvoxamine in plasma (183) 203

Antidepressant drugs
Determination of amitriptyline-N-oxide, amitriptyline and nortriptyline in serum and plasma by HPLC (183) 321

Antidepressant drugs
Reversed-phase chromatography of zimelidine and similar dibasic amines. I. Analysis in biological material (185) 593

Antidepressant drugs
Analysis of quaternary ammonium compounds and basic drugs based on ion-pair adsorption HPLC (186) 683

Antidepressant drugs
Peak tailing and retention behaviour of tricyclic antidepressant amines and related hydrophobic ammonium compounds in reversed-phase ion-pair LC on alkyl-bonded phases (189) 299

Antidepressant drugs
GLC determination of 4'-hydroxynomifensine in biological samples (190) 475

Antidepressant drugs
Determination of *cis*- and *trans*-isomers of dosulepine and dithiadene by HPLC (191) 137

Antidepressant drugs
LC of amitriptyline and related tricyclic compounds (197) 246

Antidiabetic drugs
Electron-capture GC of plasma sulphonylureas after extractive methylation (181) 17

Antidiabetic drugs
Gel chromatographic separation of insulin analogues in human serum (181) 248

Antidiabetic drugs
Detection of glibornurid in serum by reversed-phase HPLC (in German) (183) 115

Antidiabetic drugs
Determination of gliclazide in human serum by HPLC using an anion-exchange resin (183) 467

Antidotes
GC analysis of naloxone in biological fluids (183) 352

Anti-emetic drugs
Electron-capture GC assay for metoclopramide in plasma (183) 175

Anti-epileptic drugs, see also anticonvulsant drugs

Anti-epileptic drugs
HPLC separation of carbamazepine metabolites excreted in rat urine (181) 59

Anti-epileptic drugs
Microassay for primidone and its metabolites phenylethylmalondiamide, phenobarbital, and *p*-hydroxyphenobarbital in human serum, saliva, breast milk and tissues by GC-MS using selected ion monitoring (182) 71

Anti-epileptic drugs
Improved GLC method for the simultaneous determination of phenobarbital, phenytoin, carbamazepine and primidone in biological fluids (182) 116

Anti-epileptic drugs
Simple, rapid procedure for the determination of valproate and ethosuximide in plasma by GLC (182) 237

Anti-epileptic drugs
GC determination of valproic acid in human plasma (182) 241

Anti-epileptic drugs
Determination of midazolam in serum by GC with a nitrogen-selective detector (182) 439

Anti-epileptic drugs
Direct determination of valproate in serum by isotachophoresis. Comparison with a GC method (182) 496

Anti-epileptic drugs
A quantitative method for the simultaneous determination of carbamazepine, mephenytoin, phenylethylmalonamide, phenobarbital, phenytoin and primidone in serum by thin-layer chromatography. Additional comments to the analysis of carbamazepine-10,11-epoxide (183) 387

Anti-epileptic drugs
HPLC and GLC determination of diazepam and nordiazepam in plasma (183) 441

Anti-epileptic drugs
Rapid method for simultaneous determination of phenobarbital, diphenylhydantoin and their main hydroxylated metabolites by nitrogen-selective GC (183) 505

Antihistamine drugs
Plasma and urine concentration of methapyrilene by nitrogen-phosphorus GLC (183) 234

Antihistamine drugs
Determination of cinnarizine in plasma by HPLC (183) 378

Antihistamine drugs
Determination of *cis*- and *trans*-isomers of dosulepine and dithiadene by HPLC (191) 137

Antihistamine drugs
Rapid, sensitive determination of unchanged promethazine in biological material using a nitrogen-selective flame ionization detector. Identification of metabolites by GC-MS (192) 375

Antihypertensive drugs
Selective HPLC assays for hydralazine and its metabolites in plasma of man (181) 427

Antihypertensive drugs
Analysis of verapamil in plasma by LC (182) 473

Antihypertensive drugs
GC method for the simultaneous determination of hydralazine and its acetylated metabolite in serum using a nitrogen-selective detector (183) 159

Antihypertensive drugs
Determination of hydralazine metabolites: 4-hydrazino-phthalazin-1-one and N-acetylhydrazinophthalazin-1-one by GC and *s*-triazolo[3,4-*α*]phthalazine and phthalazinone by HPLC (183) 167

Antihypertensive drugs
Determination of propranolol and its major metabolites in plasma and urine by HPLC without solvent extraction (183) 213

Antihypertensive drugs
HPLC determination of prazosin in human plasma, whole blood and urine (183) 367

Antihypertensive drugs
Determination of clonidine in human plasma by glass capillary GC with electron-capture detection (187) 161

Antihypertensive drugs
Quantitative analysis of hydralazine pyruvic acid hydrazone, the major plasma metabolite of hydralazine (187) 171

Antihypertensive drugs
Determination of hydralazine and its acetylated metabolites in urine by GC and HPLC (187) 218

Antihypertensive drugs
Detection of some antihypertensive drugs and their metabolites in urine by TLC. Five commonly used beta blockers and hydralazine (187) 277

Antihypertensive drugs
Formation of electron-capturing derivatives of alprenolol by transboronation. Application to the determination of alprenolol in plasma (194) 365

Antihypertensive drugs
Detection of some antihypertensive drugs and their metabolites in urine by TLC. II. A further five beta blockers and dihydralazine (196) 189

Anti-inflammatory agents
Reversed-phase HPLC of phenylbutazone in body fluids (181) 132

Anti-inflammatory agents
Convenient and sensitive HPLC assay for ketoprofen, naproxen and other allied drugs in plasma or urine (190) 119

Anti-inflammatory agents
Quantitative determination of flufenamic acid in rat plasma and uterus by GC (192) 441

Anti-inflammatory agents
Determination of total monohydroxylated metabolites of diclofenac in urine by electron-capture GLC (195) 421

Anti-inflammatory agents
Analysis of hydrocortisone acetate ointments and creams by HPLC (198) 41

Anti-inflammatory drugs
Rapid determination of indomethacin and salicylic acid in serum by means of reversed-phase LC (181) 385

Anti-inflammatory drugs
Specific and sensitive method for the determination of aspirin and salicylic acid in plasma using reversed-phase HPLC (181) 473

Anti-inflammatory drugs
Quantitative analysis of oxepinac in human plasma, urine and saliva by GC-mass fragmentography (182) 47

Anti-inflammatory drugs
HPLC determination of indoprofen in plasma and urine (182) 262

Anti-inflammatory drugs
Determination of isofezolac in biological fluids by reversed-phase LCC (182) 369

Anti-inflammatory drugs
Determination of benoxaprofen in plasma and urine by LC (182) 478

Anti-inflammatory drugs
Rapid HPLC method for the quantitative determination of diflunisal in plasma (182) 487

Anti-inflammatory drugs
Improved GC method of determining diclofenac in plasma (183) 41

Anti-inflammatory drugs
Analysis of piroxicam in plasma by HPLC (183) 104

Anti-inflammatory drugs
High-performance liquid column chromatography of fenoprofen in serum (183) 250

Anti-inflammatory drugs
HPLC of the topical anti-inflammatory steroid halcinonide (190) 481

Antilipolytic agents
Hydrophobic chromatography and bioanalysis of some polar pyridine derivatives used as antilipolytic agents (187) 189

Antineoplastic agents
Application of an electrochemical detector to the determination of procarbazine hydrochloride by HPLC (190) 359

Antineoplastic agents
GC determination of 5-alkyl-uracils and 5-alkyl-deoxyuridines using flash methylation and glass capillary columns (191) 253

Antineoplastic drugs, see also antitumour drugs and cytostatic drugs

Antineoplastic drugs
Determination of two new metabolites of 1-hexylcarbamoyl-5-fluorouracil in biomedical specimens by HPLC (182) 125

Antineoplastic drugs
Analysis of the anticancer drugs VP 16-213 and VM 26 and their metabolites by HPLC (182) 211

Antineoplastic drugs
Improved GC of amygdalin and its diastereomer (197) 240

Antineoplastic drugs
TLC separation of potential antineoplastic agents: 1-ethoxycarbonyl-2-arylazo-2-nitroethanes (198) 224

Antineoplastic drugs
HPLC and TLC of anthracycline antibiotics. Separation and identification of components of the daunorubicin complex from fermentation broth (198) 407

Anti-oestrogenic drugs
Paired-ion chromatographic analysis of tamoxifen and two major metabolites in plasma (181) 41

Antioxidant
Determination of 2,6-di-(*tert.* -butyl)-4-methylphenol in rubber-base materials and identification of an oxidation product from 2,6-di-(*tert.* -butyl)-4-methylphenol (190) 193

Anti-oxidants
GLC retention indices of 296 non-drug substances on SE-30 or OV-1 likely to be encountered in toxicological analyses (Review) (184) 185

Anti-oxidants
HPLC determination of *tert.* -butylhydroquinone in vegetable oils (187) 436

Antirheumatic drugs
Reversed-phase HPLC of phenylbutazone in body fluids (181) 132

Antirheumatic drugs
Estimation of serum indomethacin at therapeutic levels by means of TLC and spectrophotometry (181) 282

Antirheumatic drugs
Rapid determination of indomethacin and salicylic acid in serum by means of reversed-phase LC (181) 385

Antirheumatic drugs
Quantitation of ibuprofen in biological fluids by GC-MS (181) 463

Antirheumatic drugs
Quantitative determination of diclofenac sodium salt in plasma by absorption spectroscopy with direct quantitation of thin-layer chromatograms (in German) (181) 512

Antirheumatic drugs
Improved GC method of determining diclofenac in plasma (183) 41

Antirheumatic drugs
Analysis of piroxicam in plasma by HPLC (183) 104

Antirheumatic drugs
Simultaneous determination of phenylbutazone and its metabolites in plasma and urine by HPLC (183) 331

Antirheumatic drugs
HPLC determination of benzbromarone and the main metabolite benzarone in serum (183) 383

Antispasmodic agents
Determination of adiphenine hydrochloride and diphenylacetic acid by ion-pair HPLC (200) 267

Antispasmodic drugs
Determination of aprophen and its hydrolytic by-products by ion-pair HPLC (195) 146

Antithyroid drugs
Determination of methimazole in plasma using GC-MS after extractive alkylation (182) 63

Antitumour agents
HPLC assay for methylglyoxal bis(guanylhydrazone) (methyl GAG) in plasma and urine (187) 101

Antitumour agents
HPLC of cancer chemotherapeutic agents: bis(substituted aminoalkylamino)anthraquinones (187) 212

Antitumour agents
Quantitative determination of 1,3-bis(tetrahydro-2-furanyl)-5-fluoro-2,4-pyrimidinedione and its metabolites in visceral tissues by HPLC and GC mass fragmentography (188) 270

Antitumour agents
HPLC separation of colchicine and its phenolic and N-desacetylated derivatives (188) 280

Antitumour agents
Rapid quantitative assay for 3,6-bis-(5-chloro-2-piperidyl)-2,5-piperazinedione in fermentation broth using HPLC (190) 246

Antitumour drugs, see also cytostatic drugs and antineoplastic drugs

Antitumour drugs
Determination of 6-diazo-5-oxo-L-norleucine in plasma and urine by reversed-phase HPLC of the dansyl derivative (181) 95

Antitumour drugs
Determination of hexamethylmelamine and metabolites in plasma or serum by GLC with a nitrogen-sensitive detector (181) 363

Antitumour drugs
Quantitation of the antitumor agent N-(phosphonacetyl)-L-aspartic acid in human plasma and urine by GC-MS-selected ion monitoring (182) 163

Antitumour drugs
Some aspects of the GLC analysis of cyclophosphamide in plasma (183) 49

Antitumour drugs
HPLC analysis of the anticancer agent methylglyoxal bis(guanylhydrazone) (MGBG, NSC-32946) in biological fluids (183) 363

Antitumour drugs
Plasma and tissue concentrations of hycanthone in the mouse determined by reversed-phase HPLC (183) 372

Antitumour drugs
Determination of adriamycin and daunorubicin in urine by HPLC with laser fluorometric detection (190) 377

Antitumour drugs
Determination of adriamycin in liposomes by HPLC using a fluorescence detector (190) 489

Apiezon M, hydrogenated
Evaluation of a thermally stable hydrocarbon as a non-polar base stationary phase for use in Rohrschneider-McReynolds-type schemes (186) 89

Apomorphine
Determination of apomorphine in plasma and brain tissue by ion-pair extraction and LC (185) 575

Aprophen
Determination of aprophen and its hydrolytic by-products by ion-pair HPLC (195) 146

Arabinofuranosidase
Purification of an *α*-L-arabinofuranosidase from *Aspergillus niger* by substrate affinity chromatography (197) 86

Arabinofuranosylcytosine
HPLC analysis of cytosine arabinoside and metabolites in biological samples (183) 449

Argon plasma emission source detector
Capacilities of on-line element-specific detection in HPLC using an inductively coupled argon plasma emission source detector (185) 549

Aroclor 1242
Retention behavior of chlorinated benzenes, chlorinated phenols, Aldrin, and Aroclor 1242

on Tenax-GC, Chromosorb 101, Florisil, and Carbopack C HT (200) 115

Aromatic acids
Chromatography in aqueous solution on styrene-divinylbenzene resins (193) 197

Aromatic compounds
Chromatographic behaviour of aromatic compounds on Sephadex LH gels. Calibration of gel columns for determination of molecular weight distributions (198) 131

Aromatic compounds
Chromatographic behaviour of aromatic compounds on anion-exchange resins in various carboxylate forms (198) 267

Aromatic hydrocarbons
Application of head-space analysis to the study of volatile organic impurity concentrates. GC determination of aromatic hydrocarbons in atmospheric air (186) 851

Aromatic nitro compounds
Reversed-phase LC of aromatic nitro compounds on a macroporous polystyrene gel (197) 147

Arsenate
Graphite furnace atomic absorption spectrophotometers as automated element-specific detectors for HPLC. The determination of arsenite, arsenate, methylarsonic acid and dimethylarsinic acid (191) 31

Arsenite
Graphite furnace atomic absorption spectrophotometers as automated element-specific detectors for HPLC. The determination of arsenite, arsenate, methylarsonic acid and dimethylarsinic acid (191) 31

Artifacts
Decreased accuracy in isotachophoretic analysis due to septum bleed (188) 262

Artifacts
Complexation with ferric ions: a potential source of error in the isotachophoretic determination of oxalate (188) 266

Artifacts
Background contamination by phthalates commonly encountered in the chromatographic analysis of lipid samples (189) 421

Artifacts
Analysis errors following hydrogen cleaning of an electron capture detector (190) 133

Artifacts
Correction for artifacts in the analysis of atmospheric aerosols (190) 136

Artifacts
Correction for deviation from the Lambert-Beer law in the quantitation of thin-layer chromatograms by photodensitometry (192) 208

Artifacts
Anomalous partition behaviour of sodium dodecyl sulphate monomer in Sephadex gels with high dextran concentrations. Possible role of water of hydration in the gel phase (193) 345

Artifacts
UV-induced silica gel GF photoluminescence studied by liquid scintillation spectrophotometry (195) 281

Artifacts
Syringe and column adsorption of tertiary amines in GC (196) 489

Artifacts
Error incurred in gel permeation chromatography by using the elution peak volume in lieu of the elution mean volumes in the calculation of K_{av} (198) 354

Artifacts
Correction of R_F values according to Galanos and Kapoulas (in German) (198) 357

Arylaliphatic acids
R_M values from reversed-phase TLC as parameters of lipophilicity in quantitative structure-activity relationships in four series of arylaliphatic acids (195) 329

Arylcycloalkylamines
Capillary GC with a nitrogen detector for measurement of phencyclidine, ketamine and other arylcycloalkylamines in the picogram range (193) 157

Arylhydroxylamines
Conversion of arylhydroxylamines to electrochemically-active derivatives suitable for HPLC analysis with amperometric detection (188) 159

Asarone
Reversed-phase HPLC for the determination of β-asarone (194) 245

Ascorbate phosphate
HPLC determination of L-ascorbate-2-phosphate in phosphorylation reactions (187) 421

Ascorbic acid
Application of HPLC to the separation of ascorbic acid from isoascorbic acid (196) 163

Ascorbic acid
Chemiluminescence HPLC detector applied to ascorbic acid determination (200) 153

Aspartic acid
MS technique for the determination of N-phosphonoacetyl-L-aspartic acid in serum (181) 195

Aspartic acid
GLC of free amino acids in the hyaloplasm of

rat cerebral, cerebellar and ocular tissues, and in skeletal and heart muscle (182) 147

Aspartic acid
Effect of some organic buffers on the estimation of aspartic acid and resolution in amino acid analysis (194) 76

Aspartylglucosamine
Automated ion-exchange chromatography in the detection of aspartylglucosaminuria (182) 409

Aspirin
Specific and sensitive method for the determination of aspirin and salicylic acid in plasma using reversed-phase HPLC (181) 473

Aspirin
Sample-solvent-induced peak broadening in the reversed-phase HPLC of Aspirin and related analgesics (194) 217

Association constants
Measurement of association constants for complexes by reversed-phase HPLC (186) 371

Atenolol
Electron-capture-GC determination of atenolol in plasma and urine, using a simplified procedure with improved selectivity (182) 341

Atomic absorption detector
Application of a graphite furnace atomic absorption detector automatically coupled to a high-performance liquid chromatograph for speciation of metal-containing macromolecules (185) 563

Atropine
Continuous post-column ion-pair extraction detection of some basic organic compounds in normal-phase chromatography (185) 473

Avermectins
Determination of avermectins in plasma at nanogram levels using HPLC with fluorescence detection (190) 367

Aza-arenes
Fractionation and GC analysis of aza-arenes in complex mixtures (199) 371

Azapurines
Determination of selected pyrimidines, purines and their metabolites in serum and urine by reversed-phase ion-pair chromatography (199) 345

Azathioprine
Stability-indicating assay for azathioprine and 6-mercaptopurine by reversed-phase HPLC (186) 691

Azo-pigments
Separation of some azo-pigments from their diazo- and coupling components by chromatography on Sephadex LH-20 with N,N-dimethylformamide as eluent (187) 232

Azosemide
Separation and analysis of azosemide in urine and in serum by HPLC (182) 257

B

Band dispersion
Band dispersion in high-performance exclusion chromatography (185) 289

Barban
Direct analysis of the herbicide barban in wheat products by reversed-phase LC (195) 113

Barban
Analysis of barban formulations by HPLC (195) 151

Barbiturates
Improved detectability of barbiturates in HPLC by pre-column labelling and UV detection (186) 535

Barbiturates
Analysis of therapeutic and commonly abused drugs in serum and urine by GLC using a photoionization detector (186) 637

Barbiturates
Analysis of trace amounts of barbiturates in saliva (190) 113

Barbiturates
GC separations of dialkyl barbiturate derivatives (192) 212

Barbiturates
Glass capillary column GC of barbiturates after flash-heater derivatization with dimethylformamide dimethylacetal (192) 363

Barbiturates
Direct derivatization and GC determination of barbiturates in autopsy liver tissues (195) 349

Barbiturates
Use of dual-wavelength UV detection in HPLC for the identification of barbiturates (200) 271

Basic drugs
Analysis of quaternary ammonium compounds and basic drugs based on ion-pair adsorption HPLC (186) 683

Basic drugs
Isocratic multi-column HPLC as a technique for qualitative analysis and its application to the characterisation of basic drugs using an aqueous methanol solvent (187) 65

Beer bitter acids
HPLC of beer bitter acids (197) 189

Benoxaprofen
Determination of benoxaprofen in plasma and urine by LC (182) 478

Bentonites
Inclusion compounds in chromatography (Review) (184) 347

Benzamide
TLC method for the separation and determination of the products of the reaction of amides with formaldehyde (195) 158

Benzarone
HPLC determination of benzbromarone and the main metabolite benzarone in serum (183) 383

Benzazole derivatives
Simple rapid TLC method for the detection of optical brighteners in polymers (194) 434

Benzbromarone
HPLC determination of benzbromarone and the main metabolite benzarone in serum (183) 383

Benzene
Determination of 1,2-dibromoethane, 1,2-dichloroethane and benzene in ambient air using porous polymer traps and GC-MS analysis with selected ion monitoring (190) 97

Benzene derivatives
Retention behaviour of *o*-, *m*- and *p*-isomers of benzene derivatives on a silica gel hydroxylated surface in LC (195) 205

Benzenesulphonates
Inclusion compounds in chromatography (Review) (184) 347

Benzilonium bromide
Quantitative determination of benzilonium bromide in plasma by GC-MS after oxidation to benzophenone (183) 511

Benzodiazepine derivatives
TLC of the acid hydrolysis products of nineteen benzodiazepine derivatives (194) 262

Benzodiazepines
Determination of some benzodiazepines and metabolites in serum, urine and saliva by HPLC (181) 227

Benzodiazepines
GC determination of bromo and fluoro derivatives of benzodiazepine in human body fluids (182) 81

Benzodiazepines
Determination of midazolam in serum by GC with a nitrogen-selective detector (182) 439

Benzodiazepines
LC identification of clorazepate in pharmaceutical products (188) 426

Benzodiazepines
Screening procedure for therapeutic benzodiazepines by HPLC of their benzophenones (189) 94

Benzodiazepines
GC determination of two fluorinated benzodiazepines in rats and mice (198) 180

Benzodioxole
Chromatography of 1,3-benzodioxole derivatives (193) 19

Benzoic acid
GC determination of low concentrations of benzoic acid in human plasma and urine (181) 161

Benzoic acid
Alkylation of carboxylic acids by solid-liquid phase-transfer catalysis for determination by GC. II. (196) 255

Benzoic acid derivatives
Separation of benzoic acid derivatives from plant extracts on Sephadex G-10 columns (in German) (198) 373

Benzophenones
Screening procedure for therapeutic benzodiazepines by HPLC of their benzophenones (189) 94

Benzopyrene
Determination of benzo[*a*]pyrene and other polyaromatic hydrocarbons in petroleum oils by direct LC (190) 85

Benzopyrene metabolites
Reversed-phase separation of benzo[*a*]pyrene metabolites by TLC (197) 217

Benzothiepines
Determination of *cis*- and *trans*-isomers of dosulepine and dithiadene by HPLC (191) 137

Benzoyl ecgonine
Qualitative screening procedure for the detection of benzoyl ecgonine in the urine of greyhounds (196) 337

Beta-adrenergic receptor blocking drugs
Determination of plasma mepindolol levels by HPLC and electrochemical detection (181) 67

Beta-adrenergic receptor blocking drugs
Quantification of oxprenolol in biological fluids using HPLC (181) 135

Beta-adrenergic receptor blocking drugs
Electron-capture-GC determination of atenolol in plasma and urine, using a simplified procedure with improved selectivity (182) 341

Beta-adrenergic receptor blocking drugs
Determination of metoprolol metabolites in plasma and urine by electron-capture GLC (183) 92

Beta-adrenergic receptor blocking drugs
Sensitive GC method for the determination in plasma of FM 24, 1-(2-*exo*-bicyclo[2,2,1]hept-2-ylphenoxy)-3-[(1-methylethyl)amino]-2-

propanol, a new β-adrenoceptor blocking agent (183) 99

Beta-adrenergic receptor blocking drugs
Determination of propranolol and its major metabolites in plasma and urine by HPLC without solvent extraction (183) 213

Beta-adrenergic receptor blocking drugs
Determination of pindolol in human plasma by HPLC (183) 255

Beta-adrenergic receptor blocking drugs
Detection of some antihypertensive drugs and their metabolites in urine by TLC. Five commonly used beta blockers and hydralazine (187) 277

Beta-adrenergic receptor blocking drugs
Determination of isoprenaline by ion-pair HPLC (189) 272

Beta-adrenergic receptor blocking drugs
Formation of electron-capturing derivatives of alprenolol by transboronation. Application to the determination of alprenolol in plasma (194) 365

Beta-adrenergic receptor blocking drugs
Detection of some antihypertensive drugs and their metabolites in urine by TLC. II. A further five beta blockers and dihydralazine (196) 189

Beta-adrenergic receptor blocking drugs
Semi-automated HPLC method for the determination of diacetolol hydrochloride in medicated animal diets (200) 261

Beta-induced fluorescence detector
Beta-induced fluorescence as a detection technique for LC. II. Prototype detector for normal-phase operation (200) 47

Betamethasone
Simultaneous determination of betamethasone, betamethasone acetate and hydrocortisone in biological fluids using HPLC (183) 131

Beta-stimulant
Separation of the diastereoisomers of 1-(4-hydroxy-3-methoxymethylphenyl)-2-N-(*p*-methoxyphenylisopropyl)aminoethanol by HPLC (193) 483

Bifunctional compounds
Cyclic derivatives for the selective chromatographic analysis of bifunctional compounds (Review) (184) 99

Bifunctional compounds
Comparison of electron-capturing boronic acids for the selective analysis of bifunctional compounds (186) 307

Bile acids
Rapid determination of glycine- and taurine-conjugated bile acids in human bile by HPLC (181) 177

Bile acids
Preparation and characterisation of permethylated derivatives of bile acids, and their application to GC analysis (183) 269

Bile acids
Reversed-phase HPLC separation and quantification of individual human bile acids (183) 277

Bile acids
Quantitative determination of bile acids in bile with reversed-phase HPLC (183) 287

Bile acids
GC of dimethylalkylsilyl ether derivatives of bile acid methyl esters (190) 339

Bile acids
GLC separation of bile acids and steroids on a nematic liquid crystal (193) 444

Bile acids
Separation of bile acids on Chromarods: a new qualitative and semi-quantitative TLC technique (193) 504

Bile acids
New solvent systems for separation of free and conjugated bile acids. III. Separation of bile acid methyl ester acetates (196) 150

Bile acid sulphates
Measurement of faecal bile acid sulphates (182) 293

Bilirubin conjugates
Separation and characterization of bilirubin conjugates by HPLC (182) 100

Bilirubin conjugates
Accurate and sensitive analysis of ethyl anthranilate azopigments from bile by reversed-phase HPLC (182) 105

Binary liquid phase system
Binary liquid phase system for GLC with *in situ* variable polarity (189) 255

Binding parameters
Determination of ligand-macromolecule binding parameters and the method of Hummel and Dreyer (193) 520

Binding parameters
Usefulness of chromatographic methods for the determination of drug-protein binding parameters (193) 522

Biological samples
Use of microbore columns for the separation of substances of biological origin (185) 27

Biopolymers
High-speed steric exclusion chromatography of biopolymers (185) 453

Biopolymers
Polarity of some chromatographic materials intended for the separation of biopolymers: spectroscopic label technique (196) 39

Biospecific polymers
Use of water-soluble biospecific polymers for the purification of proteins (Review) (184) 325

Bis(chloropiperidyl)piperazinedione
Rapid quantitative assay for 3,6-bis-(5-chloro-2-piperidyl)-2,5-piperazinedione in fermentation broth using HPLC (190) 246

Bis(octadecyl phosphate)zirconium
Bis(octadecyl phosphate)zirconium(IV). Novel support for reversed-phase chromatography (190) 145

Bis(substituted aminoalkylamino)anthraquinones
HPLC of cancer chemotherapeutic agents: bis(substituted aminoalkylamino)anthraquinones (187) 212

Bleomycin
High-performance paired-ion LC determination of bleomycin A_2 in urine (181) 127

Blood cells
Surface affinity chromatographic separation of blood cells. I. Separation of human and rabbit peripheral granulocytes, lymphocytes and erythrocytes using polyethylene glycol-bonded column packings (187) 351

Bonded phase
Simplified HPLC phase characterization: particle size, surface area, mean pore diameter, amount of bonded phase and activity (186) 435

Boratranes
Chromatography of organometallic and organometalloidal derivatives of amino alcohols. II. GC evidence for transannular bonds in solutes (200) 105

Boronic acids
Comparison of electron-capturing boronic acids for the selective analysis of bifunctional compounds (186) 307

Boronic acid-substituted silica
High-performance liquid affinity chromatography of nucleosides, nucleotides and carbohydrates with boronic acid-substituted microparticulate silica (200) 254

Boronic esters
Cyclic derivatives for the selective chromatographic analysis of bifunctional compounds (Review) (184) 99

Bovine serum albumin
Protein-protein interactions studied by counter-current distribution. II. Test of the theoretical model using bovine serum albumin and L-tryptophan (196) 207

Bretylium
Sensitive GC assay for the quantitation of bretylium in plasma, urine and myocardial tissue (181) 33

Bromazepam
GC determination of bromo and fluoro derivatives of benzodiazepine in human body fluids (182) 81

Bromoadenosine monophosphate
Control of 8-bromo-AMP and 8-(6-aminohexyl)amino-AMP synthesis by HPLC (196) 323

Bromoxynil
HPLC analysis of formulations of phenoxycarboxylic acids, dicamba, ioxynil and bromoxynil mixtures (in French) (190) 401

Bromperidol
Separation and quantitation of bromperidol in pharmaceutical preparations by HPLC and HPTLC (193) 255

Bronchodilator drugs
New rapid assay of theophylline in plasma by isotachophoresis (181) 147

Bronchodilator drugs
Use of flow programming in paired-ion HPLC analysis of dosage forms containing dyphylline (193) 111

Buffer effects
Mobile phase effects in reversed-phase chromatography. II. Acidic amine phosphate buffers as eluents (185) 111

Buflophemesyl chloride
tert.-Butylpentafluorophenylmethylchlorosilane as a reagent for the formation of hydrolytically stable alkylsilyl derivatives with electron-capturing properties (187) 331

Bumetanide
Determination of the diuretic bumetanide in biological fluids by HPLC (183) 118

Bupivacaine
Simultaneous determination of bupivacaine and 2,6-pipecoloxylidide in serum by GLC (182) 226

Butoprozine
HPLC in the elucidation of the metabolic fate of butoprozine. Screening for unknown metabolites in the rat (191) 109

Butylhydroquinone
HPLC determination of *tert.*-butylhydroquinone in vegetable oils (187) 436

Butylpentafluorophenylmethylchlorosilane
tert.-Butylpentafluorophenylmethylchlorosilane as a reagent for the formation of hydrolytically stable alkylsilyl derivatives with electron-capturing properties (187) 331

Butyraldehyde
GLC analysis of dimedone derivatives of formaldehyde and other aliphatic aldehydes on capillary columns (191) 239

Butyrophenones
Separation and quantitation of bromperidol in

pharmaceutical preparations by HPLC and HPTLC (193) 255

C

Caffeine
Determination of caffeine in serum by straight-phase HPLC (182) 121

Calcitonin
Paired-ion reversed-phase HPLC of human and rat calcitonin (198) 293

Calcitonin-like proteins
Use of hydrophobic interaction methods in the isolation of proteins from endocrine and paraendocrine tissues and cells by HPLC (185) 413

Calcium complex grease
Calcium complex grease as a low-polarity stationary phase (190) 291

Cannabinoids
Combined HPLC and radioimmunoassay method for the analysis of Δ^9-tetrahydrocannabinol metabolites in human urine (186) 595

Cannabinoids
Simple, rapid method for extraction of urinary cannabinoids by LSC (196) 342

Capacity ratio
Liquid adsorption chromatography with mixed mobile phases. IV. A new equation for the capacity ratio involving the solute-solvent interactions (188) 27

Capillary column preparation
Applications of surface-modified porous silicas to glass capillary column preparation (199) 97

Capillary columns
Preparation of glass capillary columns for GC (Review) (184) 235

Capillary columns
Alumina and support-coated open-tubular columns in open-tubular micro-capillary LC (189) 179

Capillary columns
Preparation of glass capillary columns for LC (191) 25

Capillary columns
Adsorption chromatography capillary columns. I. Rubidium chloride (196) 237

Capillary columns
Open-tubular microcapillary LC with 30-40 μm I.D. columns (199) 249

Capillary liquid chromatography
Techniques of capillary LC (186) 521

Caprolactam
HPLC of caprolactam and its metabolites in urine and plasma (187) 429

Capsaicin
Effective separation of capsaicin and its analogues by reversed-phase HPLC (198) 217

Capsaicinoids
Quantification of nonylic acid vanillylamide and other capsaicinoids in the pungent principle of Capsicum fruits and preparations by GLC on glass capillary columns (189) 389

Capsaicinoids
Rapid determination of nonylic acid vanillylamide and other capsaicinoids in Capsicum fruits and extracts by means of Ag^+-complexation HPLC (in German) (193) 101

Captopril
Determination of captopril in biological fluids by GLC (188) 177

Carbamate insecticides
Resolution, sensitivity and selectivity of a HPLC post-column fluorometric labeling technique for determination of carbamate insecticides (185) 615

Carbamates
Separation of methylol derivatives of imidazolidines, urea and carbamates by LC (190) 226

Carbamates
Distribution ratios and elution behavior of carbamates and polyhydroxy compounds from Sephadex G-15 (190) 448

Carbamates
Determination of carbofuran residues in potato, onion and turnip: comparative analyses using GC-selected ion monitoring or HPLC (193) 213

Carbamates
Simplified method for the analysis of some carbamate insecticides in foliage, forest soil and fish tissue by direct GLC (200) 179

Carbamates
GLC analysis of aminocarb and seven derivatives (200) 228

Carbamazepine
HPLC separation of carbamazepine metabolites excreted in rat urine (181) 59

Carbamazepine
Simultaneous LC analysis for carbamazepine and carbamazepine 10,11-epoxide in plasma and saliva by use of double internal standardization (181) 373

Carbamazepine
Improved GLC method for the simultaneous determination of phenobarbital, phenytoin,

carbamazepine and primidone in biological fluids (182) 116

Carbamazepine
A quantitative method for the simultaneous determination of carbamazepine, mephenytoin, phenylethylmalonamide, phenobarbital, phenytoin and primidone in serum by thin-layer chromatography. Additional comments to the analysis of carbamazepine-10,11-epoxide (183) 387

Carbaryl
Application of a direct aqueous acetylation technique to the GC quantitation of nitrophenols and 1-naphthol in environmental water samples (195) 105

Carbaryl
Simplified method for the analysis of some carbamate insecticides in foliage, forest soil and fish tissue by direct GLC (200) 179

Carbofuran
Determination of carbofuran residues in potato, onion and turnip: comparative analyses using GC-selected ion monitoring or HPLC (193) 213

Carbofuran
Simplified method for the analysis of some carbamate insecticides in foliage, forest soil and fish tissue by direct GLC (200) 179

Carbohydrates
Qualitative TLC separation of 1,5-anhydroglucitol in the presence of other carbohydrates on silica gel impregnated with borate buffer (187) 455

Carbohydrates
Separation of monosaccharides by HPLC: comparison of UV and RI detection (189) 414

Carbohydrates
Studies on lectins. XLV. Effect of ionic strength on the interaction of lectins with carbohydrates as determined by affinity electrophoresis (192) 408

Carbohydrates
Chromatography in aqueous solution on styrene-divinylbenzene resins (193) 197

Carbohydrates
Application of a sintered silica gel plate to the TLC of carbohydrates (193) 464

Carbohydrates
Determination of dissolved carbohydrates in natural water by GLC (194) 224

Carbohydrates
High-performance liquid affinity chromatography of nucleosides, nucleotides and carbohydrates with boronic acid-substituted microparticulate silica (200) 254

Carbon adsorbent
A PTFE-based carbon adsorbent in HPLC (191) 61

Carbon dioxide
Oxygen enhancement of the electron capture detector response of carbon dioxide (188) 89

Carbon dioxide
Response of electron-capture detector to hydrogen, oxygen, nitrogen, carbon dioxide, nitric oxide and nitrous oxide (193) 142

Carbon skeleton capillary gas chromatography
Carbon skeleton capillary GC (193) 437

Carbon skeleton chromatography
Carbon skeleton-GC techniques and their applications (187) 47

Carbon tetrachloride
Two rapid methods for the simultaneous GLC determination of carbon tetrachloride and chloroform in biological material and expired air (193) 71

Carbonyl compounds
GC determination of low-molecular-weight carbonyl compounds in aqueous solution as their O-(2,3,4,5,6-pentafluorobenzyl) oximes (187) 413

Carbonyl compounds
syn-anti Isomerisation of 2,4-dinitrophenyl-hydrazones of volatile carbonyl compounds in capillary GC-MS analyses (194) 135

Carbonyl groups, conjugated
Quantitation of non-halogenated compounds by electron-capture GC (Review) (184) 313

Carbophenothion
Improved clean-up procedure for the determination of small residues of carbophenothion in mice (190) 201

Carbowax 20M
Use of Carbowax 20M glass capillary columns below their lower temperature limit (190) 283

Carboxylic acids
Determination of free short-chain monocarboxylic acids from naturally occurring esters by GLSC on micropacked columns (in German) (189) 43

Carboxylic acids
Determination of multifunctional carboxylic acids in the presence of iron by GC (190) 188

Carboxylic acids
GC separation of chiral 2-hydroxy acids and 2-alkyl-substituted carboxylic acids (195) 292

Carboxylic acids
Alkylation of carboxylic acids by solid-liquid phase-transfer catalysis for determination by GC. II. (196) 255

Carboxylic acid surfactants
Analysis of dipeptide mixtures by the combination of ion-pair reversed-phase HPLC and GC-MS techniques (197) 31

Carboxylic cation exchangers
Ion-exchange derivatives of Spheron. III. Carboxylic cation exchangers (192) 159

Carcinogens
Combined reversed-phase and normal-phase HPLC in the purification and identification of 7,12-dimethylbenz[*a*]anthracene metabolites (185) 635

Carcinogens
Improved GC method for the quantitative determination of secondary amines as sulphonamides formed by reaction with benzenesulphonyl chloride (190) 462

Carcinogens
Analysis of N-nitroso compounds. V. Trace analysis of N-nitrosoamino acids: Gas chromatography and spectroscopy of N-nitrosoamino acid methyl esters (in German) (193) 381

Carcinogens
HPLC analysis of the mycotoxin penicillic acid and its application to biological fluids (194) 387

Carcinogens
Separation and detection of 2-aminoanthracene and its metabolites by HPLC (195) 392

Carcinogens
Quantitative GC determination of tris(2,3-dibromopropyl)phosphate in the 10-ng range by using a 0.8-mm I.D. column packed with a high liquid loaded support (196) 133

Carcinogens
The isolation of methionine and ethionine by silver ligand chromatography and application to methionine-containing peptides (200) 185

Cardiac drugs
An integrated approach to measurements of quinidine and metabolites in biological fluids (183) 514

Cardiotonic agents
HPLC analysis of amrinone and its N-acetyl derivative in plasma. Pharmacokinetics of amrinone in the dog (187) 264

Carminomycin
HPLC and TLC of anthracycline antibiotics. Separation and identification of components of the daunorubicin complex from fermentation broth (198) 407

Carpipramine
Determination of carpipramine in plasma by HPLC (183) 246

Carrier ampholytes
New developments in isoelectric focusing (Review) (184) 415

Carrier ampholytes
Molarity and ionic strength of focused carrier ampholytes in isoelectric focusing (190) 275

Carrier ampholytes
Study of the feasibility of separation of carrier ampholytes from peptides by hydrophobic interaction chromatography on octyl-Sepharose (196) 51

Carrier ampholytes
Gel chromatographic comparison of the molecular weight distributions of Ampholine, Servalyte and Pharmalyte carrier ampholytes used in isoelectric focusing (198) 301

Carrier gas selection
Influence of different carrier gases and temperatures on the overall performance of a packed column (186) 249

Catalysis reactions
Study of the kinetics for a catalyzed reaction using GC (197) 129

Catalysts
Thermoprogrammed reduction of cobalt oxide catalysts (188) 254

Catalysts
GC determination of adsorption-desorption rates. Interaction of hydrogen on a nickel catalyst (196) 355

Catecholamine metabolites
Diagnosis of neural crest tumors by reversed-phase HPLC determination of urinary catcholamine metabolites (186) 733

Catecholamines
Simplified LC-electrochemical determination of norepinephrine and dopamine in rat brain (181) 453

Catecholamines
Simple and sensitive procedure for the assay of serotonin and catecholamines in brain by HPLC using fluorescence detection (182) 277

Catecholamines
Simple method for isolating free urinary catecholamines by boric acid gel chromatography (183) 78

Catecholamines
Investigation of the performance of a HPLC system with an electrochemical detector (183) 261

Catecholamines
Semi-automated fluorimetric method for the estimation of urinary catecholamines using HPLC (183) 295

Catecholamines
Comparison of LC-electrochemical and GC-MS methods for brain dopamine and serotonin (183) 483

Catecholamines
Determination of apomorphine in plasma and brain tissue by ion-pair extraction and LC (185) 575

Catecholamines
GC of epinephrine and norepinephrine after

derivatization with chloroformates in aqueous media (190) 347

Catecholamines
Electrochemical detector for HPLC (192) 135

Catecholamines
GC of phenolic amines, 3-methoxycatecholamines, indoleamines and related amines as their N,O-ethyloxycarbonyl derivatives (194) 399

Catecholamines
Determination of catecholamines and O-methylated metabolites by reversed-phase HPLC with fluorimetric detection and its application to enzyme kinetics (196) 101

Cation exchangers
Strong non-polar cation exchangers for the separation of steroids in mixed chromatographic systems (186) 725

Caustics
Isolation of α- and β-peltatin and podophyllotoxin by LC and analysis by HPLC (189) 276

Cefoxitin
Reversed-phase HPLC assay for cefoxitin in proteinaceous biological samples (181) 120

Cefoxitin
HPLC assay for measurement of cefoxitin in serum (183) 357

Ceftizoxime
HPLC determination of ceftizoxime, a new cephalosporin antibiotic, in rat serum, bile and urine (182) 448

Cell membranes
Affinity chromatography of cells and cell membranes (Review) (184) 471

Cell pressure
Effects of dimension and density in electron capture detectors (195) 17

Cells
New developments in isoelectric focusing (Review) (184) 415

Cells
Affinity chromatography of cells and cell membranes (Review) (184) 471

Cells
New experimental approaches to the isoelectric fractionation of cells (194) 323

Cellulase complex
Affinity chromatography of a cellulase complex from *Trichoderma viride* on cross-linked cellulose (in French) (188) 377

Cellulose
New cellulose gel for chromatography (195) 221

Cellulose acetate isoelectric focusing
New developments in isoelectric focusing (Review) (184) 415

Cellulosics
High-performance gel permeation chromatography of water-soluble cellulosics (192) 275

Cephalosporin
Fast and simple method for the separation of intermediates and cofactors involved in the biosynthesis of cephalosporin C using chemically bonded C_{12} reversed-phase TLC (193) 511

Cephalosporin derivatives
HPLC determination of ceftizoxime, a new cephalosporin antibiotic, in rat serum, bile and urine (182) 448

Chain length
Role of the chain length of chemically bonded phases and the retention mechanism in reversed-phase LC (196) 21

Charge distribution
Size and charge distribution of macromolecules in living systems (193) 1

Charge numbers
Role of the charge number of the counter-ionic constituent in the separation of anions by isotachophoresis (194) 11

Charge-transfer interactions
Electron donor-acceptor properties of urea and its role in charge-transfer chromatography (188) 257

Chemiluminescence detection
Chemiluminescence HPLC detector applied to ascorbic acid determination (200) 153

Chemotherapeutic agents
HPLC of cancer chemotherapeutic agents: bis(substituted aminoalkylamino)anthraquinones (187) 212

Chemotherapeutic agents
Estimation of pipemidic acid and N-nitrosopipemidic acid by HPLC (200) 245

Chemotherapeutic drugs
Determination of methenamine in biological samples by GLC (181) 11

Chemotherapeutic drugs
Chromatographic analysis of griseofulvin and metabolites in biological fluids (181) 77

Chemotherapeutic drugs
Non-extractive fluorometric measurement of *p*-aminosalicylic acid in plasma by ion-pairing techniques and HPLC (181) 266

Chemotherapeutic drugs
Determination of nalidixic acid and its two major metabolites in human plasma and urine by reversed-phase HPLC (181) 399

Chemotherapeutic drugs
Rapid HPLC method for the determination of dapsone and monoacetyldapsone in biological fluids (182) 130

Chemotherapeutic drugs
HPLC assay for the metabolites of nitrofurantoin in plasma and urine (182) 246

Chemotherapeutic drugs
HPLC of proguanil, cycloguanil and 4-chlorophenylbiguanide using hydrophobic pairing ion and its application to serum assay (182) 359

Chemotherapeutic drugs
HPLC determination of chloramphenicol and its monosuccinate ester in plasma (182) 459

Chemotherapeutic drugs
GC determination of amantadine hydrochloride (Symmetrel) in human plasma and urine (183) 33

Chemotherapeutic drugs
Two HPLC determinations for mebendazole and its metabolites in human plasma using a rapid Sep Pak C18 extraction (183) 311

Chemotherapeutic drugs
Analysis of isoxazolyl penicillins and their metabolites in body fluids by HPLC (183) 339

Chemotherapeutic drugs
HPLC assay for measurement of cefoxitin in serum (183) 357

Chemotherapeutic drugs
Investigation of vinca alkaloid-plasma membrane interactions by detergent gel chromatography (191) 287

China, developments in chromatography
The Third National Symposium on Chromatography in China (Review) (184) 215

Chiral additive
Reversed-phase HPLC resolution of D- and L-Dns-amino acids by mixed chelate complexation (199) 295

Chiral additives
Reversed-phase separation of optical isomers of Dns-amino acids and peptides using chiral metal chelate additives (185) 323

Chiral bonded stationary phases
Ligand-exchange chromatography of enantiomeric amino acids on copper-loaded chiral bonded silica gel and of amino acids on copper(II)-modified silica gel (185) 345

Chiral hydroxy acids
Structural requirements for enantioselectivity in GC of chiral α-hydroxy acids (200) 189

Chiral stationary phases
LC resolution of enantiomers on normal-phase chiral amide-bonded silica gel. Retentions of optically active α-amino acid derivatives on N-acyl homologues of L-valylaminopropylsilanized silica phases (186) 543

Chiral stationary phases
Synthesis and properties of a novel chiral stationary phase for the resolution of amino acid enantiomers (186) 611

Chiral stationary phases
Broad spectrum resolution of optical isomers using chiral HPLC bonded phases (192) 143

Chitin
New method of chitin determination based on deacetylation and GLC assay of liberated acetic acid (190) 67

Chloramine-T reagent
Alkaline chloramine-T reagent for the detection of phenolic compounds on thin-layer plates (196) 309

Chloramphenicol
HPLC micro-assay for chloramphenicol in human blood plasma and cerebrospinal fluid (181) 123

Chloramphenicol
HPLC determination of chloramphenicol and its monosuccinate ester in plasma (182) 459

Chlordiazepoxide
Determination of chlordiazepoxide and its metabolites in human plasma by reversed-phase HPLC (181) 141

Chlorinated aromatic compounds
General approach to the fractionation and class determination of complex mixtures of chlorinated aromatic compounds (197) 155

Chlorinated aromatic compounds
HPLC in the analysis of chorinated dibenzodioxins and dibenzofurans in chicken liver and wood shaving samples (197) 171

Chlorinated benzenes
Retention behavior of chlorinated benzenes, chlorinated phenols, Aldrin, and Aroclor 1242 on Tenax-GC, Chromosorb 101, Florisil, and Carbopack C HT (200) 115

Chlorinated dioxins
Rapid analysis of chlorinated dioxins in complex organic mixtures by application of a GC-MS-calculator system with user-developed software (189) 53

Chlorinated hydrocarbons
Evaluation of macroreticular resins for the determination of low concentrations of chlorinated hydrocarbons in sea water and tap water (193) 357

Chlorinated phenols
Retention behavior of chlorinated benzenes, chlorinated phenols, Aldrin, and Aroclor 1242 on Tenax-GC, Chromosorb 101, Florisil, and Carbopack C HT (200) 115

Chlorinated quinones
GC identification and determination of chlorinated quinones formed during chlorination of dihydric phenols with hypochlorite in dilute aqueous solution (200) 137

Chlormethiazole
Simultaneous quantitation of chlormethiazole and two of its metabolites in blood and plasma by GLC (182) 55

Chlormethylisatin
Derivatization of fatty acids with 1-chlormethylisatin for HPLC (187) 208

Chlorocresol
TLC of 2-methyl-4-chlorophenoxyacetic acid and its soil metabolites (189) 73

Chloroform
Two rapid methods for the simultaneous GLC determination of carbon tetrachloride and chloroform in biological material and expired air (193) 71

Chlorohydrins
GC-MS of the flophemesyl derivatives of a series of fatty acid methyl ester chlorohydrins (189) 86

Chloromethylnitrophthalimide
N-Chloromethyl-4-nitrophthalimide as derivatization reagent in HPLC (in German) (198) 367

Chlorophenylbiguanide
HPLC of proguanil, cycloguanil and 4-chlorophenylbiguanide using hydrophobic pairing ion and its application to serum assay (182) 359

Chlororesorcinol
Analysis of non-volatile organic compounds in water by HPLC (185) 671

Chlorosalicylic acid
Analysis of non-volatile organic compounds in water by HPLC (185) 671

Chlorouracil
Analysis of non-volatile organic compounds in water by HPLC (185) 671

Chlorouridine
Analysis of non-volatile organic compounds in water by HPLC (185) 671

Chlorpyrifos
Use of chlorpyrifos as an evaluation standard for GC detectors (188) 305

Chlorthalidone
Rapid and sensitive determination of chlorthalidone in blood, plasma and urine of man using HPLC (181) 497

Cholecalciferol
GC-MS properties of boronate esters of 24*R*,25-dihydroxycholecalciferol (192) 434

Cholesterol
Lipid content of human platelets quantitated by TLC in combination with flame ionization detection (182) 301

Cholesterol
Fractionation of steroid digitonides by TLC and CC on silica gel (188) 445

Cholesterol
Pregnenolone separation from cholesterol using Sephadex LH-20 minicolumns (190) 256

Cholesterol
Capillary GC studies of cholesterol biosynthesis in rats treated with EGYT-1299 (191) 261

Cholic acid
Covalent coupling of cholic acid to aminohexylamino-Sepharose 4B and its use in affinity chromatography of serum albumin (187) 409

Choline
Paired-ion HPLC assay for plasma choline (181) 329

Cholinergic drugs
Reversed-phase, ion-pair LC of quaternary ammonium compounds. Determination of pyridostigmine, neostigmine and edrophonium in biological fluids (183) 193

Cholinergic drugs
Determination of pilocarpine, physostigmine, its degradation product rubreserine and preservatives by HPLC (198) 529

Chromatin proteins
Chromatography of chromatin proteins on Cibacron Blue F3G-A-agarose (192) 494

Chromatin proteins
Quantitative analysis of non-histone chromosomal proteins HMG 14 and HMG 17 by polyacrylamide gel electrophoresis (198) 80

Chromatography, developments in China
The Third National Symposium on Chromatography in China (Review) (184) 215

Chromia layers
GC and LC on silica-supported chromia layers (197) 21

Chromium
GC of volatile metal dialkyldithiophosphates (193) 53

Chromium chelates
Minimizing the retention of chromium chelates with some fluorinated β-diketones (191) 231

Chromium complexes
Paper electrophoretic study of ion-pair formation. XIII. Behaviour of anionic trisoxalatocobalt(III) and trisoxalatochromium(III) in optically active electrolytes (198) 76

Chromium complexes
Resolution of neutral complexes of transition metals by stereoselective adsorption on optically active complexes. I. Partial resolution of tris(acetylacetonato)cobalt(III) and tris(acetylacetonato)chromium(III) on cobalt(III) complexes (198) 172

Chymotrypsin
Affinity chromatographic interactions of proteases with low-molecular-weight soybean protease inhibitors (195) 385

Ciba-Geigy
The role of chromatography in Ciba-Geigy (Review) (184) 207

Cigarette smoke
Study of nitrogen-containing compounds in cigarette smoke by GC-MS (198) 188

Cinchona alkaloids
TLC separation of *Cinchona* alkaloids (Review) (184) 79

Cinchona alkaloids
Reversed-phase HPLC analysis of cinchona alkaloids in pharmaceuticals (189) 241

Cinchona alkaloids
Analysis of cinchona alkaloids by HPLC (in French) (192) 479

Cinerin
Examination of the conversion products of pyrethrins and allethrin formulations exposed to sunlight by GC and MS (198) 317

Cinnarizine
Determination of cinnarizine in plasma by HPLC (183) 378

Citric acid
Determination of multifunctional carboxylic acids in the presence of iron by GC (190) 188

Citric acid cycle intermediates
TLC of citric acid cycle intermediates, lactic acid, and the amino acid taurine (189) 286

Citrinin
HPLC analysis of the mycotoxin citrinin and its application to biological fluids (190) 419

Clobazam
Use of PTFE coils in post-column photochemical reactors for LC: application to pharmaceuticals (199) 239

Clonidine
Determination of clonidine in human plasma by glass capillary GC with electron-capture detection (187) 161

Clopenthixol
Specific HPLC method for estimation of the *cis*(*Z*)- and *trans*(*E*)-isomers of clopenthixol and a N-dealkyl metabolite (183) 239

Clorazepate
LC identification of clorazepate in pharmaceutical products (188) 426

Clovoxamine
The use of a pre-column for the direct HPLC determination of the antidepressants clovoxamine and fluvoxamine in plasma (183) 203

Coal distillate
GC study of methylstyrenes in the light-oil fraction of coal distillate using selective hydrogenation (193) 397

Coating technique
Coating of glass capillary chromatographic columns: a new variant of the static method (190) 445

Coating techniques
Synthesis and GC use of metal-containing surfaces (189) 13

Coating thickness
GC retention characteristics of different coating thicknesses on a glass capillary column (194) 277

Cobalamins
TLC of acid-labile cobalamins (194) 273

Cobalt complexes
Effect of the composition and structure of cobalt(III) complexes on their R_F values obtained by TLC on silica gel (196) 59

Cobalt complexes
Paper electrophoretic study of ion-pair formation. XIII. Behaviour of anionic trisoxalatocobalt(III) and trisoxalatochromium(III) in optically active electrolytes (198) 76

Cobalt complexes
Resolution of neutral complexes of transition metals by stereoselective adsorption on optically active complexes. I. Partial resolution of tris(acetylacetonato)cobalt(III) and tris(acetylacetonato)chromium(III) on cobalt(III) complexes (198) 172

Cobalt oxide catalysts
Thermoprogrammed reduction of cobalt oxide catalysts (188) 254

Cocaine
Qualitative screening procedure for the detection of benzoyl ecgonine in the urine of greyhounds (196) 337

Cocaine
Qualitative differentiation between cocaine, lidocaine and cocaine-lidocaine mixtures ("rock-cocaine") using TLC (198) 534

Cocaines
Positive identification and quantitation of isomeric cocaines by HPLC (193) 371

Coil planet centrifuge
New horizontal flow-through coil planet centrifuge for counter-current chromatography. I. Principle of design and analysis of acceleration (188) 33

Coil planet centrifuge
New horizontal flow-through coil planet centrifuge for counter-current chromatography. II. The apparatus and its partition capabilities (188) 43

Coil planet centrifuge
Toroidal coil planet centrifuge for counter-current chromatography (192) 75

Coil planet centrifuge
New horizontal flow-through coil planet centrifuge for counter-current chromatography. III. Separation and purification of dinitrophenyl amino acids and peptides (193) 37

Colchicine
HPLC separation of colchicine and its phenolic and N-desacetylated derivatives (188) 280

Collagen
Employment of GLC for the analysis of collagen amino acids in biopsy tissue (182) 155

Collagen
Separation of cyanogen bromide peptides of collagen by means of HPLC (190) 393

Collagens
Chromatographic separation and quantification of type I and type III collagens (182) 94

Colloids
Size-exclusion HPLC of inorganic colloids (185) 273

Column coating
Preparation of glass capillary columns for GC (Review) (184) 235

Column coating
Glass capillary columns coated with non-soluble methyl silicone gums (186) 159

Column coating
Alumina and support-coated open-tubular columns in open-tubular micro-capillary LC (189) 179

Column coating
Marine sterols. XIII. Evaluation of glass support-coated open tubular columns, prepared by the mercury-plug dynamic technique, for marine sterol analysis (198) 111

Column coating
Coating support-coated open tubular capillary columns with Silar 10C or Alltech CS-10 and Silica T40 for separation of isomers of fatty acid methyl esters (198) 153

Column coatings
Comparison of capillary columns coated with C_{87} hydrocarbon and squalane in the analysis of *n*-pentadecene isomers (191) 199

Column deactivation
Polydentate phosphonium salts for the deactivation of wall-coated open-tubular columns. Comparison of temperature stability with other deactivation methods (187) 57

Column deactivation
Simple device to facilitate the deactivation of a GC system by PEG 20M via the gas phase (192) 418

Column dead time
Evaluation of procedures for the estimation of dead time (Review) (184) 1

Column efficiency
Preparation and operation of LC columns of very high efficiency (185) 3

Column efficiency
High-performance GC columns packed with micro-particles (186) 25

Column efficiency
Ultra-micro HPLC (190) 150

Column loading
In situ loading and reloading of GC columns with stationary liquids (186) 285

Column packings
Retention characteristics of several bonded-phase LC columns for some polycyclic aromatic hydrocarbons (188) 115

Column performance
Influence of different carrier gases and temperatures on the overall performance of a packed column (186) 249

Column performance
Enhancement of GSC column performance by inert solid dilution (188) 79

Column preparation
Preparation of glass capillary columns for GC (Review) (184) 235

Column preparation
Preparation and operation of LC columns of very high efficiency (185) 3

Column preparation
Preparation of glass capillary columns for LC (191) 25

Column preparation
Adsorption chromatography capillary columns. I. Rubidium chloride (196) 237

Column preparation
Design and use of short microbore columns in LC (198) 93

Column preparation
Glass surface analytical studies in the preparation of open tubular columns for GC (199) 355

Columns
Contamination by use of Extrelut columns (in German) (183) 1

Columns
Use of microbore columns for the separation of substances of biological origin (185) 27

Columns
Use of microbore columns for rapid LC separations (186) 475

Columns
Glass columns with high resistance to internal pressure for HPLC (193) 128

Columns
HPLC columns of small diameter (197) 97

Columns, capillary adsorption
Glass capillary adsorption columns for GC (186) 183

Column switching
HPLC column switching technique in the analysis of medicated feed for an automated clean-up procedure (198) 162

Column treatment
Studies of open-tubular micro-capillary LC. IV. Soda-lime glass columns treated with alkaline solution (185) 73

Complex structure
Effect of the composition and structure of cobalt(III) complexes on their R_F values obtained by TLC on silica gel (196) 59

Compound chromatography
Compound chromatography (191) 95

Computerised chromatography
Software improvements for GC-MS calculator system used in the analysis of trace organic compounds from environmental samples (190) 311

Concanavalin A
Non-bonded affinity chromatography: a biospecific oil-soluble coated support for concanavalin A (188) 61

Concentration effects
Concentration effects in gel permeation chromatography. VI. Non-Newtonian behaviour of polymer solutions (187) 21

Conductimetric detector
Conductimetric detection of organic acids in reversed-phase chromatography (185) 625

Conductivity detection
High-frequency contactless conductivity detection in isotachophoresis (192) 253

Congeners
GC column for the rapid determination of congeners in potable spirits (198) 347

Congeners
HPLC separation of urushiol congeners in poison ivy and poison oak (200) 163

Contact spotting
Contact spotting. A new approach to sample application in TLC (186) 555

Continuous flow electrophoresis
Continuous flow electrophoresis. The crescent phenomena revisited. I. Isothermal effects (195) 165

Contraceptives
HPLC and GLC of some norgestrel intermediates. Physical properties of the isolated *syn*- and *anti*-isomers of oximes (191) 145

Contractile proteins
HPLC of myosin's subfragment 1 and light chains on Spherogel TSK-type columns (197) 207

Contrast media
Rapid determination of renal contrast media in biological fluids by means of HPLC (182) 492

Convicine
Chromogenic reagent for vicine and convicine on thin-layer plates (194) 251

Copolymers
Determination of the composition of copolymers as a function of molecular weight by pyrolysis GC size-exclusion chromatography (194) 163

Copper
Quantitative analysis of zinc, copper and nickel diethyldithiocarbamates by GLC (187) 363

Copper
Binding of zinc and copper to some gel filtration media (188) 405

Copper-modified silica gel
Ligand-exchange chromatography of enantiomeric amino acids on copper-loaded chiral bonded silica gel and of amino acids on copper(II)-modified silica gel (185) 345

Corn syrups
A rapid and high-resolution method to determine the composition of corn syrups by LC (187) 381

Corticosteroids
Quantitative analysis of corticosteroids in adrenal cell cultures by capillary column GC combined with MS (186) 581

Corticosteroids
Reversed- and normal-phase HPLC of 18-hydroxylated steroids and their derivatives. Comparison of selectivity, efficiency and recovery from biological samples (198) 23

Corticosteroids
Analysis of hydrocortisone acetate ointments and creams by HPLC (198) 41

Corticosteroids
PC systems for the separation of aldosterone, 18-hydroxycorticosterone, 18-hydroxydeoxycorticosterone, corticosterone and deoxycorticosterone (198) 70

Corticosterone
PC systems for the separation of aldosterone, 18-hydroxycorticosterone, 18-hydroxydeoxycorticosterone, corticosterone and deoxycorticosterone (198) 70

Cortisol
Rapid quantitative assay of plasma 11-

deoxycortisol and cortisol by HPLC for use in the metyrapone test (183) 87

Coumarin
HPLC of coumarin and its metabolites (196) 501

Coumarin derivatives
Simple rapid TLC method for the detection of optical brighteners in polymers (194) 434

Counter-ions
Role of the charge number of the counter-ionic constituent in the separation of anions by isotachophoresis (194) 11

Coupling, on-line
On-line coupling of HPLC and NMR (186) 497

Crescent phenomena
Continuous flow electrophoresis. The crescent phenomena revisited. I. Isothermal effects (195) 165

Cross-correlation gas chromatography
Comparison between conventional and cross-correlation GC for evolved gas analysis (186) 145

Cross-linkage effect
Peptide separations on substituted polystyrene resins. Effect of cross-linkage (187) 27

Crude oil fractions
Preparative separation of C_9 and C_{10} monoaromatic hydrocarbon mixtures by HPLC on silica gel (191) 319

Cyanide
Determination of cyanide and thiocyanate in biological fluids by GC-MS (188) 357

Cyanoazoxybenzene alkyl carbonates
Properties of liquid crystalline cyanoazoxybenzene alkyl carbonates as stationary phases in GC (200) 65

Cyanogen bromide peptides
Separation of cyanogen bromide peptides of collagen by means of HPLC (190) 393

Cyclic alcohols
HPLC of adamantanols and other cyclic alcohols (198) 457

Cyclic amides
HPLC of caprolactam and its metabolites in urine and plasma (187) 429

Cyclic derivatives
Cyclic derivatives for the selective chromatographic analysis of bifunctional compounds (Review) (184) 99

Cycloalkenylnaphthalenes
GC of some alkenyl- and cycloalkenylnaphthalenes on Bentone 34, silicone fluid DC 550, and Apiezon L phases (198) 1

Cyclobutadipyrimidines
Separation of cyclobutyl dimers of thymine and thymidine by HPLC and TLC (195) 139

Cyclodextrin polyurethane resins
Preparation and sorption behaviour of cyclodextrin polyurethane resins (194) 153

Cyclodextrins
Inclusion compounds in chromatography (Review) (184) 347

Cyclodextrins
Analysis of cyclodextrins by HPLC (189) 98

Cycloguanil
HPLC of proguanil, cycloguanil and 4-chlorophenylbiguanide using hydrophobic pairing ion and its application to serum assay (182) 359

Cyclophosphamide
Some aspects of the GLC analysis of cyclophosphamide in plasma (183) 49

Cyclosporin
HPLC determination of cyclosporin A in human plasma and urine (182) 454

Cyclosteroids
Reversed-phase HPLC separation and analysis of some physalins (13,14-*seco*-16,24-cyclosteroids) (198) 203

Cysteic acid
Analysis of cysteic acid and cysteine sulfinic acid by GLC (187) 239

Cysteine
Estimation of glutathione in rat liver by reversed-phase HPLC: separation from cysteine and γ-glutamylcysteine (194) 424

Cysteine derivatives
Isotachophoretic analysis of some sulfur-containing amino acids in human urine (183) 226

Cysteine sulphinic acid
Analysis of cysteic acid and cysteine sulfinic acid by GLC (187) 239

Cytidylic acid
Application of a weakly basic dimethylamino-modified silica ion exchanger to the separation of oligonucleotides (185) 403

Cytokinins
High-performance steric exclusion chromatography of plant hormones (198) 57

Cytosine
Analysis of pyrimidine bases in biological materials by GC-MS (190) 107

Cytosine arabinoside
HPLC analysis of cytosine arabinoside and metabolites in biological samples (183) 449

Cytostatic drugs, see also antitumour drugs and antineoplastic drugs

Cytostatic drugs
High-performance paired-ion LC determination of bleomycin A_2 in urine (181) 127

Cytostatic drugs
Analysis of methotrexate by isotachophoresis (181) 441

Cytostatic drugs
Reversed-phase HPLC analysis of methotrexate and 7-hydroxymethotrexate in serum (181) 478

Cytostatic drugs
Determination of estramustine and its 17-keto metabolite in plasma by HPLC (182) 387

D

Dalapon
Improved determination of dalapon residues in plant tissues and natural water by derivatization and electron capture GC (187) 244

Dapsone
Rapid HPLC method for the determination of dapsone and monoacetyldapsone in biological fluids (182) 130

Data analysis
Numerical analysis of chromatographic-spectrometric data. Purity of partially resolved peaks and pattern separation of the pure component spectra (186) 443

Daunorubicin
Determination of adriamycin and daunorubicin in urine by HPLC with laser fluorometric detection (190) 377

Daunorubicin
HPLC and TLC of anthracycline antibiotics. Separation and identification of components of the daunorubicin complex from fermentation broth (198) 407

Deactivation
Simple device to facilitate the deactivation of a GC system by PEG 20M via the gas phase (192) 418

Dead time estimation
Evaluation of procedures for the estimation of dead time (Review) (184) 1

Decalines
Identification of the pyrolysis products of *cis*-decaline (in French) (192) 216

Dehydrogenase
Study of the interaction between NADP-dependent dehydrogenase and immobilized adenosine 2'-monophosphate by means of affinity electrophoresis (196) 85

Dehydroglaucine
HPLC determination of glaucine and its metabolite, dehydroglaucine in microbial culture (193) 170

Dehydroretinals
HPLC analysis of *cis-trans* stereoisomeric 3-dehydroretinals in the presence of retinal isomers (192) 395

Denatonium benzoate
HPLC determination of denatonium benzoate in rapeseed oil (195) 243

Denaturants
HPLC determination of denatonium benzoate in rapeseed oil (195) 243

Deoxycholate
Use of deoxycholate in gel filtration chromatography of synaptosomal membrane glycoproteins (193) 316

Deoxycorticosterone
PC systems for the separation of aldosterone, 18-hydroxycorticosterone, 18-hydroxydeoxycorticosterone, corticosterone and deoxycorticosterone (198) 70

Deoxycortisol
Rapid quantitative assay of plasma 11-deoxycortisol and cortisol by HPLC for use in the metyrapone test (183) 87

Deoxyriboadenylic acids
Preparative isolation of deoxyriboadenylic acids from hydrolysates of oxidized herringsperm DNA using template-chromatography (in German) (187) 119

Deoxyriboguanylic acids
Preparation of oligomers of deoxyriboguanylic acids by partial chemical hydrolysis of DNA (in German) (196) 435

Deoxyribonuclease
Specificity of deoxyribonuclease hydrolysis determined by anion-exchange HPLC (193) 175

Deoxyribonucleic acids, see DNA

Deoxyribonucleotides
Determination of pyrimidine ribotide and deoxyribotide pools in cultured cells and mouse liver by HPLC (188) 149

Deoxyribonucleotides
Analysis of purine ribonucleotides and deoxyribonucleotides in cell extracts by HPLC (198) 435

Deoxyuridines
GC determination of 5-alkyl-uracils and 5-alkyl-deoxyuridines using flash methylation and glass capillary columns (191) 253

Derivative structure
Effect of derivative structure on flame-ionization detector response of amino acid oxazolidinones (197) 226

Derivatization
Cyclic derivatives for the selective chromatographic analysis of bifunctional compounds (Review) (184) 99

Derivatization
Estimation of neutral sugars and sugar alcohols in biological fluids by GLC (Review) (184) 457

Derivatization
Derivatization of fatty acids with 1-chlormethylisatin for HPLC (187) 208

Derivatization
Comparative study of derivatisation procedures for the quantitative determination of the auxin, phenylacetic acid, by GC (192) 429

Derivatization
Dimethoxymethylsilyl ethers: stable silyl derivatives for GC and MS (196) 156

Derivatization
GC of trimethylsilylesters of trifluoroacetylamino acids. III. Simple method for the preparation of the trimethylsilylesters of N-trifluoroacetyl amino acids (in German) (196) 160

Derivatization reagents
tert.-Butylpentafluorophenylmethyl-chlorosilane as a reagent for the formation of hydrolytically stable alkylsilyl derivatives with electron-capturing properties (187) 331

Derivatization reagents
GC determination of low-molecular-weight carbonyl compounds in aqueous solution as their O-(2,3,4,5,6-pentafluorobenzyl) oximes (187) 413

Derivatization reagents
N-Chloromethyl-4-nitrophthalimide as derivatization reagent in HPLC (in German) (198) 367

Derivatization reagents
New electron-capturing pentafluorophenyldialkylchlorosilanes as versatile derivatizing reagents for GC (199) 123

Derivatization reagents
Derivatization of dithiols and certain monothiols with phenylarsine oxide for GC (200) 171

Desipramine
Technical considerations in the GC analysis of desipramine (183) 149

Desmosterol
Capillary GC studies of cholesterol biosynthesis in rats treated with EGYT-1299 (191) 261

Desorption
GC technique for temperature-programmed desorption of reversibly and irreversibly adsorbed species (192) 420

Desoxycholic acid
Inclusion compounds in chromatography (Review) (184) 347

Detection
Solid-phase reagents for LC detection (185) 497

Detection system
High-frequency contactless conductivity detection in isotachophoresis (192) 253

Detector performance
Use of chlorpyrifos as an evaluation standard for GC detectors (188) 305

Detector response
Non-linear responses of the electron-capture detector to alkyl monochlorides (192) 117

Detector response
Alteration of the electron-capture detector response to polycyclic aromatic hydrocarbons by oxygen doping of the carrier gas (197) 51

Detector response
Effect of derivative structure on flame-ionization detector response of amino acid oxazolidinones (197) 226

Detector response
Application of a unified theory of GC photoionization detector response to a 10.2-eV detector (200) 35

Detectors
Small-volume electrochemical detector for microcolumn LC (181) 287

Detectors
Capacilities of on-line element-specific detection in HPLC using an inductively coupled argon plasma emission source detector (185) 549

Detectors
Application of a graphite furnace atomic absorption detector automatically coupled to a high-performance liquid chromatograph for speciation of metal-containing macromolecules (185) 563

Detectors
Conductimetric detection of organic acids in reversed-phase chromatography (185) 625

Detectors
Analysis of kinetics and concentration dependence of electron-capture detector (186) 99

Detectors
Design and selective application of a dropping mercury electrode amperometric detector in column LC (186) 489

Detectors
Response of nitramines in the thermal analyzer (187) 229

Detectors
Studies and speculations on an alternative

mechanism for the response of direct-current electron-capture detectors (188) 1

Detectors
Oxygen enhancement of the electron capture detector response of carbon dioxide (188) 89

Detectors
New results with a flow-impedance bridge detector (189) 3

Detectors
Performance of an evaporative analyser detector for gel permeation chromatography (189) 259

Detectors
Analysis errors following hydrogen cleaning of an electron capture detector (190) 133

Detectors
The photoionization detector. Theory, performance and application as a low-level monitor of oil vapour (190) 263

Detectors
Application of an electrochemical detector to the determination of procarbazine hydrochloride by HPLC (190) 359

Detectors
Graphite furnace atomic absorption spectrophotometers as automated element-specific detectors for HPLC. The determination of arsenite, arsenate, methylarsonic acid and dimethylarsinic acid (191) 31

Detectors
Polarographic detection for HPLC using a flow-through detector (191) 47

Detectors
Electrochemical detector for HPLC (192) 135

Detectors
Comparative investigation of radioactive sources for the electron capture detector (195) 1

Detectors
Effects of dimension and density in electron capture detectors (195) 17

Detectors
Survey of selected hydrides as doping agents for a hydrogen-atmosphere flame-ionization detector (195) 211

Detectors
Use of the image analyser Optomax for the quantitative evaluation of antibiotics separated by gel electrophoresis and by TLC (195) 297

Detectors
Synchronized accumulating radioisotope detector for radio GC (196) 144

Detectors
Reaction and response in electron-capture detectors (196) 397

Detectors
Effect of the sensitivity setting of a katharometer on response factors at low concentrations (197) 121

Detectors
Specific detection in GLC effluents with on-line HPLC optical detectors (199) 113

Detectors
Use of PTFE coils in post-column photochemical reactors for LC: application to pharmaceuticals (199) 239

Detectors
The electron-capture detector. I. Theoretical model (200) 15

Detectors
Beta-induced fluorescence as a detection technique for LC. II. Prototype detector for normal-phase operation (200) 47

Detectors
Chemiluminescence HPLC detector applied to ascorbic acid determination (200) 153

Detector time constant
Influence of detector time constant variations on efficiency calculations in the standardisation of HPLC columns (198) 235

Dexamethasone sodium phosphate
Analysis of dexamethasone sodium phosphate formulations by HPLC (193) 160

Dexsil
Applications of Dexsil and other GC porous-layer open tubular columns to metal compound separations and pyrolysis (196) 403

Dextran
Separation range and separation efficiency in high-speed gel filtration on TSK-GEL SW columns (190) 297

Diacetolol hydrochloride
Semi-automated HPLC method for the determination of diacetolol hydrochloride in medicated animal diets (200) 261

Diadenosine tetraphosphate
Detection and quantitation of diadenosine tetraphosphate by HPLC (190) 212

Diamines
Adsorption-derivatization GC-MS determination of aqueous 1,6-hexanediamine at the low and sub-ppm level (193) 271

Diamines
GLC determination of toxic diamines in permanent hair dyes (193) 277

Diaminoanisole
GLC determination of toxic diamines in permanent hair dyes (193) 277

Diaminobenzene
GLC determination of toxic diamines in permanent hair dyes (193) 277

Diaminotoluene
GLC determination of toxic diamines in permanent hair dyes (193) 277

Diazepam
Determination of some benzodiazepines and metabolites in serum, urine and saliva by HPLC (181) 227

Diazepam
HPLC and GLC determination of diazepam and nordiazepam in plasma (183) 441

Diazooxonorleucine
Determination of 6-diazo-5-oxo-L-norleucine in plasma and urine by reversed-phase HPLC of the dansyl derivative (181) 95

Dibenzodioxins
Rapid analysis of chlorinated dioxins in complex organic mixtures by application of a GC-MS-calculator system with user-developed software (189) 53

Dibenzodioxins
HPLC in the analysis of chorinated dibenzodioxins and dibenzofurans in chicken liver and wood shaving samples (197) 171

Dibenzofurans
HPLC in the analysis of chorinated dibenzodioxins and dibenzofurans in chicken liver and wood shaving samples (197) 171

Dibenzopyrones
HPLC determination of major mycotoxins produced by *Alternaria* molds (194) 89

Dibenzyl sulphoxide
Study of the interactions between substituted phenols and alcohols and dibenzyl sulphoxide by means of GC (198) 495

Dibromoethane
Determination of 1,2-dibromoethane, 1,2-dichloroethane and benzene in ambient air using porous polymer traps and GC-MS analysis with selected ion monitoring (190) 97

Dibutylmethylphenol
Determination of 2,6-di-(*tert.* -butyl)-4-methylphenol in rubber-base materials and identification of an oxidation product from 2,6-di-(*tert.* -butyl)-4-methylphenol (190) 193

Dicamba
HPLC analysis of formulations of phenoxycarboxylic acids, dicamba, ioxynil and bromoxynil mixtures (in French) (190) 401

Dicarboxylic acids
GC-MS of saturated and unsaturated dicarboxylic acids in urine (199) 181

Dichlobenil
Determination of 2,6-dichlorobenzonitrile residues in soil and water (in German) (193) 320

Dichlorobenzonitrile
Determination of 2,6-dichlorobenzonitrile residues in soil and water (in German) (193) 320

Dichloroethane
Determination of 1,2-dibromoethane, 1,2-dichloroethane and benzene in ambient air using porous polymer traps and GC-MS analysis with selected ion monitoring (190) 97

Dichloronitrosalicylanilide
Extractive alkylation of 5,2'-dichloro-4-nitrosalicylanilide (niclosamide) for GLC analysis (189) 375

Dichlorophenylmethylurea
Determination of residues of methazole and its metabolites 1-(3,4-dichlorophenyl)-3-methylurea and 1-(3,4-dichlorophenyl)urea in soil by HPLC (197) 267

Dichloropropionic acid
Improved determination of dalapon residues in plant tissues and natural water by derivatization and electron capture GC (187) 244

Diclofenac
Quantitative determination of diclofenac sodium salt in plasma by absorption spectroscopy with direct quantitation of thin-layer chromatograms (in German) (181) 512

Diclofenac
Improved GC method of determining diclofenac in plasma (183) 41

Diclofenac
Determination of total monohydroxylated metabolites of diclofenac in urine by electron-capture GLC (195) 421

Diepoxybutane
Determination of Kováts retention indices with a capillary column and electron-capture detection: application to the assay of the enzymatic conversion of 3,4-epoxy-1-butene into diepoxybutane (186) 81

Diethanolamine
GC detection of diethanolamine and its degradation products (187) 145

Diethyldithiocarbamate
Determination of disulfiram and metabolites from biological fluids by HPLC (181) 407

Diethylene glycol
GC study of solution and adsorption of hydrocarbons on glycols. I. Diethylene glycol and triethylene glycol (197) 135

Di(ethylhexyl) phthalate
Determination by electron-capture GC of mono- and di(2-ethylhexyl) phthalate in intravenous solutions stored in poly(vinyl chloride) bags (193) 405

Diffusion constant
The dynamic diffusion constant within fluid flow in an open straight tube with an elliptical cross-section (196) 349

Diflunisal
Rapid HPLC method for the quantitative determination of diflunisal in plasma (182) 487

Digitonides
Fractionation of steroid digitonides by TLC and CC on silica gel (188) 445

Diglycerides
HPLC of diglyceride *p*-nitrobenzoates. An approach to molecular analysis of phospholipids (198) 520

Digoxigenin
HPLC method for isolation of tritiated digoxin and metabolites in urine (182) 465

Digoxin
HPLC method for isolation of tritiated digoxin and metabolites in urine (182) 465

Dihydralazine
Detection of some antihypertensive drugs and their metabolites in urine by TLC. II. A further five beta blockers and dihydralazine (196) 189

Dihydroergot alkaloids
Detection and assay of dihydroergot alkaloids by TLC using *o*-phthalaldehyde-sulphuric acid reagent (193) 500

Dihydroxycholecalciferol
GC-MS properties of boronate esters of 24*R*,25-dihydroxycholecalciferol (192) 434

Diisocyanate
Determination of free toluene diisocyanate in polyurethane prepolymers by HPLC (198) 212

Dimethoxymethylsilyl ethers
Dimethoxymethylsilyl ethers: stable silyl derivatives for GC and MS (196) 156

Dimethoxypoly(ethylene glycol)
Study of the adsorption effects at the surface of poly(ethylene glycol)-coated column packings (186) 63

Dimethylarsinic acid
Graphite furnace atomic absorption spectrophotometers as automated element-specific detectors for HPLC. The determination of arsenite, arsenate, methylarsonic acid and dimethylarsinic acid (191) 31

Dimethylbenzanthracene metabolites
Combined reversed-phase and normal-phase HPLC in the purification and identification of 7,12-dimethylbenz[*a*]anthracene metabolites (185) 635

Dimethylolurea
Separation of urea-formaldehyde addition products by gel permeation chromatography (195) 154

Dimethylxanthine
New rapid assay of theophylline in plasma by isotachophoresis (181) 147

Dimethylxanthine
Micro estimation of plasma theophylline by GLC with on-column butylation and nitrogen-specific detection (182) 232

Dinitrogen
Analysis of dinitrogen-nitrogen oxide mixtures employing direct vacuum line/gas chromatograph injection (200) 224

Dinitrophenyl amino acids
New horizontal flow-through coil planet centrifuge for counter-current chromatography. III. Separation and purification of dinitrophenyl amino acids and peptides (193) 37

Diols
Use of bromine as an aid in the identification of unsaturated fatty alcohols and diols (189) 263

Dioxins
Rapid analysis of chlorinated dioxins in complex organic mixtures by application of a GC-MS-calculator system with user-developed software (189) 53

Dipalmitoyl lecithin
Quantitative determination of natural dipalmitoyl lecithin with dimyristoyl lecithin as internal standard by capillary GLC (192) 185

Dipeptides
Ligand-exchange chromatography of small peptides on copper(II)-modified silica gel: application to the study of the enzymatic degradation of methionine-enkephalin (186) 677

Dipeptides
New sensitive technique for the quantitative analysis of initiation peptides (193) 491

Dipeptides
Separation of underivatized dipeptides by HPLC on a weak anion-exchange bonded phase (195) 119

Dipeptides
Analysis of dipeptide mixtures by the combination of ion-pair reversed-phase HPLC and GC-MS techniques (197) 31

Dipeptidyl peptidase
Rapid chromatographic purification of dipeptidyl peptidase IV in human submaxillary gland (189) 233

Diphenylacetic acid
Determination of adiphenine hydrochloride and diphenylacetic acid by ion-pair HPLC (200) 267

Diphenylhydantoin
Rapid method for simultaneous determination of phenobarbital, diphenylhydantoin and their main hydroxylated metabolites by nitrogen-selective GC (183) 505

Dipole-dipole interactions
Fundamental, predictive approach to dipole-dipole interactions based on the dipole moment of the solute and the dielectric constant of the solvent (194) 105

Dipyridamole
HPLC analysis of dipyridamole in plasma and whole blood (183) 57

Disaccharides
HPLC separation of unsaturated disaccharides derived from heparan sulfate and heparin (195) 402

Disopyramide
Simultaneous analysis of disopyramide and quinidine in plasma by HPLC (183) 65

Dispersion
Early phases of the dispersion of a sample injected in Poiseuille flow (186) 353

Distribution phenomena
Role of organic modifier sorption on retention phenomena in reversed-phase LC (199) 259

Disulfiram
Determination of disulfiram and metabolites from biological fluids by HPLC (181) 407

Disulphide-polyacrylamide gels
Dissolvable disulphide-polyacrylamide gels for the electrophoretic analysis of chromosomal proteins and for affino-electrophoresis of thiol-proteins (190) 437

Disulphides
Solid-phase reagents for LC detection (185) 497

Disulphur decafluoride
Adsorption-desorption GC-IR determination of trace disulfur decafluoride in sulfur hexafluoride (193) 265

Diterpenes
Rapid and micro identification of biologically active diterpenes in *Rabdosia umbrosus* var. *excisinflexus* (Labiatae) by reversed-phase HPLC (200) 250

Dithiadene
Determination of *cis*- and *trans*-isomers of dosulepine and dithiadene by HPLC (191) 137

Dithiols
Derivatization of dithiols and certain monothiols with phenylarsine oxide for GC (200) 171

Diuretic drugs
Rapid and sensitive determination of chlorthalidone in blood, plasma and urine of man using HPLC (181) 497

Diuretic drugs
Quantitative GC determination of two oxidized metabolites of the diuretic mefruside in human urine, plasma and red blood cells (182) 179

Diuretic drugs
Separation and analysis of azosemide in urine and in serum by HPLC (182) 257

Diuretic drugs
Determination of the diuretic bumetanide in biological fluids by HPLC (183) 118

Divinylbenzene
Structure and sorption properties of porous copolymers of styrene, acrylonitrile and divinylbenzene (196) 75

DNA
Preparative isolation of deoxyriboadenylic acids from hydrolysates of oxidized herringsperm DNA using template-chromatography (in German) (187) 119

DNA
Preparation of oligomers of deoxyriboguanylic acids by partial chemical hydrolysis of DNA (in German) (196) 435

DNA fragments
Separation and sequencing of the sequence isomers of pyrimidine deoxypentanucleoside tetraphosphates by HPLC (188) 273

DNase
Specificity of deoxyribonuclease hydrolysis determined by anion-exchange HPLC (193) 175

Dns-amino acids
Reversed-phase separation of optical isomers of Dns-amino acids and peptides using chiral metal chelate additives (185) 323

Dns-amino acids
Two-dimensional TLC of Dns-amino acids on reversed-phase silica gel (193) 421

Dns-amino acids
Reversed-phase HPLC resolution of D- and L-Dns-amino acids by mixed chelate complexation (199) 295

Docosanoylvalinemethylheptadecylamide
N-Docosanoyl-L-valine-2(2-methyl)-*n*-heptadecylamide as a stationary phase for the resolution of optical isomers in GLC (195) 317

Documentation
Instant copy technique for documentation of thin-layer chromatograms (189) 284

DOPA
Electrochemical detector for HPLC (192) 135

DOPA
Rapid and sensitive assay of tyrosine 3-monooxygenase activity by HPLC using the native fluorescence of DOPA (198) 511

Dopamine
Simplified LC-electrochemical determination of norepinephrine and dopamine in rat brain (181) 453

Dopamine
Simple and sensitive procedure for the assay

of serotonin and catecholamines in brain by HPLC using fluorescence detection (182) 277

Dopamine
Simple method for isolating free urinary catecholamines by boric acid gel chromatography (183) 78

Dopamine
Investigation of the performance of a HPLC system with an electrochemical detector (183) 261

Dopamine
Comparison of LC-electrochemical and GC-MS methods for brain dopamine and serotonin (183) 483

Dopamine
Electrochemical detector for HPLC (192) 135

Dopamine
GC-MS determination of dopamine in subregions of rat brain (198) 329

Dopants
Oxygen enhancement of the electron capture detector response of carbon dioxide (188) 89

Dopants
Effect of dopants on the linearity of a d.c. electron-capture detector (192) 413

Dopants
Alteration of the electron-capture detector response to polycyclic aromatic hydrocarbons by oxygen doping of the carrier gas (197) 51

Doping agents
Survey of selected hydrides as doping agents for a hydrogen-atmosphere flame-ionization detector (195) 211

Doping drugs
Reversed-phase HPLC of phenylbutazone in body fluids (181) 132

Doping drugs
Qualitative screening procedure for the detection of benzoyl ecgonine in the urine of greyhounds (196) 337

Dosulepine
Determination of *cis*- and *trans*-isomers of dosulepine and dithiadene by HPLC (191) 137

Dothiepin
Chemical ionisation mass fragmentographic measurement of dothiepin plasma concentrations following a single oral dose in man (183) 141

Doxapram
Quantitation of doxapram in blood, plasma and urine (182) 191

Doxycycline
Rapid analysis of doxycycline from biological samples by HPLC (187) 439

Droplet counter-current chromatography
Application of droplet counter-current chromatography to the isolation of natural products (186) 529

Dropping mercury electrode amperometric detector
Design and selective application of a dropping mercury electrode amperometric detector in column LC (186) 489

Drug monitoring
HPLC in the elucidation of the metabolic fate of butoprozine. Screening for unknown metabolites in the rat (191) 109

Drug-protein binding parameters
Usefulness of chromatographic methods for the determination of drug-protein binding parameters (193) 522

Drug-protein interactions
Mutual displacement interactions in the binding of two drugs of human serum albumin by frontal affinity chromatography (188) 347

Drugs, basic
Use of ninhydrin as a spray reagent for the detection of some basic drugs on thin-layer chromatograms (195) 133

Drug screening
GLC retention indices of 296 non-drug substances on SE-30 or OV-1 likely to be encountered in toxicological analyses (Review) (184) 185

Drugs of abuse
Analysis of therapeutic and commonly abused drugs in serum and urine by GLC using a photoionization detector (186) 637

Drugs of abuse
Mass screening and confirmation of methaqualone and its metabolites in urine by radioimmunoassay-TLC (190) 129

Drugs of abuse
Separation and determination of opium alkaloids by HPLC (190) 251

Drugs of abuse
Positive identification and quantitation of isomeric cocaines by HPLC (193) 371

Drugs of abuse
Simple, rapid method for extraction of urinary cannabinoids by LSC (196) 342

Drugs of abuse
Qualitative differentiation between cocaine, lidocaine and cocaine-lidocaine mixtures ("rock-cocaine") using TLC (198) 534

Drugs of abuse
Use of dual-wavelength UV detection in HPLC for the identification of barbiturates (200) 271

Durable press agents
Separation of methylol derivatives of imidazolidines, urea and carbamates by LC (190) 226

Dyes
Resolution and retention behaviour of some dyes in overpressured TLC (191) 293

Dyes
GLC determination of toxic diamines in permanent hair dyes (193) 277

Dyes
Surface ion exchange and adsorption of some dyes on α-$Zr(HPO_4)_2 \cdot H_2O$ micro-crystals (195) 270

Dyes, organic
TLC of slightly soluble substances at elevated temperature (186) 575

Dynamic diffusion constant
The dynamic diffusion constant within fluid flow in an open straight tube with an elliptical cross-section (196) 349

Dyphylline
Use of flow programming in paired-ion HPLC analysis of dosage forms containing dyphylline (193) 111

E

Ecdysteroids
Practical aspects of the preparation and chromatography of the trimethylsilyl ethers of ecdysteroids (194) 43

Ecdysteroids
Comparison of HPLC and GC for the analysis of ecdysteroids (194) 343

Edrophonium
Reversed-phase, ion-pair LC of quaternary ammonium compounds. Determination of pyridostigmine, neostigmine and edrophonium in biological fluids (183) 193

Efficiency calculations
Influence of detector time constant variations on efficiency calculations in the standardisation of HPLC columns (198) 235

Elastase
One-step purification of human leukocyte elastase by biospecific affinity chromatography at subzero temperatures (192) 236

Electrochemical detector
Small-volume electrochemical detector for microcolumn LC (181) 287

Electrochemical detector
Application of an electrochemical detector to the determination of procarbazine hydrochloride by HPLC (190) 359

Electrochemical detector
Electrochemical detector for HPLC (192) 135

Electrokinetic detection
Measurement of exclusion volumes of packed columns by means of electrokinetic detection (191) 3

Electron-capture detector
Analysis of kinetics and concentration dependence of electron-capture detector (186) 99

Electron-capture detector
Studies and speculations on an alternative mechanism for the response of direct-current electron-capture detectors (188) 1

Electron capture detector
Oxygen enhancement of the electron capture detector response of carbon dioxide (188) 89

Electron capture detector
Analysis errors following hydrogen cleaning of an electron capture detector (190) 133

Electron-capture detector
Effect of dopants on the linearity of a d.c. electron-capture detector (192) 413

Electron capture detector
Comparative investigation of radioactive sources for the electron capture detector (195) 1

Electron capture detector
Effects of dimension and density in electron capture detectors (195) 17

Electron-capture detector
Reaction and response in electron-capture detectors (196) 397

Electron-capture detector
Alteration of the electron-capture detector response to polycyclic aromatic hydrocarbons by oxygen doping of the carrier gas (197) 51

Electron-capture detector
The electron-capture detector. I. Theoretical model (200) 15

Electron-capture sensitization
Selective electron-capture sensitization (186) 3

Electronic signal differentiation
Electronic singal differentiation as an aid to quantitation in GC (197) 1

Electrophoretic fractionation systems
Mass, charge, momentum and energy conservation in electrophoretic fractionation systems (Review) (184) 21

Electrophoretic mobility
Determination of the free electrophoretic mobility of proteins by polyacrylamide gradient gel electrophoresis: A new approach (192) 402

Element-specific detection
Capacilities of on-line element-specific detection in HPLC using an inductively coupled argon plasma emission source detector (185) 549

Eluent composition
Mobile phase effects in reversed-phase chromatography. III. Changes in conformation and retention of oligo(ethylene glycol) derivatives with temperature and eluent composition (185) 129

Eluents
Anion chromatography with low-conductivity eluents (186) 509

Eluents
Anion chromatography with low-conductivity eluents. II (187) 35

Elution curves
Influence of sorption on the shape of chromatographic elution curves (198) 377

Elution, non-linear
Study of non-linear elution in preparative LC (in French) (192) 53

Emetic drugs
Determination of apomorphine in plasma and brain tissue by ion-pair extraction and LC (185) 575

Enantiomer separation
Separation of enantiomers by HPLC on triacetylcellulose (193) 308

Enantiomer separation
Quantitative determination of D = L mixtures of optical enantiomers by GC (198) 185

Enantiomer separation
Structural requirements for enantioselectivity in GC of chiral α-hydroxy acids (200) 189

Enantiomer separation
Resolution of amino acid enantiomers by glass capillary GC on easily prepared optically active stationary phases (200) 195

Endogenous compounds
GLC retention indices of 296 non-drug substances on SE-30 or OV-1 likely to be encountered in toxicological analyses (Review) (184) 185

Energy of interaction
Determination of the energy of intermolecular interactions in GC from retention indices (190) 1

Enkephalin
Ligand-exchange chromatography of small peptides on copper(II)-modified silica gel: application to the study of the enzymatic degradation of methionine-enkephalin (186) 677

Enkephalin
Combination of enzymatic and mass fragmentographic assays for the identification and measurement of [met^5]-enkephalin (190) 43

Enzyme inhibitor
Determination of captopril in biological fluids by GLC (188) 177

Enzyme kinetics
Determination of catecholamines and O-methylated metabolites by reversed-phase HPLC with fluorimetric detection and its application to enzyme kinetics (196) 101

Enzymes
Use of water-soluble biospecific polymers for the purification of proteins (Review) (184) 325

Enzymes
Purification of enzymes by heparin-Sepharose affinity chromatography (Review) (184) 335

Enzymes
Preparative separation of proteins and enzymes in the mean molecular-weight range of 10,000-100,000 on LiChrosorb Diol packing by high-performance size-exclusion chromatography (185) 445

Enzymes
High-speed steric exclusion chromatography of biopolymers (185) 453

Enzymes
Solid-phase reagents for LC detection (185) 497

Enzymes
Experimental taxonomy of *Biomphalaria* (gastropoda: planorbidae). I. Methods for experimental taxonomic studies on *Biomphalaria* carried out by horizontal starch gel electrophoresis and staining of twelve enzymes (188) 169

Enzymes
Rapid method for the determination of urokinase activity by HPLC (188) 185

Enzymes
Assay of phospholipase D on a TLC plate (188) 292

Enzymes
Affinity chromatography of a cellulase complex from *Trichoderma viride* on cross-linked cellulose (in French) (188) 377

Enzymes
Assay method for urokinase activity by capillary-tube isotachophoresis using a synthetic substrate (188) 383

Enzymes
Improved GC method for measuring phenylethylene glycol. Application to the determination of styrene monooxygenase and epoxide hydrase activities (188) 400

Enzymes
Polyacrylamide-boronate beads saturated with biomolecules: a new general support for affinity chromatography of enzymes (189) 225

Enzymes
Rapid chromatographic purification of dipeptidyl peptidase IV in human submaxillary gland (189) 233

Enzymes
Efficient purification of a microbial steroid-1-dehydrogenase by electrophoretic desorption from the affinity matrix on a preparative scale (189) 279

Enzymes
Purification of enzymes by high-speed gel filtration on TSK-GEL SW columns (190) 305

Enzymes
High-capacity method for purification of human liver hexosaminidase B using hydrophobic chromatography (190) 385

Enzymes
Purification of grape polyphenoloxidase with hydrophobic chromatography (192) 232

Enzymes
One-step purification of human leukocyte elastase by biospecific affinity chromatography at subzero temperatures (192) 236

Enzymes
Study of the interaction between L-lactate dehydrogenase isoenzymes and immobilized 8-substituted adenosine 5'-monophosphate by means of affinity electrophoresis (192) 351

Enzymes
Specificity of deoxyribonuclease hydrolysis determined by anion-exchange HPLC (193) 175

Enzymes
Reversed-phase HPLC of adenine nucleotides: application to the kinetics of an adenosine 3'-phosphate 5'-sulphatophosphate sulphotransferase from plants (193) 285

Enzymes
Affinity chromatography of rat liver lactate dehydrogenase on the Remazol derivative of bead cellulose (194) 95

Enzymes
Interaction of trypsin with immobilized *p*-aminobenzamidine derivatives studied by means of affinity electrophoresis (194) 175

Enzymes
Determination of the interaction of lactate dehydrogenase with high-molecular-weight derivatives of AMP by affinity electrophoresis (194) 183

Enzymes
Affinity chromatographic interactions of proteases with low-molecular-weight soybean protease inhibitors (195) 385

Enzymes
Study of the interaction between NADP-dependent dehydrogenase and immobilized adenosine 2'-monophosphate by means of affinity electrophoresis (196) 85

Enzymes
Purification of an α-L-arabinofuranosidase from *Aspergillus niger* by substrate affinity chromatography (197) 86

Enzymes
Rapid and sensitive assay of tyrosine 3-monooxygenase activity by HPLC using the native fluorescence of DOPA (198) 511

Enzymes
In vivo studies of the tryptophan-5-hydroxylase system. Quantitation of serotonin and tryptamine using GC-mass fragmentography (199) 171

Enzymes
Optimized assay for purine nucleoside phosphorylase by reversed-phase HPLC (199) 275

Ephedrine
Analysis of perfluoroacyl derivatives of ephedrine, pseudoephedrine and analogues by GC and MS. I. Trifluoroacetyl derivatives (190) 53

Epinephrine
GC of epinephrine and norepinephrine after derivatization with chloroformates in aqueous media (190) 347

Epioxytetracycline
Reversed-phase ion pair chromatography of oxytetracycline, epioxytetracycline and anhydrooxytetracycline (190) 486

Epoxide hydrase
Improved GC method for measuring phenylethylene glycol. Application to the determination of styrene monooxygenase and epoxide hydrase activities (188) 400

Epoxides
Determination of trace amounts of hydrogen chloride by derivatization with epoxides and GC separation (186) 219

Epoxybutene
Determination of Kováts retention indices with a capillary column and electron-capture detection: application to the assay of the enzymatic conversion of 3,4-epoxy-1-butene into diepoxybutane (186) 81

Epoxy resins
Separation and characterization of by-product oligomers in epoxy resins by reversed-phase HPLC (193) 243

Equivalent chain length
GC of esters. XIII. Interrelationship of equivalent chain length *(ECL)* and retention index values of fatty esters (187) 297

Ergot alkaloids
Detection and assay of dihydroergot alkaloids

by TLC using *o*-phthalaldehyde-sulphuric acid reagent (193) 500

Ergotamine
Continuous post-column ion-pair extraction detection of some basic organic compounds in normal-phase chromatography (185) 473

Ergotoxine alkaloids
Separation of ergotoxine alkaloids by HPLC on silica (191) 101

Erythrocytes
Surface affinity chromatographic separation of blood cells. I. Separation of human and rabbit peripheral granulocytes, lymphocytes and erythrocytes using polyethylene glycol-bonded column packings (187) 351

Erythrocytes
Electrofocusing of stroma-free hemolyzates of human erythrocytes; comparison of fresh, stored and chemically modified samples (193) 515

Essential oils
Study of three types of essential oil of *Valeriana officinalis* L. s.l by combined GC-negative ion chemical ionization MS (190) 321

Essential oils
Quantitative determination of minor components in essential oils: determination of pulegone in peppermint oils (190) 471

Estramustine
Determination of estramustine and its 17-keto metabolite in plasma by HPLC (182) 387

Estrogens, see oestrogens

Ethanol
Simple rapid and sensitive method for the simultaneous quantitation of ethanol and acetaldehyde in biological materials using head-space GC (190) 197

Ethaverine
Determination of the antispasmodic agent ethaverine in human plasma by HPLC (182) 379

Ethionine
The isolation of methionine and ethionine by silver ligand chromatography and application to methionine-containing peptides (200) 185

Ethosuximide
Simple, rapid procedure for the determination of valproate and ethosuximide in plasma by GLC (182) 237

Ethoxycarbonylarylazonitroethanes
TLC separation of potential antineoplastic agents: 1-ethoxycarbonyl-2-arylazo-2-nitroethanes (198) 224

Ethyl anthranilate azopigments
Accurate and sensitive analysis of ethyl anthranilate azopigments from bile by reversed-phase HPLC (182) 105

Ethylenediaminetetraacetic acid
Determination of multifunctional carboxylic acids in the presence of iron by GC (190) 188

Ethylene oxide condensates
Improved HPLC analysis of ethylene oxide condensates by their esterification with 3,5-dinitrobenzoyl chloride (187) 261

Ethylphenols
Resolution of ethylphenols following *in situ* reaction on a thin-layer plate with Fast Blue B Salt (192) 425

Evaporative analyser detector
Performance of an evaporative analyser detector for gel permeation chromatography (189) 259

Exclusion volumes
Measurement of exclusion volumes of packed columns by means of electrokinetic detection (191) 3

Expectorant drugs
The determination of bifunctional compounds. IX. A selective reaction for the determination of guaifenesin in plasma by GC (183) 433

Extra-column effects
Extra-column effects in HPLC. I. Theoretical study of the injection problem (185) 79

Extrelut columns
Contamination by use of Extrelut columns (in German) (183) 1

F

Fatty acids
GC determination of the fatty acid pattern of red cell membrane plasmalogens in healthy children (182) 27

Fatty acids
The use of formic acid in carrier gas. Rapid method for identification and determination of phytanic acid by GC-MS-chemical ionization (182) 269

Fatty acids
Detection of tuberculostearic acid in biological specimens by means of glass capillary GC-electron and chemical ionization MS, utilizing selected ion monitoring (182) 402

Fatty acids
Applications of reversed-phase chromatography and nephelometric detection to analysis of non-polar mixtures at microgram level (187) 111

Fatty acids
Derivatization of fatty acids with 1-chlormethylisatin for HPLC (187) 208

Fatty acids
Fractionation of fish oil fatty acid methyl esters by means of argentation and reversed-phase HPLC, and its utility in total fatty acid analysis (187) 307

Fatty acids
GC-MS of the flophemesyl derivatives of a series of fatty acid methyl ester chlorohydrins (189) 86

Fatty acids
Use of bromine as an aid in the identification of unsaturated fatty alcohols and diols (189) 263

Fatty acids
Phenanthrimidazoles as fluorescent derivatives in the analysis of fatty acids by HPLC (189) 359

Fatty acids
Identification of characteristic branched-chain fatty acids of *Mycobacterium kansasii* and *gordonae* by GC-MS (190) 183

Fatty acids
Modified preparation of 2-naphthacyl esters of fatty acids and their fluorometric detection in HPLC (in German) (192) 240

Fatty acids
Fatty acids. XIX. A quantitative treatment of saturated triglycerides by reversed-phase HPLC (192) 457

Fatty acids
Isolation of 14-methylhexadecanoic acid from wool fat by preparative GLC (193) 153

Fatty acids
Separation of long-chain fatty acids extracted from peat by gel permeation chromatography (in French) (193) 329

Fatty acids
GLC separation of geometric isomers of unsaturated fatty acid methyl esters using a glass capillary column (194) 404

Fatty acids
Metabolic profiles of penaeid shrimp: dietary lipids and ovarian maturation (195) 359

Fatty acids
Analysis of fatty acid methyl esters by high-resolution GC-chemical ionisation MS (196) 415

Fatty acids
Coating support-coated open tubular capillary columns with Silar 10C or Alltech CS-10 and Silica T40 for separation of isomers of fatty acid methyl esters (198) 153

Fatty alcohols
Use of bromine as an aid in the identification of unsaturated fatty alcohols and diols (189) 263

Fatty alcohols
4-(*p*-methoxycinnamyloxy)-4'-methoxy-azobenzene: a nematic liquid crystal for the GLC analysis of the stereochemistry of lepidopterous sex pheromones and related unsaturated fatty alcohols and derivatives (190) 35

Fatty esters
GC of esters. XIII. Interrelationship of equivalent chain length *(ECL)* and retention index values of fatty esters (187) 297

Feed mixtures, animal
HPLC column switching technique in the analysis of medicated feed for an automated clean-up procedure (198) 162

Fenoprofen
High-performance liquid column chromatography of fenoprofen in serum (183) 250

Ferric ions
Complexation with ferric ions: a potential source of error in the isotachophoretic determination of oxalate (188) 266

Ferrocene alcohols
TLC separation of some ferrocene alcohols (198) 536

Field-flow fractionation
Retention characteristics of various polymers in thermal field-flow fractionation (186) 37

Fingerprinting
Fingerprinting of heparins by low-amperage electrophoresis in barium acetate (195) 398

Firemaster FF-1
GC-MS analysis of polybrominated biphenyl constituents of Firemaster FF-1 (197) 258

Fish oil
Fractionation of fish oil fatty acid methyl esters by means of argentation and reversed-phase HPLC, and its utility in total fatty acid analysis (187) 307

Flame-ionization detector
Survey of selected hydrides as doping agents for a hydrogen-atmosphere flame-ionization detector (195) 211

Flavones
Separation of polymethoxylated flavones by straight-phase HPLC (190) 233

Flavonoids
HPLC, GLC and TLC of naturally occurring flavonoids, phenolic and related compounds (187) 255

Flavonoids
Analysis of flavonoids by HPLC (189) 217

"Flip-flop" chromatography
"Flip-flop" chromatography (186) 15

Flow-impedance bridge detector
New results with a flow-impedance bridge detector (189) 3

Flow segmentation
Coated-open-tubular chromatography with flow segmentation. I. Theory (185) 43

Flow segmentation
Coated-open-tubular chromatography with flow segmentation. II. Experimental study and optimization for size-exclusion separation (185) 57

Flow-through polarographic detector
Polarographic detection for HPLC using a flow-through detector (191) 47

Flufenamic acid
Quantitative determination of flufenamic acid in rat plasma and uterus by GC (192) 441

Flunitrazepam
GC determination of bromo and fluoro derivatives of benzodiazepine in human body fluids (182) 81

Fluorescence detector
Beta-induced fluorescence as a detection technique for LC. II. Prototype detector for normal-phase operation (200) 47

Fluorescence micro disc electrophoresis
Fluorescence micro disc electrophoresis in sodium dodecyl sulfate. A simple and sensitive method applicable for routine laboratory analysis (181) 259

Fluorescent compounds
Separation of fluorescent compounds in members of the genus *Rhizopus* by TLC (187) 399

Fluorescent derivatives
Phenanthrimidazoles as fluorescent derivatives in the analysis of fatty acids by HPLC (189) 359

Fluoropyridinecarboxylic acid
Hydrophobic chromatography and bioanalysis of some polar pyridine derivatives used as antilipolytic agents (187) 189

Fluoropyrimidinedione derivatives
Quantitative determination of 1,3-bis(tetrahydro-2-furanyl)-5-fluoro-2,4-pyrimidinedione and its metabolites in visceral tissues by HPLC and GC mass fragmentography (188) 270

Fluorouracil derivatives
Determination of two new metabolites of 1-hexylcarbamoyl-5-fluorouracil in biomedical specimens by HPLC (182) 125

Fluorouracil nucleosides
Reversed-phase LC of 5-fluorouracil nucleosides and nucleotides in the presence of quaternary ammonium ions (194) 305

Fluphenazine
HPLC separation of the N- and S-oxides of fluphenazine and fluphenazine decanoate (194) 416

Fluvoxamine
The use of a pre-column for the direct HPLC determination of the antidepressants clovoxamine and fluvoxamine in plasma (183) 203

FM 24
Sensitive GC method for the determination in plasma of FM 24, 1-(2-*exo*-bicyclo[2,2,1]hept-2-ylphenoxy)-3-[(1-methylethyl)amino]-2-propanol, a new β-adrenoceptor blocking agent (183) 99

Folate derivatives
Separation of folate derivatives by *in situ* paired-ion HPLC (190) 237

Folic acid derivatives
HPLC separation of clinically important folic acid derivatives using ion-pair chromatography (190) 241

Food additives
GLC retention indices of 296 non-drug substances on SE-30 or OV-1 likely to be encountered in toxicological analyses (Review) (184) 185

Food contamination
Determination of styrene in foods by reversed-phase HPLC (198) 198

Food preservatives
GC determination of nitrite in foods as trimethylsilyl derivative of 1*H*-benzotriazole (194) 21

Formaldehyde
GLC analysis of dimedone derivatives of formaldehyde and other aliphatic aldehydes on capillary columns (191) 239

Formaldehyde
Separation of urea-formaldehyde addition products by gel permeation chromatography (195) 154

Formaldehyde
TLC method for the separation and determination of the products of the reaction of amides with formaldehyde (195) 158

Formaldehyde
Determination of formaldehyde, acetaldehyde and associated components in solution and in vapours by GC (200) 206

Fractionation of cells
New experimental approaches to the isoelectric fractionation of cells (194) 323

Free displacement electrophoresis
Free displacement electrophoresis (isotachophoresis): an absolute determination of the Kohlrausch functions and their use in interaction studies (194) 1

Fructose
Separation and quantitative determination of

fructose as the O-methyloxime by GLC using glass capillary columns (192) 446

Functional group behaviour
Functional group behaviour in ion-pair reversed-phase HPLC using surface-active pairing ions (185) 197

Fungicides
Analysis of pentachlorophenol in waste water using HPLC (190) 203

Fungicides
Simple TLC method for detection of pentachlorophenol in sawdust and woodshavings (195) 441

Fungicides
GC determination of Metalaxyl (Ridomil) residues in tobacco (200) 313

Fungicides
Estimation of the systemic fungicide Ridomil by TLC (200) 317

Furazolidone
HPLC analysis of furazolidone in liver and kidney (198) 526

Furfural
Determination of furfural in spent sulfite liquor by GC (194) 228

G

Galactosidase
Purification of enzymes by high-speed gel filtration on TSK-GEL SW columns (190) 305

Galacturonic acid oligomers
Separation of α-D-galacturonic acid oligomers by chromatography on polyacrylamide gel (194) 315

Galanthamine
New method for the quantitative analysis of galanthamine in drug extracts from *Leucojum aestivum* (in German) (188) 452

Gangliosides
Brain gangliosides: an improved simple method for their extraction and identification (195) 257

Gas analysis
Comparison between conventional and cross-correlation GC for evolved gas analysis (186) 145

Gel particles
New cellulose gel for chromatography (195) 221

Gentamicin
Micro determination of gentamicin in serum by HPLC with UV detection (182) 201

Gentiana root
Determination of the bitter constituents of the Gentiana root by HPLC (193) 163

Gentiobioside
Determination of the bitter constituents of the Gentiana root by HPLC (193) 163

Gentiopicroside
Determination of the bitter constituents of the Gentiana root by HPLC (193) 163

Geomatrices
Elucidation of geomatrices by laser pyrolysis-GC and pyrolysis-MS (186) 831

Germanium compounds
GC detection of organogermanium compounds by surface or gas-phase luminescence (186) 299

Germatranes
Chromatography of organometallic and organometalloidal derivatives of amino alcohols. II. GC evidence for transannular bonds in solutes (200) 105

Gibberellins
High-performance steric exclusion chromatography of plant hormones (198) 57

Gibberellins
HPLC of gibberellins (198) 449

Ginsenosides
GLC method for determination of ginsenosides in *Panax ginseng* (196) 121

Glass capillary columns
Coating of glass capillary chromatographic columns: a new variant of the static method (190) 445

Glass coating
Use of silicone-coated porous glass for exclusion chromatography in an aqueous medium (196) 485

Glass columns
Glass columns with high resistance to internal pressure for HPLC (193) 128

Glass surfaces
Glass surface analytical studies in the preparation of open tubular columns for GC (199) 355

Glaucine
HPLC determination of glaucine and its metabolite, dehydroglaucine in microbial culture (193) 170

Glibornurid
Detection of glibornurid in serum by reversed-phase HPLC (in German) (183) 115

Gliclazide
Determination of gliclazide in human serum by HPLC using an anion-exchange resin (183) 467

Glipizide
Electron-capture GC of plasma sulphonylureas after extractive methylation (181) 17

Glucanases
Affinity chromatography of a cellulase complex from *Trichoderma viride* on cross-linked cellulose (in French) (188) 377

Glucose
Thermal degradation of sugars in a GC injection port (193) 83

Glucosiduronic esters
Liquid ion exchanges in PC of steroidal glucosiduronic esters (189) 33

Glucosinolates
Separation of glucosinolates by HPLC (197) 199

Glucuronic acid
New method for the quantitative assay of the enzymatic hydrolysis of UDPglucuronic acid using analytical capillary isotachophoresis (188) 235

Glucuronide conjugates
HPLC of glucuronide and sulfate conjugates using ion-pair chromatography (195) 431

Glucuronides
Separation of steroid glucuronides by reversed-phase LCC (194) 80

Glucuronides
Evaluation of Amberlite XAD-4 for recovery of glucuronides and sulfates from aqueous solution (196) 303

Glutamylcysteine
Estimation of glutathione in rat liver by reversed-phase HPLC: separation from cysteine and γ-glutamylcysteine (194) 424

Glutathione
Estimation of glutathione in rat liver by reversed-phase HPLC: separation from cysteine and γ-glutamylcysteine (194) 424

Glycidylmethacrylate copolymers
Reactive polymers. XXXI. Thermally treated polymeric sorbents based on glycidylmethacrylate copolymers (194) 297

Glycine
Simple and rapid mass fragmentographic method for the determination of glycine in brain tissue (193) 95

Glycine conjugates
Rapid determination of glycine- and taurine-conjugated bile acids in human bile by HPLC (181) 177

Glycine oligomers
Ligand-exchange chromatography of small peptides on copper(II)-modified silica gel: application to the study of the enzymatic degradation of methionine-enkephalin (186) 677

Glycoalkaloids
Optical brighteners as TLC detection reagents for glycoalkaloids and steroid alkaloids in *Solanum* species. II. Blankophor BA 267%, BBU neu and KU, and Tinopal CBS-X and 5 BMS-X (189) 406

Glycolic acid
Quantitative determination of quinic acid and derivatives by HPLC after derivatization with *p*-bromophenacyl bromide (190) 411

Glycollic acid
Decreased accuracy in isotachophoretic analysis due to septum bleed (188) 262

Glycols
GC study of solution and adsorption of hydrocarbons on glycols. I. Diethylene glycol and triethylene glycol (197) 135

Glycoproteins
Differentiation of the polysaccharide side-chains of glycoproteins by a fingerprinting technique: heterogeneity of human gastric mucin (181) 108

Glycoproteins
Purification of ovomucoid by hydroxyapatite chromatography (187) 224

Glycoproteins
Isolation of water-soluble plant glycoproteins by affinity chromatography on immobilised concanavalin A (in French) (188) 435

Glycoproteins
Use of deoxycholate in gel filtration chromatography of synaptosomal membrane glycoproteins (193) 316

Glycoproteins
Interactions of insolubilized lectins with membrane glycoproteins in presence of detergents (195) 197

Glycosaminoglycans
Determination of molecular-weight distribution of aorta glycosaminoglycans by automated gel filtration (192) 307

Glycosides
Application of droplet counter-current chromatography to the isolation of natural products (186) 529

Glycosphingolipids
Brain gangliosides: an improved simple method for their extraction and identification (195) 257

Glyoxylic acid
Decreased accuracy in isotachophoretic analysis due to septum bleed (188) 262

Gradient elution
Influence of organic modifiers on the retention behaviour in reversed-phase LC and its consequences for gradient elution (185) 179

Gradient elution
Gradient elution in LC. XI. Influence of the

adjustable gradient parameters on the chromatographic behaviour of sample compounds (192) 1

Gradient elution
Gradient elution in LC. XII. Optimization of conditions for gradient elution (192) 19

Gradient elution
Gradient elution in LC. XIII. Instrumental errors in gradient elution chromatography (192) 37

Gradient spreader
New spreader for the preparation of gradient layers in TLC (189) 293

Granulocytes
Surface affinity chromatographic separation of blood cells. I. Separation of human and rabbit peripheral granulocytes, lymphocytes and erythrocytes using polyethylene glycol-bonded column packings (187) 351

Graphite furnace atomic absorption detector
Application of a graphite furnace atomic absorption detector automatically coupled to a high-performance liquid chromatograph for speciation of metal-containing macromolecules (185) 563

Graphite furnace atomic absorption spectrophotometer
Graphite furnace atomic absorption spectrophotometers as automated element-specific detectors for HPLC. The determination of arsenite, arsenate, methylarsonic acid and dimethylarsinic acid (191) 31

Griseofulvin
Chromatographic analysis of griseofulvin and metabolites in biological fluids (181) 77

Guaiacols
TLC of chlorinated guaiacols (194) 55

Guaifenesin
The determination of bifunctional compounds. IX. A selective reaction for the determination of guaifenesin in plasma by GC (183) 433

H

Haemoglobin
Quantitative determination of hemoglobin A_1c by thin-layer isoelectric focusing (182) 325

Haemoglobin
Electrofocusing of stroma-free haemoglobin on a thin layer of agarose gel (196) 142

Haemoglobin Bart's
Carboxymethyl-cellulose microchromatography for the quantitation of hemoglobin Bart's (γ_4) and its use in the detection of the α-thalassemia conditions (198) 443

Halcinonide
HPLC of the topical anti-inflammatory steroid halcinonide (190) 481

Halides
Use of isothermal GC for the determination of the adsorption enthalpies and entropies of inorganic halides at high temperatures (187) 319

Hallucinogenic compounds
Determination of hallucinogenic components of *Psilocybe* mushrooms using HPLC (196) 180

Halogenated anilines
Determination of halogenated anilines in urine by HPLC with an electrochemical detector (188) 412

Halogenated uracil derivatives
Separation of halogenated uracil derivatives from nucleobases and nucleosides by TLC on silica gel (187) 453

Harmane alkaloids
Fluorimetric and HPLC determination of harmane alkaloids in *Peganum harmala* cell cultures (194) 234

HA-966
Assay of HA-966 in rat plasma by capillary GLC with nitrogen-selective detection (183) 185

Heater displacement chromatography
Quantitative aspects of heater displacement chromatography (197) 109

Heparan sulphate
HPLC separation of unsaturated disaccharides derived from heparan sulfate and heparin (195) 402

Heparin
Purification of enzymes by heparin-Sepharose affinity chromatography (Review) (184) 335

Heparin
HPLC separation of unsaturated disaccharides derived from heparan sulfate and heparin (195) 402

Heparin
Fractionation and identification of heparin and other acidic mucopolysaccharides by a new discontinuous electrophoretic method (196) 455

Heparins
Fingerprinting of heparins by low-amperage electrophoresis in barium acetate (195) 398

Heptadienes
Identification of *n*-hepta- and *n*-octadienes by high-resolution GC using structure-retention correlations and MS (191) 187

Heptaene macrolide antibiotics
Improved HPLC method for the comparison of heptaene macrolide antibiotics (189) 249

Herbicides
Improved determination of dalapon residues in plant tissues and natural water by derivatization and electron capture GC (187) 244

Herbicides
Comparison of the HPLC behaviour of *s*-triazine derivatives on various stationary phases (187) 341

Herbicides
GC determination of Lenacil and Pyrazon in soil and bacterial cultures (189) 89

Herbicides
Analysis of pentachlorophenol in waste water using HPLC (190) 203

Herbicides
HPLC analysis of formulations of phenoxycarboxylic acids, dicamba, ioxynil and bromoxynil mixtures (in French) (190) 401

Herbicides
Determination of 2,6-dichlorobenzonitrile residues in soil and water (in German) (193) 320

Herbicides
GC measurement of volatility of herbicide esters (195) 75

Herbicides
Direct analysis of the herbicide barban in wheat products by reversed-phase LC (195) 113

Herbicides
Analysis of barban formulations by HPLC (195) 151

Herbicides
Determination of residues of methazole and its metabolites, 1-(3,4-dichlorophenyl)-3-methylurea and 1-(3,4-dichlorophenyl)urea in soil by HPLC (197) 267

Herbicides
Determination of amitrole by ion-pair HPLC (200) 234

HETP
The HETP of an unretained component of fluid flow in a coiled round pipe (186) 341

Hexacyanoferrate compounds
Gel filtration of soluble coloured metal hexacyanoferrate(II) compounds (187) 458

Hexamethylenetetramine
Determination of methenamine in biological samples by GLC (181) 11

Hexamethylmelamine
Determination of hexamethylmelamine and metabolites in plasma or serum by GLC with a nitrogen-sensitive detector (181) 363

Hexanediamine
Adsorption-derivatization GC-MS determination of aqueous 1,6-hexanediamine at the low and sub-ppm level (193) 271

Hexosaminidase
High-capacity method for purification of human liver hexosaminidase B using hydrophobic chromatography (190) 385

Hexylcarbamoylfluorouracil
Determination of two new metabolites of 1-hexylcarbamoyl-5-fluorouracil in biomedical specimens by HPLC (182) 125

High-performance liquid chromatography
A hazard in high-performance liquid chromatography (189) 108

High-speed isotachophoresis
High-speed isotachophoresis. Effect of driving current on the qualitative interpretation of isotachopherograms (191) 271

High-temperature gas chromatography
High-temperature quantitative glass capillary GC. Analysis of piperine and of quinine-quinidine (199) 105

Hippuric acid
GC determination of low concentrations of benzoic acid in human plasma and urine (181) 161

Hippuric acid
HPLC determination of hippuric acid in human urine. Preliminary results for normal urine levels (182) 88

Histamine
Quantitative analysis of histamine in biological samples by GC-MS (181) 153

Histidine
Presence of sodium azide during acid hydrolysis of protein samples causes the destruction of tyrosine, phenylalanine and histidine (189) 106

Histidine
GC separation of 3-methylhistidine (196) 166

Histidine diastereomers
Resolution of histidine diastereomers by GC (188) 408

Histones
Dissolvable disulphide-polyacrylamide gels for the electrophoretic analysis of chromosomal proteins and for affino-electrophoresis of thiol-proteins (190) 437

Histones
Quantitative staining of submicrogram amounts of histone and high-mobility group proteins on sodium dodecylsulphate-polyacrylamide gels (200) 200

Homologous series
Influence of temperature on the retention behaviour of members of homologous series in reversed-phase HPLC (196) 1

Homovanillic acid
Determination of homovanillic, isohomovanillic and vanillylmandelic acids in human urine by means of glass capillary GLC with temperature-programmed electron-capture detection (183) 391

Homovanillic acid
Highly sensitive method for the quantitation of homovanillic acid in cerebrospinal fluid (199) 191

Horizontal flow-through coil planet centrifuge
New horizontal flow-through coil planet centrifuge for counter-current chromatography. I. Principle of design and analysis of acceleration (188) 33

Horizontal flow-through coil planet centrifuge
New horizontal flow-through coil planet centrifuge for counter-current chromatography. II. The apparatus and its partition capabilities (188) 43

Horizontal flow-through coil planet centrifuge
New horizontal flow-through coil planet centrifuge for counter-current chromatography. III. Separation and purification of dinitrophenyl amino acids and peptides (193) 37

Hormones
Combination of enzymatic and mass fragmentographic assays for the identification and measurement of [met^5]-enkephalin (190) 43

Hormones
HPLC of the reduction products of progesterone (190) 169

Hormones
Pregnenolone separation from cholesterol using Sephadex LH-20 minicolumns (190) 256

Hormones
HPLC of plant hormones. I. Separation of plant hormones of the indole type (191) 129

Hormones
Reversed-phase HPLC purification and characterization of the adrenocorticotropin/lipotropin precursor and its fragments (193) 291

Hormones
Quantitative HPLC determination of Δ^4-3-ketosteroids in adrenocortical extracts (193) 413

Hormones
Separation of vasopressin analogues by reversed-phase HPLC (193) 427

Hormones
Improved HPLC separation of decomposition products of prednisolone by adding sulphite to the mobile phase (193) 480

Hormones
Practical aspects of the preparation and chromatography of the trimethylsilyl ethers of ecdysteroids (194) 43

Hormones
Intermediate- and large-scale reversed-phase preparative HPLC on an axially compressed column: a facile, quantitative separation of 7α- and 7β-methyl-17β-acetoxy-3-oxoandrost-4-enes (194) 239

Hormones
Comparison of HPLC and GC for the analysis of ecdysteroids (194) 343

Hormones
Glass capillary column GC method for simultaneous quantitative determination of insect juvenile hormones at the picogram level: a comparitive study of various halogenated derivatives (196) 425

Hormones
Separation of metabolites of testosterone hydroxylation by HPLC (196) 494

Hormones
Reversed- and normal-phase HPLC of 18-hydroxylated steroids and their derivatives. Comparison of selectivity, efficiency and recovery from biological samples (198) 23

Hormones
High-performance steric exclusion chromatography of plant hormones (198) 57

Hormones
Separation of enantiomeric iodinated thyronines by LC of diastereomers (198) 471

Hormones
Separation of urinary steroids by GLC on packed and mixed-bed columns (200) 221

Hyaluronic acid
HPLC of hyaluronic acid and oligosaccharides produced by bovine testes hyaluronidase (187) 373

Hycanthone
Plasma and tissue concentrations of hycanthone in the mouse determined by reversed-phase HPLC (183) 372

Hydralazine
Selective HPLC assays for hydralazine and its metabolites in plasma of man (181) 427

Hydralazine
GC method for the simultaneous determination of hydralazine and its acetylated metabolite in serum using a nitrogen-selective detector (183) 159

Hydralazine
Determination of hydralazine metabolites: 4-hydrazino-phthalazin-1-one and N-acetylhydrazinophthalazin-1-one by GC and *s*-triazolo[3,4-*a*]phthalazine and phthalazinone by HPLC (183) 167

Hydralazine
Determination of hydralazine and its

acetylated metabolites in urine by GC and HPLC (187) 218

Hydralazine
Detection of some antihypertensive drugs and their metabolites in urine by TLC. Five commonly used beta blockers and hydralazine (187) 277

Hydralazine metabolites
Quantitative analysis of hydralazine pyruvic acid hydrazone, the major plasma metabolite of hydralazine (187) 171

Hydralazine pyruvate hydrazone
Stability of hydralazine pyruvate hydrazone (198) 150

Hydrazinophthalazinone
Determination of hydralazine metabolites: 4-hydrazino-phthalazin-1-one and N-acetylhydrazinophthalazin-1-one by GC and *s*-triazolo[3,4-*a*]phthalazine and phthalazinone by HPLC (183) 167

Hydrides
Survey of selected hydrides as doping agents for a hydrogen-atmosphere flame-ionization detector (195) 211

Hydrocarbons
Evaluation of a thermally stable hydrocarbon as a non-polar base stationary phase for use in Rohrschneider-McReynolds-type schemes (186) 89

Hydrocarbons
Glass capillary adsorption columns for GC (186) 183

Hydrocarbons
GLC-MS studies on extracts and distillates from Posidonomia shales (186) 843

Hydrocarbons
Application of head-space analysis to the study of volatile organic impurity concentrates. GC determination of aromatic hydrocarbons in atmospheric air (186) 851

Hydrocarbons
Applications of reversed-phase chromatography and nephelometric detection to analysis of non-polar mixtures at microgram level (187) 111

Hydrocarbons
Quantification of petroleum-type hydrocarbons in avian tissue (187) 153

Hydrocarbons
Determination of some molecular structure parameters for hydrocarbons from the retention volumes on graphitized carbon black (191) 169

Hydrocarbons
Preparative separation of C_9 and C_{10} monoaromatic hydrocarbon mixtures by HPLC on silica gel (191) 319

Hydrocarbons
Chromatography of hydrocarbons on molybdenum sulphide (192) 323

Hydrocarbons
Evaluation of macroreticular resins for the determination of low concentrations of chlorinated hydrocarbons in sea water and tap water (193) 357

Hydrocarbons
Influence of temperature on the retention behaviour of members of homologous series in reversed-phase HPLC (196) 1

Hydrocarbons
GC study of solution and adsorption of hydrocarbons on glycols. I. Diethylene glycol and triethylene glycol (197) 135

Hydrocarbons
Separation of C_6 cyclic hydrocarbons before the inert peak on microporous silica: a case of negative adsorption (199) 23

Hydrocarbons, gasoline-range
Automatic system for the high-resolution GC analysis of gasoline-range hydrocarbon mixtures (186) 207

Hydrocarbons, gasoline-range
Applications of surface-modified porous silicas to glass capillary column preparation (199) 97

Hydrocortisone
Simultaneous determination of betamethasone, betamethasone acetate and hydrocortisone in biological fluids using HPLC (183) 131

Hydrocortisone acetate
Analysis of hydrocortisone acetate ointments and creams by HPLC (198) 41

Hydrogen
Response of electron-capture detector to hydrogen, oxygen, nitrogen, carbon dioxide, nitric oxide and nitrous oxide (193) 142

Hydrogen chloride
Determination of trace amounts of hydrogen chloride by derivatization with epoxides and GC separation (186) 219

Hydrophobicity
Use of reversed-phase TLC to asses hydrophobicity in studies of quantitative structure-activity relationships for drugs (196) 391

Hydroxyacetophenones
Separation and identification of some hydroxyacetophenones and their derivatives by TLC (188) 289

Hydroxy acids
GC separation of chiral 2-hydroxy acids and 2-alkyl-substituted carboxylic acids (195) 292

Hydroxy acids
Structural requirements for enantioselectivity in GC of chiral *a*-hydroxy acids (200) 189

Hydroxyamino acids
Gas chromatography of trifluoroacetyl amino acid trimethylsilyl esters. II. Problems in preparing specified derivatives of hydroxyamino acids (in German) (188) 251

Hydroxyaminopyrrolidone
Assay of HA-966 in rat plasma by capillary GLC with nitrogen-selective detection (183) 185

Hydroxyatrazine
Continuous post-column ion-pair extraction detection of some basic organic compounds in normal-phase chromatography (185) 473

Hydroxybenzoic acids
Analysis of hydroxybenzoic and hydroxycinnamic acids in plant material. I. Sample preparation and TLC (195) 85

Hydroxybenzoic acids
Analysis of hydroxybenzoic and hydroxycinnamic acids in plant material. II. Determination by GLC (195) 95

Hydroxycinnamic acids
Analysis of hydroxybenzoic and hydroxycinnamic acids in plant material. I. Sample preparation and TLC (195) 85

Hydroxycinnamic acids
Analysis of hydroxybenzoic and hydroxycinnamic acids in plant material. II. Determination by GLC (195) 95

Hydroxycorticosterone
PC systems for the separation of aldosterone, 18-hydroxycorticosterone, 18-hydroxydeoxycorticosterone, corticosterone and deoxycorticosterone (198) 70

Hydroxydeoxycorticosterone
PC systems for the separation of aldosterone, 18-hydroxycorticosterone, 18-hydroxydeoxycorticosterone, corticosterone and deoxycorticosterone (198) 70

Hydroxydiketotetranorprostanedioic acid
Capillary column GLC-MS assay for 7α-hydroxy-5,11-diketotetranorprostane-1,16-dioic acid, the major human urinary metabolite of prostaglandins E_1 and E_2 (181) 85

Hydroxyethanethiol
GLC determination of 2-mercaptoethanol (200) 211

Hydroxyindoleacetic acid
HPLC determination of 5-hydroxyindole-3-acetic acid in urine using Sephadex G-10 for isolation (181) 456

Hydroxylamine
Method for the measurement of hydroxylamine in colonic fluid using derivatisation and GC (181) 449

Hydroxylated steroids
Reversed- and normal-phase HPLC of 18-hydroxylated steroids and their derivatives. Comparison of selectivity, efficiency and recovery from biological samples (198) 23

Hydroxylysine
Employment of GLC for the analysis of collagen amino acids in biopsy tissue (182) 155

Hydroxylysine
Chromatographic method for the measurement of hydroxylysine, hydroxylysine glycosides and 3-methylhistidine in human urine (187) 271

Hydroxymethotrexate
Reversed-phase HPLC analysis of methotrexate and 7-hydroxymethotrexate in serum (181) 478

(Hydroxymethoxymethylphenyl)-(methoxyphenylisopropyl)aminoethanol
Separation of the diastereoisomers of 1-(4-hydroxy-3-methoxymethylphenyl)-2-N-(*p*-methoxyphenylisopropyl)aminoethanol by HPLC (193) 483

Hydroxynomifensine
GLC determination of 4'-hydroxynomifensine in biological samples (190) 475

Hydroxynorleucine
Employment of GLC for the analysis of collagen amino acids in biopsy tissue (182) 155

Hydroxyphenobarbital
Microassay for primidone and its metabolites phenylethylmalondiamide, phenobarbital, and *p*-hydroxyphenobarbital in human serum, saliva, breast milk and tissues by GC-MS using selected ion monitoring (182) 71

Hydroxyphenylbutazone
Simultaneous determination of phenylbutazone and its metabolites in plasma and urine by HPLC (183) 331

Hydroxyproline
Chromatographic separation and quantification of type I and type III collagens (182) 94

Hydroxyproline
Employment of GLC for the analysis of collagen amino acids in biopsy tissue (182) 155

Hydroxyproline
Determination of specific urinary hydroxyproline-containing peptides in patients with lung cancer (183) 221

Hydroxyproline
Improved method for the preparation of 3-hydroxyproline, and some of its chromatographic properties (190) 457

Hydroxysteroids
Enzymatic color development of 3α-hydroxysteroids on thin-layer chromatograms for determination of excretion pattern of 3α-hydroxysteroids in patients with some adrenogenital syndrome (182) 430

Hydroxytryptamine
Trimethylsilylation of biogenic indoleamines. Practical kinetic and analytical aspects related to tryptamine and 5-hydroxytryptamine (186) 619

Hypnotic drugs
Determination of nitromethaqualone in blood by electron-capture-GC (182) 435

Hypnotic drugs
Determination of propiomazine and its N-demethyl metabolite in plasma by GC with alkali flame ionization detection (183) 229

Hypnotic drugs
Improved detectability of barbiturates in HPLC by pre-column labelling and UV detection (186) 535

Hypoglycemic drugs
Electron-capture GC of plasma sulphonylureas after extractive methylation (181) 17

Hypoglycin
Separation of hypoglycin A from leucine and other amino acids on Sephadex G-10 (193) 148

Hypothalamic oligopeptides
HPLC and field desorption MS of hypothalamic oligopeptides (195) 369

I

Ibuprofen
Quantitation of ibuprofen in biological fluids by GC-MS (181) 463

Image analyser
Use of the image analyser Optomax for the quantitative evaluation of antibiotics separated by gel electrophoresis and by TLC (195) 297

Imidazolidines
Separation of methylol derivatives of imidazolidines, urea and carbamates by LC (190) 226

Immunosuppressive agents
HPLC determination of cyclosporin A in human plasma and urine (182) 454

Imogolite
GC adsorption studies on synthetic imogolite (188) 97

Impregnants
Chromatographic behaviour of aliphatic amines on phenol-impregnated thin layers (196) 225

Impurities
GC quantitation of residual impurities in raw material matrixes (193) 207

Inclusion compounds
Inclusion compounds in chromatography (Review) (184) 347

Indanes
Preparative separation of C_9 and C_{10} monoaromatic hydrocarbon mixtures by HPLC on silica gel (191) 319

Indoleacetic acid
High-performance steric exclusion chromatography of plant hormones (198) 57

Indoleamines
Trimethylsilylation of biogenic indoleamines. Practical kinetic and analytical aspects related to tryptamine and 5-hydroxytryptamine (186) 619

Indoleamines
GC of phenolic amines, 3-methoxycatecholamines, indoleamines and related amines as their N,O-ethyloxycarbonyl derivatives (194) 399

Indole derivatives
HPLC of plant hormones. I. Separation of plant hormones of the indole type (191) 129

Indole metabolites
Determination of tryptophan and metabolites in rat brain and pineal tissue by reversed-phase HPLC with electrochemical detection (181) 187

Indomethacin
Estimation of serum indomethacin at therapeutic levels by means of TLC and spectrophotometry (181) 282

Indomethacin
Rapid determination of indomethacin and salicylic acid in serum by means of reversed-phase LC (181) 385

Indoprofen
HPLC determination of indoprofen in plasma and urine (182) 262

Inert solid dilution
Enhancement of GSC column performance by inert solid dilution (188) 79

Information content
Information content of multi-dimensional switching systems in GC (186) 129

Injection effect
Extra-column effects in HPLC. I. Theoretical study of the injection problem (185) 79

Injection system
Capillary GC injection system for large sample volumes (186) 197

Injection technique
Direct aqueous injection GC for the analysis of trace organics in water (186) 785

Injection technique
Syringe and column adsorption of tertiary amines in GC (196) 489

Injection techniques
Factors affecting the accuracy and precision of cold on-column injections in capillary GC (189) 109

Injection techniques
Should the septum part of vaporizing injectors be kept at lower temperatures? (198) 64

Injection techniques
Analysis of dinitrogen-nitrogen oxide mixtures employing direct vacuum line/gas chromatograph injection (200) 224

Inorganic anions
Anion chromatography with low-conductivity eluents. II (187) 35

Inorganic colloids
Size-exclusion HPLC of inorganic colloids (185) 273

Inorganic halides
Use of isothermal GC for the determination of the adsorption enthalpies and entropies of inorganic halides at high temperatures (187) 319

Inositol hexaphosphoric acid
Analysis of phytic acid by HPLC (197) 274

Insecticides
Resolution, sensitivity and selectivity of a HPLC post-column fluorometric labeling technique for determination of carbamate insecticides (185) 615

Insecticides
Improved clean-up procedure for the determination of small residues of carbophenothion in mice (190) 201

Insecticides
Analysis of pentachlorophenol in waste water using HPLC (190) 203

Insecticides
Analysis of aminocarb formulations by HPLC (190) 208

Insecticides
Determination of carbofuran residues in potato, onion and turnip: comparative analyses using GC-selected ion monitoring or HPLC (193) 213

Insecticides
Evaluation of macroreticular resins for the determination of low concentrations of chlorinated hydrocarbons in sea water and tap water (193) 357

Insecticides
Detection of aminocarb and its major metabolites by TLC (194) 100

Insecticides
Direct determination of propoxur in plant tissues by GC with an alkali flame-ionization detector (197) 263

Insecticides
Simplified method for the analysis of some carbamate insecticides in foliage, forest soil and fish tissue by direct GLC (200) 179

Insecticides
GLC analysis of aminocarb and seven derivatives (200) 228

Insect juvenile hormones
Glass capillary column GC method for simultaneous quantitative determination of insect juvenile hormones at the picogram level: a comparitive study of various halogenated derivatives (196) 425

Insect pheromones
GC detection and structure analysis of volatile pheromones in insects (186) 605

Insect pheromones
4-(*p*-methoxycinnamyloxy)-4'-methoxy-azobenzene: a nematic liquid crystal for the GLC analysis of the stereochemistry of lepidopterous sex pheromones and related unsaturated fatty alcohols and derivatives (190) 35

Insect venoms
Application of high-speed aqueous gel permeation chromatography to insect venoms (197) 278

Instrumental errors
Gradient elution in LC. XIII. Instrumental errors in gradient elution chromatography (192) 37

Insulin analogues
Gel chromatographic separation of insulin analogues in human serum (181) 248

Insulin analogues
Release of insulin analogues in man, stimulated with glucose. In vitro and in vivo studies (182) 419

Interaction studies
Free displacement electrophoresis (isotachophoresis): an absolute determination of the Kohlrausch functions and their use in interaction studies (194) 1

Interelectrode distance
Effects of dimension and density in electron capture detectors (195) 17

Intermedine
Preparative separation of the pyrrolizidine alkaloids, intermedine and lycopsamine, as their borate complexes (195) 379

Intermolecular interactions
Determination of the energy of intermolecular interactions in GC from retention indices (190) 1

Iodamide
Rapid determination of renal contrast media

in biological fluids by means of HPLC (182) 492

Iodopyracet
Rapid determination of renal contrast media in biological fluids by means of HPLC (182) 492

Iodothyronines
Separation of iodinated compounds of L-tyrosyl-L-tyrosine from iodothyronines by Biogel P-2 CC (188) 431

Iodothyronines
Separation of enantiomeric iodinated thyronines by LC of diastereomers (198) 471

Iodotyrosines
Simple and fast separation of the iodotyrosines by TLC (194) 270

Ion exchangers
Stability of ion exchangers against treatment with nitric acid and radiation. II. Change in sorption properties of Dowex A-1, Chelex 100 and Wof MC 50 after treatment with nitric acid (in German) (198) 257

Ioxynil
HPLC analysis of formulations of phenoxycarboxylic acids, dicamba, ioxynil and bromoxynil mixtures (in French) (190) 401

Iridoid glycosides
Application of droplet counter-current chromatography to the isolation of natural products (186) 529

Isoalkanes
Calculation of GC retention indices of isoalkanes based on a topological approach (188) 297

Isoascorbic acid
Application of HPLC to the separation of ascorbic acid from isoascorbic acid (196) 163

Isocyanates
Determination of airborne free monomeric aromatic and aliphatic isocyanates by HPLC (190) 175

Isoelectric focusing
New developments in isoelectric focusing (Review) (184) 415

Isofezolac
Determination of isofezolac in biological fluids by reversed-phase LCC (182) 369

Isoflavones
HPLC method for the analysis of isoflavones (198) 193

Isohomovanillic acid
Determination of homovanillic, isohomovanillic and vanillylmandelic acids in human urine by means of glass capillary GLC with temperature-programmed electron-capture detection (183) 391

Isomer separation
Analysis of *Z* and *E* isomers of urocanic acid by HPLC (183) 83

Isomer separation
Reversed-phase separation of optical isomers of Dns-amino acids and peptides using chiral metal chelate additives (185) 323

Isomer separation
LC resolution of enantiomers on normal-phase chiral amide-bonded silica gel. Retentions of optically active α-amino acid derivatives on N-acyl homologues of L-valylaminopropylsilanized silica phases (186) 543

Isomer separation
Resolution of histidine diastereomers by GC (188) 408

Isomer separation
Separation of a series of positional and geometrical isomers of olefinic aliphatic primary alcohols and acetates by capillary GC (189) 199

Isomer separation
4-(*p*-methoxycinnamyloxy)-4'-methoxy-azobenzene: a nematic liquid crystal for the GLC analysis of the stereochemistry of lepidopterous sex pheromones and related unsaturated fatty alcohols and derivatives (190) 35

Isomer separation
Improved separation of uroporphyrin isomers by HPLC (190) 221

Isomer separation
Use of Carbowax 20M glass capillary columns below their lower temperature limit (190) 283

Isomer separation
Determination of *cis*- and *trans*-isomers of dosulepine and dithiadene by HPLC (191) 137

Isomer separation
Comparison of capillary columns coated with C_{87} hydrocarbon and squalane in the analysis of *n*-pentadecene isomers (191) 199

Isomer separation
Broad spectrum resolution of optical isomers using chiral HPLC bonded phases (192) 143

Isomer separation
Ligand-exchange chromatography. I. Resolution of L- and D-proline on a copper(II)-proline complex bound to microparticulate silica gel (192) 228

Isomer separation
TLC estimation of 2,4-xylenol in 2,5-xylenol (192) 250

Isomer separation
HPLC analysis of *cis-trans* stereoisomeric 3-dehydroretinals in the presence of retinal isomers (192) 395

Isomer separation
Resolution of ethylphenols following *in situ* reaction on a thin-layer plate with Fast Blue B Salt (192) 425

Isomer separation
Separation of xylenol isomers on nitrofluorenes and nitro-9-fluorenones by GLC. I. Mixed columns and effect of temperature on retention behaviour (193) 300

Isomer separation
Separation of enantiomers by HPLC on triacetylcellulose (193) 308

Isomer separation
Separation of isomeric ureas and oxazoles by TLC (193) 338

Isomer separation
Positive identification and quantitation of isomeric cocaines by HPLC (193) 371

Isomer separation
syn-anti Isomerisation of 2,4-dinitrophenylhydrazones of volatile carbonyl compounds in capillary GC-MS analyses (194) 135

Isomer separation
GLC separation of geometric isomers of unsaturated fatty acid methyl esters using a glass capillary column (194) 404

Isomer separation
Separation of underivatized dipeptides by HPLC on a weak anion-exchange bonded phase (195) 119

Isomer separation
Retention behaviour of *o*-, *m*- and *p*-isomers of benzene derivatives on a silica gel hydroxylated surface in LC (195) 205

Isomer separation
GC separation of chiral 2-hydroxy acids and 2-alkyl-substituted carboxylic acids (195) 292

Isomer separation
N-Docosanoyl-L-valine-2(2-methyl)-*n*-heptadecylamide as a stationary phase for the resolution of optical isomers in GLC (195) 317

Isomer separation
Separation behaviour of some isomeric organic compounds on sugars, sugar alcohols and their mixed phases by GLC (197) 251

Isomer separation
Coating support-coated open tubular capillary columns with Silar 10C or Alltech CS-10 and Silica T40 for separation of isomers of fatty acid methyl esters (198) 153

Isomer separation
Separation of enantiomeric iodinated thyronines by LC of diastereomers (198) 471

Isomer separation
Reversed-phase HPLC resolution of D- and L-Dns-amino acids by mixed chelate complexation (199) 295

Isoprenaline
Determination of isoprenaline by ion-pair HPLC (189) 272

Isopropoxyphenylmethyl carbamate
Direct determination of propoxur in plant tissues by GC with an alkali flame-ionization detector (197) 263

Isotachopherograms
High-speed isotachophoresis. Effect of driving current on the qualitative interpretation of isotachopherograms (191) 271

Isotherm linearity
Isotherm linearity and sample capacity in LC (193) 181

Isoxazolyl penicillins
Analysis of isoxazolyl penicillins and their metabolites in body fluids by HPLC (183) 339

J

JADO carbon
JADO carbon as an adsorbent in sampling columns (193) 137

Jasmolin
Examination of the conversion products of pyrethrins and allethrin formulations exposed to sunlight by GC and MS (198) 317

Juvenile hormones
Glass capillary column GC method for simultaneous quantitative determination of insect juvenile hormones at the picogram level: a comparitive study of various halogenated derivatives (196) 425

K

Kale anaemia factor
TLC of S-methylcysteine sulphoxide (kale anaemia factor) in the presence of common amino acids (189) 436

Katharometer
Effect of the sensitivity setting of a katharometer on response factors at low concentrations (197) 121

Kawa-lactones
Simple method for separation and quantitative determination of kawa-lactones by HPLC (in German) (193) 486

Ketamine
Capillary GC with a nitrogen detector for measurement of phencyclidine, ketamine and other arylcycloalkylamines in the picogram range (193) 157

Keto acids
Origin of the two peaks for 2-keto-3-methylvaleric acid produced by the oximation of the keto acids occurring in maple syrup urine disease (182) 221

Ketobemidone
Syringe and column adsorption of tertiary amines in GC (196) 489

Ketomethylvaleric acid
Origin of the two peaks for 2-keto-3-methylvaleric acid produced by the oximation of the keto acids occurring in maple syrup urine disease (182) 221

Ketones
GC determination of low-molecular-weight carbonyl compounds in aqueous solution as their O-(2,3,4,5,6-pentafluorobenzyl) oximes (187) 413

Ketones
Separation of saturated and unsaturated aldehyde and ketone 2,4-dinitrophenylhydrazone derivatives by reversed-phase HPLC (193) 432

Ketones, cyclic
HPLC of adamantanones and other cyclic ketones (200) 238

Ketoprofen
Convenient and sensitive HPLC assay for ketoprofen, naproxen and other allied drugs in plasma or urine (190) 119

Ketosteroids
Quantitative HPLC determination of Δ^4-3-ketosteroids in adrenocortical extracts (193) 413

Kinetics parameters
Chromatographic peak shape. III. Influence of kinetics parameters on solute profiles (186) 117

Kohlrausch functions
Free displacement electrophoresis (isotachophoresis): an absolute determination of the Kohlrausch functions and their use in interaction studies (194) 1

Kováts retention indices
Determination of Kováts retention indices with a capillary column and electron-capture detection: application to the assay of the enzymatic conversion of 3,4-epoxy-1-butene into diepoxybutane (186) 81

L

Labile complexes
Isotachophoresis of kinetically labile complexes (199) 81

Lactam antibiotics
Fast and simple method for the separation of intermediates and cofactors involved in the biosynthesis of cephalosporin C using chemically bonded C_{12} reversed-phase TLC (193) 511

Lactate dehydrogenase
Affinity chromatography of rat liver lactate dehydrogenase on the Remazol derivative of bead cellulose (194) 95

Lactate dehydrogenase
Determination of the interaction of lactate dehydrogenase with high-molecular-weight derivatives of AMP by affinity electrophoresis (194) 183

Lactate dehydrogenase isoenzymes
Study of the interaction between L-lactate dehydrogenase isoenzymes and immobilized 8-substituted adenosine 5'-monophosphate by means of affinity electrophoresis (192) 351

Lactic acid
TLC of citric acid cycle intermediates, lactic acid, and the amino acid taurine (189) 286

Lactones
Simple method for separation and quantitative determination of kawa-lactones by HPLC (in German) (193) 486

Lantadene
TLC of lantadene A and some related triterpenoids (196) 515

Layer preparation
New spreader for the preparation of gradient layers in TLC (189) 293

Layer thickness
Empirical correlation of layer thickness and support pore size in "chemically bonded" phases (195) 25

Lecithin
Miniature two-dimensional TLC separation of lecithin and sphingomyelin (182) 311

Lecithins
Quantitative determination of natural dipalmitoyl lecithin with dimyristoyl lecithin as internal standard by capillary GLC (192) 185

Lectins
Affinity chromatography of cells and cell membranes (Review) (184) 471

Lectins
Studies on lectins. XLV. Effect of ionic

strength on the interaction of lectins with carbohydrates as determined by affinity electrophoresis (192) 408

Lectins
Interactions of insolubilized lectins with membrane glycoproteins in presence of detergents (195) 197

Lenacil
GC determination of Lenacil and Pyrazon in soil and bacterial cultures (189) 89

Leucine
Determination of leucine flux in vivo by GC-MS utilizing stable isotopes for trace and internal standard (183) 403

Leucine
Separation of hypoglycin A from leucine and other amino acids on Sephadex G-10 (193) 148

Lidocaine
Qualitative differentiation between cocaine, lidocaine and cocaine-lidocaine mixtures ("rock-cocaine") using TLC (198) 534

Ligand-macromolecule binding parameters
Determination of ligand-macromolecule binding parameters and the method of Hummel and Dreyer (193) 520

Ligands
Covalent coupling of cholic acid to aminohexylamino-Sepharose 4B and its use in affinity chromatography of serum albumin (187) 409

Light-oil fraction
GC study of methylstyrenes in the light-oil fraction of coal distillate using selective hydrogenation (193) 397

Lipids
Lipids: TLC separation in twelve fractions by three successive unidirectional developments on the same plate (181) 254

Lipids
Lipid content of human platelets quantitated by TLC in combination with flame ionization detection (182) 301

Lipids
Miniature two-dimensional TLC separation of lecithin and sphingomyelin (182) 311

Lipids
GC-MS analysis of neutral lipids from methanogenic and thermoacidophilic bacteria (186) 795

Lipids
Investigation of the use of argentation HPLC for the analysis of triglycerides (188) 205

Lipids
Background contamination by phthalates commonly encountered in the chromatographic analysis of lipid samples (189) 421

Lipids
Investigation of vinca alkaloid-plasma membrane interactions by detergent gel chromatography (191) 287

Lipids
Quantitative determination of natural dipalmitoyl lecithin with dimyristoyl lecithin as internal standard by capillary GLC (192) 185

Lipids
Effect of solvents on the resolution of neutral lipids on chromarods (198) 279

Lipophilic ion-pairing reagents
Perfluoroalkanoic acids as lipophilic ion-pairing reagents in reversed-phase LC of peptides including secretin (195) 181

Lipophilicity
R_M values from reversed-phase TLC as parameters of lipophilicity in quantitative structure-activity relationships in four series of arylaliphatic acids (195) 329

Lipopolysaccharides
Separation of amino sugars and related compounds by two-dimensional TLC (195) 127

Lipoproteins
High-speed steric exclusion chromatography of biopolymers (185) 453

Lipotropin precursor
Reversed-phase HPLC purification and characterization of the adrenocorticotropin/lipotropin precursor and its fragments (193) 291

Liquid crystals
Utility of high-temperature thermotropic liquid crystals as stationary phases for novel GLC separations (186) 237

Liquid crystals
Effect of the support surface on the structure of the film of liquid crystalline stationary phase (188) 107

Liquid crystals
4-(*p*-methoxycinnamyloxy)-4'-methoxy-azobenzene: a nematic liquid crystal for the GLC analysis of the stereochemistry of lepidopterous sex pheromones and related unsaturated fatty alcohols and derivatives (190) 35

Liquid crystals
Liquid crystals as the stationary phase in GC. Adsorption phenomena caused by an electric field applied across the column (192) 89

Liquid crystals
Solution behaviour of liquid crystal phases in capillary GC. II. Solution properties of smectic A-, B- and C-modifications (in German) (192) 97

Liquid crystals
GLC separation of bile acids and steroids on a nematic liquid crystal (193) 444

Liquid crystals
Liquid crystals. I. Synthesis and application as stationary phases in GLC (195) 309

Liquid crystals
Properties of liquid crystalline cyanoazoxybenzene alkyl carbonates as stationary phases in GC (200) 65

Low-conductivity eluents
Anion chromatography with low-conductivity eluents (186) 509

Low-conductivity eluents
Anion chromatography with low-conductivity eluents. II (187) 35

Lycopsamine
Preparative separation of the pyrrolizidine alkaloids, intermedine and lycopsamine, as their borate complexes (195) 379

Lymphocytes
Surface affinity chromatographic separation of blood cells. I. Separation of human and rabbit peripheral granulocytes, lymphocytes and erythrocytes using polyethylene glycol-bonded column packings (187) 351

Lysine
Employment of GLC for the analysis of collagen amino acids in biopsy tissue (182) 155

Lysophosphatidylcholine
Comparative determination of plasma phospholipids by automated GLC and manual colorimetric phosphorus methods (182) 1

Lysophosphoglycerides
Isocratic HPLC separation of phosphoglycerides and lysophosphoglycerides (197) 79

Lysosomes
Separation of rat thyroid lysosome subpopulations on Sepharose 2B (189) 101

M

Macrolide antibiotics
Improved HPLC method for the comparison of heptaene macrolide antibiotics (189) 249

Macrolide antibiotics
TLC of macrolide antibiotics (193) 119

Macromolecules
Application of a graphite furnace atomic absorption detector automatically coupled to a high-performance liquid chromatograph for speciation of metal-containing macromolecules (185) 563

Macromolecules
Size and charge distribution of macromolecules in living systems (193) 1

Magnesium complexes
HPLC of nucleotide-Mg(II) complexes. Separation and mechanism of separation (199) 283

Magnesium dihexylphosphinate
Coordination polymers as adsorbents and stationary phases in GC. VI. Separation properties of alkaline-earth metal di-*n*-hexylphosphinates (188) 323

Maridomycin
Analytical studies of maridomycin. III. Spectrophotometric determination of maridomycins and 9-propionylmaridomycins as iodine complexes after TLC (190) 493

Mass transfer
Mass transfer in ideal and geometrically deformed open tubes. III. Deformed metal and plastic tubes (199) 3

Matrix material
Affinity chromatography of cells and cell membranes (Review) (184) 471

Mean pore diameter
Simplified HPLC phase characterization: particle size, surface area, mean pore diameter, amount of bonded phase and activity (186) 435

Mebendazole
Two HPLC determinations for mebendazole and its metabolites in human plasma using a rapid Sep Pak C18 extraction (183) 311

Mecillinam
Determination of mecillinam in urine by reversed-phase HPLC (181) 272

Mefruside
Quantitative GC determination of two oxidized metabolites of the diuretic mefruside in human urine, plasma and red blood cells (182) 179

Melanotropin inhibiting factor
Separation of alkylaminonaphthylenesulfonyl peptides and amino acids by HPLC. Methods for measuring melanotropin inhibiting factor breakdown (192) 341

Melatonin
Determination of melatonin by HPLC with electrochemical detection (190) 217

Menaquinones
HPLC assay of vitamin K in human serum (186) 749

Mephenytoin
A quantitative method for the simultaneous determination of carbamazepine, mephenytoin,

phenylethylmalonamide, phenobarbital, phenytoin and primidone in serum by thin-layer chromatography. Additional comments to the analysis of carbamazepine-10,11-epoxide (183) 387

Mepindolol
Determination of plasma mepindolol levels by HPLC and electrochemical detection (181) 67

Mercaptoethanol
GLC determination of 2-mercaptoethanol (200) 211

Mercaptopurine
Stability-indicating assay for azathioprine and 6-mercaptopurine by reversed-phase HPLC (186) 691

Mercaptopurines
Approach to the quantitative analysis of nucleotides by GC-MS (182) 395

Mescaline
Cactus alkaloids. XL. Identification of mescaline and other β-phenethylamines in *Pereskia, Pereskiopsis* and *Islaya* by use of fluorescamine conjugates (189) 79

Metabolic disorders
Diagnosis of neural crest tumors by reversed-phase HPLC determination of urinary catcholamine metabolites (186) 733

Metabolic profiles
Metabolic profiles of penaeid shrimp: dietary lipids and ovarian maturation (195) 359

Metalaxyl
GC determination of Metalaxyl (Ridomil) residues in tobacco (200) 313

Metalaxyl
Estimation of the systemic fungicide Ridomil by TLC (200) 317

Metal chelate affinity chromatography
Metal chelate affinity chromatography. I. Influence of various parameters on the retention of nucleotides and related compounds (198) 247

Metal complexes
Quantitative analysis of zinc, copper and nickel diethyldithiocarbamates by GLC (187) 363

Metal complexes
Binding of zinc and copper to some gel filtration media (188) 405

Metal complexes
Minimizing the retention of chromium chelates with some fluorinated β-diketones (191) 231

Metal complexes
GC of volatile metal dialkyldithiophosphates (193) 53

Metal complexes
Effect of the composition and structure of cobalt(III) complexes on their R_F values obtained by TLC on silica gel (196) 59

Metal complexes
GC with plasma emission spectroscopic detection of Friedel-Crafts catalyzed alkyl group redistribution products among Si, Ge, Sn and Pb atoms (196) 265

Metal complexes
Paper electrophoretic study of ion-pair formation. XIII. Behaviour of anionic trisoxalatocobalt(III) and trisoxalatochromium(III) in optically active electrolytes (198) 76

Metal complexes
Resolution of neutral complexes of transition metals by stereoselective adsorption on optically active complexes. I. Partial resolution of tris(acetylacetonato)cobalt(III) and tris(acetylacetonato)chromium(III) on cobalt(III) complexes (198) 172

Metal complexes
Isotachophoresis of kinetically labile complexes (199) 81

Metal complexes
HPLC of nucleotide-Mg(II) complexes. Separation and mechanism of separation (199) 283

Metal complexes
Reversed-phase HPLC resolution of D- and L-Dns-amino acids by mixed chelate complexation (199) 295

Metal compounds
Applications of Dexsil and other GC porous-layer open tubular columns to metal compound separations and pyrolysis (196) 403

Metal-containing surfaces
Synthesis and GC use of metal-containing surfaces (189) 13

Metal hexacyanoferrate compounds
Gel filtration of soluble coloured metal hexacyanoferrate(II) compounds (187) 458

Metal ions
Chromatographic separation of metal ions on macroreticular anion-exchange resins of a low capacity (188) 391

Metal oxides
LC on metal oxides with silica geometry (189) 139

Metal-solute complexes
HPLC of nucleotides and nucleosides using outersphere and innersphere metal-solute complexes (185) 361

[met]-enkephalin
Combination of enzymatic and mass fragmentographic assays for the identification and measurement of [met^5]-enkephalin (190) 43

Methacrylic acid esters
Modification of the Rohrschneider method for

the classification of the polarity of polymeric sorbents based on methacrylic acid esters (190) 13

Methampetamine
Microdetermination of stimulant drugs in urine by HPLC (196) 334

Methane thiolsulphonates
GC and GC-MS characterisation of methane thiolsulphonates carrying further functional groups (198) 287

Methapyrilene
Plasma and urine concentration of methapyrilene by nitrogen-phosphorus GLC (183) 234

Methaqualone
Mass screening and confirmation of methaqualone and its metabolites in urine by radioimmunoassay-TLC (190) 129

Methazole
Determination of residues of methazole and its metabolites, 1-(3,4-dichlorophenyl)-3-methylurea and 1-(3,4-dichlorophenyl)urea in soil by HPLC (197) 267

Methenamine
Determination of methenamine in biological samples by GLC (181) 11

Methimazole
Determination of methimazole in plasma using GC-MS after extractive alkylation (182) 63

Methionine
The isolation of methionine and ethionine by silver ligand chromatography and application to methionine-containing peptides (200) 185

Methionine-enkephalin
Ligand-exchange chromatography of small peptides on copper(II)-modified silica gel: application to the study of the enzymatic degradation of methionine-enkephalin (186) 677

Methomyl
Simplified method for the analysis of some carbamate insecticides in foliage, forest soil and fish tissue by direct GLC (200) 179

Methotrexate
Analysis of methotrexate by isotachophoresis (181) 441

Methotrexate
Reversed-phase HPLC analysis of methotrexate and 7-hydroxymethotrexate in serum (181) 478

Methoxycatecholamines
GC of phenolic amines, 3-methoxycatecholamines, indoleamines and related amines as their N,O-ethyloxycarbonyl derivatives (194) 399

Methoxycinnamyloxymethoxyazobenzene
4-(*p*-methoxycinnamyloxy)-4'-methoxyazobenzene: a nematic liquid crystal for the GLC analysis of the stereochemistry of lepidopterous sex pheromones and related unsaturated fatty alcohols and derivatives (190) 35

Methoxyhydroxyphenylglycol
Modification of assays for the routine analysis of 3-methoxy-4-hydroxyphenylglycol in urine by electron-capture GC (181) 1

Methylacetoxyoxoandrostenes
Intermediate- and large-scale reversed-phase preparative HPLC on an axially compressed column: a facile, quantitative separation of 7α- and 7β-methyl-17β-acetoxy-3-oxoandrost-4-enes (194) 239

Methylarsonic acid
Graphite furnace atomic absorption spectrophotometers as automated element-specific detectors for HPLC. The determination of arsenite, arsenate, methylarsonic acid and dimethylarsinic acid (191) 31

Methylchlorocatechol
TLC of 2-methyl-4-chlorophenoxyacetic acid and its soil metabolites (189) 73

Methylchlorophenoxyacetic acid
TLC of 2-methyl-4-chlorophenoxyacetic acid and its soil metabolites (189) 73

Methylcysteine
The enantiomer as internal standard for the quantitation of the alkylated amino acid S-methyl-L-cysteine in haemoglobin by GC-chemical ionisation MS with single ion detection (200) 145

Methylcysteine sulphoxide
TLC of S-methylcysteine sulphoxide (kale anaemia factor) in the presence of common amino acids (189) 436

Methylene number
Correlation of GC specific retention volumes of homologous compounds with temperature and methylene number (199) 209

Methylene Units
Concentration-dependent changes in Methylene Units in steroid analysis (in German) (197) 232

Methylglyoxal bis(guanylhydrazone)
HPLC analysis of the anticancer agent methylglyoxal bis(guanylhydrazone) (MGBG, NSC-32946) in biological fluids (183) 363

Methylglyoxal bis(guanylhydrazone)
HPLC assay for methylglyoxal bis(guanylhydrazone) (methyl GAG) in plasma and urine (187) 101

Methylhexadecanoic
Isolation of 14-methylhexadecanoic acid from wool fat by preparative GLC (193) 153

Methylhistidine
Rapid HPLC method for quantitation of 3-methylhistidine (182) 414

Methylhistidine
Chromatographic method for the measurement of hydroxylysine, hydroxylysine glycosides and 3-methylhistidine in human urine (187) 271

Methylhistidine
GC separation of 3-methylhistidine (196) 166

Methylmalonic acid
Simple estimation of urinary methylmalonic acid by isotachophoretic analysis (190) 501

Methyloctadecanoic acid
Detection of tuberculostearic acid in biological specimens by means of glass capillary GC-electron and chemical ionization MS, utilizing selected ion monitoring (182) 402

Methyl rhamnopyranoside
Preparative LCC of methyl ethers of methyl α-L-rhamnopyranoside (196) 331

Methyl silicone gums
Glass capillary columns coated with non-soluble methyl silicone gums (186) 159

Methylstyrenes
GC study of methylstyrenes in the light-oil fraction of coal distillate using selective hydrogenation (193) 397

Methyltetrahydrofolic acid
Determination of 5-methyltetrahydrofolic acid in plasma and spinal fluid by HPLC, using on-column concentration and electrochemical detection (182) 35

Methyltriazolophthalazine
GC method for the simultaneous determination of hydralazine and its acetylated metabolite in serum using a nitrogen-selective detector (183) 159

Metoclopramide
Electron-capture GC assay for metoclopramide in plasma (183) 175

Metoprolol
Determination of metoprolol metabolites in plasma and urine by electron-capture GLC (183) 92

Metyrapone test
Rapid quantitative assay of plasma 11-deoxycortisol and cortisol by HPLC for use in the metyrapone test (183) 87

Mexacarbate
Simplified method for the analysis of some carbamate insecticides in foliage, forest soil and fish tissue by direct GLC (200) 179

Mexiletine
Mexiletine analysis in blood and plasma using GC and nitrogen-selective detection (181) 469

Mianserin
Simplified method for determination of the tetracyclic antidepressant mianserin in human plasma using GC with nitrogen detection (181) 25

Mianserin
Comparative statistical study of assay methods using mass fragmentography and GC with nitrogen detection for determination of the tetracyclic antidepressant mianserin in human plasma (181) 115

Microbore columns
Use of microbore columns for the separation of substances of biological origin (185) 27

Microbore columns
Use of microbore columns for rapid LC separations (186) 475

Microbore columns
Design and use of short microbore columns in LC (198) 93

Midazolam
Determination of midazolam in serum by GC with a nitrogen-selective detector (182) 439

Mixed solvents
Study of mixed solvents: squalane-stearonitrile. I. Comparison of results by GC and surface tensions (in French) (188) 315

Mixed stationary phases
Separation behaviour of some isomeric organic compounds on sugars, sugar alcohols and their mixed phases by GLC (197) 251

Mobile phase composition
Reversed-phase ion-pair HPLC of drugs and related compounds using underivatized silica as the stationary phase (186) 705

Mobile phase composition
Influence of mobile phase composition and nature of sample on retention data from HPLC (191) 81

Mobile phase composition
Formulation of multicomponent mobile solvents for LC (198) 489

Mobile phase composition
Role of organic modifier sorption on retention phenomena in reversed-phase LC (199) 259

Mobile phase effect
Mechanism of solute retention in liquid-solid chromatography and the role of the mobile phase in affecting separation. Competition *versus* "sorption" (Review) (184) 363

Mobile phase effects
Mobile phase effects in reversed-phase chromatography. I. Concomitant dependence of retention on column temperature and eluent composition (185) 99

Mobile phase effects
Mobile phase effects in reversed-phase

chromatography. II. Acidic amine phosphate buffers as eluents (185) 111

Mobile phase effects
Mobile phase effects in reversed-phase chromatography. III. Changes in conformation and retention of oligo(ethylene glycol) derivatives with temperature and eluent composition (185) 129

Mobile phases
Reversed-phase ion-pair chromatography with an adsorbing stationary phase and a hydrophobic quaternary ammonium ion in the mobile phase. I. Retention studies with tetrabutylammonium as cationic component (185) 225

Mobile phase segmentation
Coated-open-tubular chromatography with flow segmentation. I. Theory (185) 43

Mobile phase segmentation
Coated-open-tubular chromatography with flow segmentation. II. Experimental study and optimization for size-exclusion separation (185) 57

Molecular interactions
Study of the interactions between substituted phenols and alcohols and dibenzyl sulphoxide by means of GC (198) 495

Molecular retention index
Elimination of the reference stationary phase in classification schemes using the molecular retention index (199) 199

Molecular sieve effects
Theoretical considerations of molecular sieve effects (187) 285

Molecular size parameters
Empirical approach to the universal calibration of isomers of phthalate esters and similar compounds in size exclusion chromatography (192) 295

Molecular structure
Determination of some molecular structure parameters for hydrocarbons from the retention volumes on graphitized carbon black (191) 169

Molecular weight calibration
Fractionation of polymers by gel permeation chromatography: an experimental and theoretical approach (192) 259

Molecular-weight distribution
High-performance gel permeation chromatography of water-soluble cellulosics (192) 275

Molecular-weight distribution
Determination of molecular-weight distribution of aorta glycosaminoglycans by automated gel filtration (192) 307

Molecular weight distributions
Chromatographic behaviour of aromatic compounds on Sephadex LH gels. Calibration of gel columns for determination of molecular weight distributions (198) 131

Molecular weight distributions
Gel chromatographic comparison of the molecular weight distributions of Ampholine, Servalyte and Pharmalyte carrier ampholytes used in isoelectric focusing (198) 301

Molluscicides
Extractive alkylation of 5,2'-dichloro-4-nitrosalicylanilide (niclosamide) for GLC analysis (189) 375

Molybdenum sulphide
Chromatography of hydrocarbons on molybdenum sulphide (192) 323

Monoacetyldapsone
Rapid HPLC method for the determination of dapsone and monoacetyldapsone in biological fluids (182) 130

Monoamines
Quantitative structure-retention relationships in cation-exchange chromatography (198) 241

Monocarboxylic acids
Determination of free short-chain monocarboxylic acids from naturally occurring esters by GLSC on micropacked columns (in German) (189) 43

Mono(ethylhexyl) phthalate
Determination by electron-capture GC of mono- and di(2-ethylhexyl) phthalate in intravenous solutions stored in poly(vinyl chloride) bags (193) 405

Monoglycerides
HPLC of diglyceride *p*-nitrobenzoates. An approach to molecular analysis of phospholipids (198) 520

Monomethylolurea
Separation of urea-formaldehyde addition products by gel permeation chromatography (195) 154

Monosaccharides
Separation of monosaccharides by HPLC: comparison of UV and RI detection (189) 414

Monosaccharides
Separation of monosaccharides as their alditol acetates by capillary column GLC (194) 410

Monoterpenes
Study of three types of essential oil of *Valeriana officinalis* L. s. l by combined GC-negative ion chemical ionization MS (190) 321

Monothiols
Derivatization of dithiols and certain monothiols with phenylarsine oxide for GC (200) 171

Morphine
Amperometric HPLC method for narcotic alkaloids (188) 420

Mucin
Differentiation of the polysaccharide side-chains of glycoproteins by a fingerprinting technique: heterogeneity of human gastric mucin (181) 108

Mucolytic drugs
Determination of a new mucolytic drug Adamexina in biological liquids by photodensitometry (181) 504

Mucopolysaccharides
Fractionation and identification of heparin and other acidic mucopolysaccharides by a new discontinuous electrophoretic method (196) 455

Multi-dimensional chromatography
Two-dimensional (Letter to the Editor) (187) 466

Multifactor optimization
Multifactor optimization of reversed-phase LC separations (199) 317

Multilayer adsorption
Solute-solvent interactions on the surface of silica gel. III. Multilayer adsorption of water on the surface of silica gel (196) 193

Multivitamin tablets
Comparison of reversed-phase and adsorption modes of HPLC for the assay of fat soluble vitamins in multivitamin tablets (200) 300

Mycolic acids
TLC of methanolysates of mycolic acid-containing bacteria (188) 221

Mycotoxins
HPLC analysis of the mycotoxin citrinin and its application to biological fluids (190) 419

Mycotoxins
Capillary GC-MS determination of some mycotoxins causing fusariotoxicoses in animals (191) 327

Mycotoxins
HPLC determination of major mycotoxins produced by *Alternaria* molds (194) 89

Mycotoxins
HPLC analysis of the mycotoxin penicillic acid and its application to biological fluids (194) 387

Mycotoxins
Separation and purification of metabolites of the mycotoxin, sporidesmin, by HPLC (200) 288

Myoglobin
Human oxymyoglobin: isolation and characterization (195) 277

Myoneural blocking agents
Determination of Org NC 45 (a myoneural blocking agent) in human plasma using normal-phase HPLC (183) 459

Myosins
HPLC of myosin's subfragment 1 and light chains on Spherogel TSK-type columns (197) 207

N

NADP-dependent dehydrogenase
Study of the interaction between NADP-dependent dehydrogenase and immobilized adenosine 2'-monophosphate by means of affinity electrophoresis (196) 85

Nalidixic acid
Determination of nalidixic acid and its two major metabolites in human plasma and urine by reversed-phase HPLC (181) 399

Naloxone
GC analysis of naloxone in biological fluids (183) 352

Naloxone hydrochloride
HPLC analysis of naloxone hydrochloride in injectable solutions (200) 277

Naphthalene derivatives
Qualitative and quantitative analysis of propylnaphthalenes by GC on silicone stationary phases (197) 92

Naphthalene derivatives
GC of some alkenyl- and cycloalkenylnaphthalenes on Bentone 34, silicone fluid DC 550, and Apiezon L phases (198) 1

Naphthol
Application of a direct aqueous acetylation technique to the GC quantitation of nitrophenols and 1-naphthol in environmental water samples (195) 105

Naproxen
Convenient and sensitive HPLC assay for ketoprofen, naproxen and other allied drugs in plasma or urine (190) 119

Narcotic alkaloids
Amperometric HPLC method for narcotic alkaloids (188) 420

Narcotic antagonists
HPLC analysis of naloxone hydrochloride in injectable solutions (200) 277

Natural products
Application of droplet counter-current chromatography to the isolation of natural products (186) 529

Nebramycin
Aminoglycoside antibiotics: TLC

bioautographic detection and quantitative assay (195) 251

Neoamygdalin
Improved GC of amygdalin and its diastereomer (197) 240

Neostigmine
Reversed-phase, ion-pair LC of quaternary ammonium compounds. Determination of pyridostigmine, neostigmine and edrophonium in biological fluids (183) 193

Nerve gases
Stimulated sequential dehydroxy-fluorination: a safe, convenient method for GC analysis of phosphate diesters with possible application to hydrolysates of organophosphorus pesticides and nerve gases (192) 173

Neurohypophyseal hormones
Separation of vasopressin analogues by reversed-phase HPLC (193) 427

Neuroleptic drugs
Specific HPLC method for estimation of the *cis*(*Z*)- and *trans*(*E*)-isomers of clopenthixol and a N-dealkyl metabolite (183) 239

Neuroleptic drugs
Reliable routine method for determination of perazine in serum by TLC with an internal standard (183) 475

Neuroleptic drugs
Separation and quantitation of bromperidol in pharmaceutical preparations by HPLC and HPTLC (193) 255

Neuroleptic drugs
HPLC separation of the N- and S-oxides of fluphenazine and fluphenazine decanoate (194) 416

Neuropeptides
Ion-pair HPLC of the insect neuropeptide proctolin and some analogs (194) 421

Nickel
Quantitative analysis of zinc, copper and nickel diethyldithiocarbamates by GLC (187) 363

Nickel
GC of volatile metal dialkyldithiophosphates (193) 53

Nickel catalysts
GC determination of adsorption-desorption rates. Interaction of hydrogen on a nickel catalyst (196) 355

Niclosamide
Extractive alkylation of 5,2'-dichloro-4-nitrosalicylanilide (niclosamide) for GLC analysis (189) 375

Ninhydrin
Use of ninhydrin as a spray reagent for the detection of some basic drugs on thin-layer chromatograms (195) 133

Nitramines
Response of nitramines in the thermal analyzer (187) 229

Nitrates, organic
Sintered TLC (Review) (184) 37

Nitric acid treatment
Stability of ion exchangers against treatment with nitric acid and radiation. II. Change in sorption properties of Dowex A-1, Chelex 100 and Wof MC 50 after treatment with nitric acid (in German) (198) 257

Nitric oxide
Response of electron-capture detector to hydrogen, oxygen, nitrogen, carbon dioxide, nitric oxide and nitrous oxide (193) 142

Nitrile group-solute interactions
GLC determination of equilibrium constants of interactions between solutes and free nitrile groups (in French) (193) 9

Nitrite
GC determination of nitrite in foods as trimethylsilyl derivative of 1*H*-benzotriazole (194) 21

Nitrobenzylarenes
TLC of nitrobenzylarenes (in German) (188) 448

Nitro compounds
Reversed-phase LC of aromatic nitro compounds on a macroporous polystyrene gel (197) 147

Nitrofluorenes
Separation of xylenol isomers on nitrofluorenes and nitro-9-fluorenones by GLC. I. Mixed columns and effect of temperature on retention behaviour (193) 300

Nitrofurans
HPLC analysis of furazolidone in liver and kidney (198) 526

Nitrofurantoin
HPLC assay for the metabolites of nitrofurantoin in plasma and urine (182) 246

Nitrogen
Response of electron-capture detector to hydrogen, oxygen, nitrogen, carbon dioxide, nitric oxide and nitrous oxide (193) 142

Nitrogen-containing compounds
Study of nitrogen-containing compounds in cigarette smoke by GC-MS (198) 188

Nitrogenous compounds
Continuous automatic nitrogen determination for gel chromatography of protein enzymatic hydrolysates (193) 496

Nitrogen oxide
Analysis of dinitrogen-nitrogen oxide mixtures employing direct vacuum line/gas chromatograph injection (200) 224

Nitroimidazoles
TLC of some 5-nitroimidazoles of pharmaceutical interest (193) 340

Nitromethaqualone
Determination of nitromethaqualone in blood by electron-capture-GC (182) 435

Nitrophenol
Measurement of the formation of paracetamol and *p*-nitrophenol glucuronides *in vitro*, by ion-pair HPLC (192) 247

Nitrophenols
Application of a direct aqueous acetylation technique to the GC quantitation of nitrophenols and 1-naphthol in environmental water samples (195) 105

Nitrophenols
Quantitative analysis of polynitrophenols in water in the micro- to nanogram range by reversed-phase ion-pair LC (195) 435

Nitrosoamino acids
Analysis of N-nitroso compounds. V. Trace analysis of N-nitrosoamino acids: Gas chromatography and spectroscopy of N-nitrosoamino acid methyl esters (in German) (193) 381

Nitrosoamino acids
GC determination of N-nitrosoamino acids (195) 416

Nitrosopipemidic acid
Estimation of pipemidic acid and N-nitrosopipemidic acid by HPLC (200) 245

Nitrous oxide
Response of electron-capture detector to hydrogen, oxygen, nitrogen, carbon dioxide, nitric oxide and nitrous oxide (193) 142

Nitroxylenes
Separation behaviour of some isomeric organic compounds on sugars, sugar alcohols and their mixed phases by GLC (197) 251

Nomifensine metabolites
GLC determination of 4'-hydroxynomifensine in biological samples (190) 475

Non-halogenated compounds
Quantitation of non-halogenated compounds by electron-capture GC (Review) (184) 313

Non-histone proteins
Quantitative staining of submicrogram amounts of histone and high-mobility group proteins on sodium dodecylsulphate-polyacrylamide gels (200) 200

Non-polar mixtures
Applications of reversed-phase chromatography and nephelometric detection to analysis of non-polar mixtures at microgram level (187) 111

Nonylic acid vanillylamide
Quantification of nonylic acid vanillylamide and other capsaicinoids in the pungent principle of Capsicum fruits and preparations by GLC on glass capillary columns (189) 389

Nonylic acid vanillylamide
Rapid determination of nonylic acid vanillylamide and other capsaicinoids in Capsicum fruits and extracts by means of Ag^+-complexation HPLC (in German) (193) 101

Noradrenaline
Simple method for isolating free urinary catecholamines by boric acid gel chromatography (183) 78

Noradrenaline
Semi-automated fluorimetric method for the estimation of urinary catecholamines using HPLC (183) 295

Noradrenaline
Electrochemical detector for HPLC (192) 135

Nordiazepam
HPLC and GLC determination of diazepam and nordiazepam in plasma (183) 441

Norepinephrine
Simplified LC-electrochemical determination of norepinephrine and dopamine in rat brain (181) 453

Norepinephrine
Simple and sensitive procedure for the assay of serotonin and catecholamines in brain by HPLC using fluorescence detection (182) 277

Norepinephrine
Investigation of the performance of a HPLC system with an electrochemical detector (183) 261

Norepinephrine
GC of epinephrine and norepinephrine after derivatization with chloroformates in aqueous media (190) 347

Norgestrel
HPLC and GLC of some norgestrel intermediates. Physical properties of the isolated *syn*- and *anti*-isomers of oximes (191) 145

Nortriptyline
Determination of amitriptyline-N-oxide, amitriptyline and nortriptyline in serum and plasma by HPLC (183) 321

Norverapamil
HPLC analysis of verapamil. II. Simultaneous quantitation of verapamil and its active metabolite, norverapamil (181) 484

Norzimelidine
Reversed-phase chromatography of zimelidine and similar dibasic amines. I. Analysis in biological material (185) 593

Nuclear magnetic resonance
On-line coupling of HPLC and NMR (186) 497

Nucleic acid bases
Nucleic acid bases and derivatives: detection by Dns derivatization TLC (195) 287

Nucleobases
Separation of halogenated uracil derivatives from nucleobases and nucleosides by TLC on silica gel (187) 453

Nucleoside phosphates
Rapid separation of nucleoside mono-, di- and triphosphates on ion-exclusion/exchange columns (192) 490

Nucleosides
Approach to the quantitative analysis of nucleotides by GC-MS (182) 395

Nucleosides
Quantitative HPLC of nucleosides and bases in human plasma (183) 421

Nucleosides
HPLC analysis of cytosine arabinoside and metabolites in biological samples (183) 449

Nucleosides
Identification of adenosine and eight modified adenine nucleosides using reversed-phase HPLC and enzymatic peak shift with adenosine deaminase (183) 499

Nucleosides
HPLC of nucleotides and nucleosides using outersphere and innersphere metal-solute complexes (185) 361

Nucleosides
Identification and quantitation of nucleosides, bases and other UV-absorbing compounds in serum, using reversed-phase HPLC. I. Chromatographic methodology (186) 647

Nucleosides
Identification and quantitation of nucleosides, bases and other UV-absorbing compounds in serum, using reversed-phase HPLC. II. Evaluation of human sera (186) 659

Nucleosides
Separation of halogenated uracil derivatives from nucleobases and nucleosides by TLC on silica gel (187) 453

Nucleosides
Chromatography of nucleosides (188) 129

Nucleosides
Rapid purification of radiolabeled trinucleoside diphosphates by reversed-phase HPLC (192) 485

Nucleosides
HPLC and MS of transfer RNA bases for isotopic abundance (194) 205

Nucleosides
Reversed-phase LC of 5-fluorouracil nucleosides and nucleotides in the presence of quaternary ammonium ions (194) 305

Nucleosides
GC retention data for silyl and acyl derivatives of nucleosides (196) 367

Nucleosides
High-performance liquid affinity chromatography of nucleosides, nucleotides and carbohydrates with boronic acid-substituted microparticulate silica (200) 254

Nucleotides
Simultaneous analysis of ATP, ADP, AMP, and other purines in human erythrocytes by HPLC (181) 103

Nucleotides
Approach to the quantitative analysis of nucleotides by GC-MS (182) 395

Nucleotides
HPLC of nucleotides and nucleosides using outersphere and innersphere metal-solute complexes (185) 361

Nucleotides
Determination of pyrimidine ribotide and deoxyribotide pools in cultured cells and mouse liver by HPLC (188) 149

Nucleotides
Separation and sequencing of the sequence isomers of pyrimidine deoxypentanucleoside tetraphosphates by HPLC (188) 273

Nucleotides
Simple anion-exchange chromatography for the determination of adenine nucleotides by using AG MP-1 resin (192) 193

Nucleotides
Reversed-phase HPLC of adenine nucleotides: application to the kinetics of an adenosine 3'-phosphate 5'-sulphatophosphate sulphotransferase from plants (193) 285

Nucleotides
Reversed-phase LC of 5-fluorouracil nucleosides and nucleotides in the presence of quaternary ammonium ions (194) 305

Nucleotides
Metal chelate affinity chromatography. I. Influence of various parameters on the retention of nucleotides and related compounds (198) 247

Nucleotides
Analysis of purine ribonucleotides and deoxyribonucleotides in cell extracts by HPLC (198) 435

Nucleotides
Operational parameters of anion-exchange chromatography using AG MP-1 resin for rapid assay of adenine nucleotides (198) 500

Nucleotides
HPLC of nucleotide-Mg(II) complexes. Separation and mechanism of separation (199) 283

Nucleotides
High-performance liquid affinity chromatography of nucleosides, nucleotides and carbohydrates with boronic acid-substituted microparticulate silica (200) 254

O

Octadecyl bonded silicas
GC evidence for phase transitions in very compact octadecyl bonded silicas (200) 95

Octadienes
Identification of *n*-hepta- and *n*-octadienes by high-resolution GC using structure-retention correlations and MS (191) 187

Octylamine hydrochloride
Combined effects of pH and surface-active-ion concentration in reversed-phase LC (199) 307

Oestrogenic drugs
Determination of residues of anabolic drugs in meat by GC-MS (195) 231

Oil vapour
The photoionization detector. Theory, performance and application as a low-level monitor of oil vapour (190) 263

Oligo(ethylene glycol) derivatives
Mobile phase effects in reversed-phase chromatography. III. Changes in conformation and retention of oligo(ethylene glycol) derivatives with temperature and eluent composition (185) 129

Oligogalacturonic acids
Separation of α-D-galacturonic acid oligomers by chromatography on polyacrylamide gel (194) 315

Oligomers
Determination of oligomers in vinyl chloride polymers by steric exclusion chromatography (193) 235

Oligomers
Separation and characterization of by-product oligomers in epoxy resins by reversed-phase HPLC (193) 243

Oligonucleotides
Application of a weakly basic dimethylamino-modified silica ion exchanger to the separation of oligonucleotides (185) 403

Oligonucleotides
Separation and sequencing of the sequence isomers of pyrimidine deoxypentanucleoside tetraphosphates by HPLC (188) 273

Oligopeptides
HPLC and field desorption MS of hypothalamic oligopeptides (195) 369

Oligosaccharides
HPLC of hyaluronic acid and oligosaccharides produced by bovine testes hyaluronidase (187) 373

Oligosaccharides
A rapid and high-resolution method to determine the composition of corn syrups by LC (187) 381

Oligosaccharides
LC of non-reducing oligosaccharides: a new detection principle (191) 313

Oligosaccharides
A homologous series of non-reducing oligosaccharides in *Artemisia absinthium* L. roots (200) 309

On-line trace enrichment
On-line trace enrichment in HPLC using a pre-column (185) 483

Open-tubular columns
Kinetic optimization of straight open-tubular LC (186) 405

Open tubular columns
Open tubular columns in LC (191) 17

Open tubular columns
Glass surface analytical studies in the preparation of open tubular columns for GC (199) 355

Open-tubular microcapillary columns
Studies on open-tubular microcapillary LC. VI. Styrene-divinylbenzene copolymer stationary phase (192) 127

Open tubular microcapillary columns
Open-tubular microcapillary LC with 30-40 μm I.D. columns (199) 249

Opiate alkaloids
Amperometric HPLC method for narcotic alkaloids (188) 420

Opium alkaloids
Separation and determination of opium alkaloids by HPLC (190) 251

Optical brighteners
Optical brighteners as TLC detection reagents for glycoalkaloids and steroid alkaloids in *Solanum* species. II. Blankophor BA 267%, BBU neu and KU, and Tinopal CBS-X and 5 BMS-X (189) 406

Optical brighteners
Simple rapid TLC method for the detection of optical brighteners in polymers (194) 434

Optical detectors
Specific detection in GLC effluents with on-line HPLC optical detectors (199) 113

Optimization
Coated-open-tubular chromatography with

flow segmentation. II. Experimental study and optimization for size-exclusion separation (185) 57

Optimization
Systematic approach to optimizing resolution in reversed-phase LC with emphasis on the role of temperature (185) 153

Optimization
Optimization of the instrumental parameters of a combined liquid chromatograph-mass spectrometer coupled by an interface for direct liquid introduction. I. Performance of the vacuum equipment (185) 529

Optimization
New approach to chromatographic optimization in glass capillary GC (186) 189

Optimization
Kinetic optimization of straight open-tubular LC (186) 405

Optimization
Influence of particle parameters on the efficiency of LC systems (191) 9

Optimization
Gradient elution in LC. XII. Optimization of conditions for gradient elution (192) 19

Optimization
Optimization of solvent strength and selectivity for reversed-phase LC using an interactive mixture-design statistical technique (199) 57

Optimization
Multifactor optimization of reversed-phase LC separations (199) 317

Organelles
New developments in isoelectric focusing (Review) (184) 415

Organelles
Affinity chromatography of cells and cell membranes (Review) (184) 471

Organic acids
Conductimetric detection of organic acids in reversed-phase chromatography (185) 625

Organic acids
Stevia rebaudiana. I. Use of *p*-bromophenacyl bromide to enhance UV detection of water-soluble organic acids (steviolbioside and rebaudioside B) in HPLC analysis (192) 387

Organic acids
Trimethylsilyl esters of plant acids and their application in GLC. Preparation, kinetics of silylation and solvent effects on quantitative reaction and stability of the derivatives (in German) (194) 33

Organic acids
Concentration and determination of organic acids in complex aqueous samples (199) 143

Organic compounds
Continuous post-column ion-pair extraction detection of some basic organic compounds in normal-phase chromatography (185) 473

Organic compounds
Analysis of non-volatile organic compounds in water by HPLC (185) 671

Organic compounds
Direct determination of organic compounds in water using steam-solid chromatography (191) 221

Organic compounds
JADO carbon as an adsorbent in sampling columns (193) 137

Organic compounds
Sampling and HPLC analysis of organic compounds in liquified carbon dioxide (194) 335

Organic impurities
Detection of trace organic impurities in binary solvent systems: a solvent purity test (188) 193

Organic modifiers
Influence of organic modifiers on the retention behaviour in reversed-phase LC and its consequences for gradient elution (185) 179

Organic modifier sorption
Role of organic modifier sorption on retention phenomena in reversed-phase LC (199) 259

Organic substances
TLC of slightly soluble substances at elevated temperature (186) 575

Organochlorine compounds
Carbon skeleton capillary GC (193) 437

Organochlorine pesticides
OV-17-QF-1 capillary column for organochlorine pesticide analysis (195) 265

Organogermanium compounds
GC detection of organogermanium compounds by surface or gas-phase luminescence (186) 299

Organometallic compounds
Applications of Dexsil and other GC porous-layer open tubular columns to metal compound separations and pyrolysis (196) 403

Organophosphoric acids
Determination of organophosphoric and organophosphorothioic acids in aqueous solutions by ion chromatography (189) 61

Organophosphorothioic acids
Determination of organophosphoric and organophosphorothioic acids in aqueous solutions by ion chromatography (189) 61

Organophosphorus compounds
Structure-response relationship of GC-flame photometric detection to some organophosphorus compounds (189) 331

Organophosphorus pesticides
Stimulated sequential dehydroxy-fluorination:

a safe, convenient method for GC analysis of phosphate diesters with possible application to hydrolysates of organophosphorus pesticides and nerve gases (192) 173

Organosilicon bonded phases
"Bonded phases" from cyclic organosilicons for GC and LC (200) 3

Organotin compounds
Comparative assessment of GLC and HPLC for the separation of tin tetraalkyls and alkyltin halides (200) 305

Organotin silicates
Application of a graphite furnace atomic absorption detector automatically coupled to a high-performance liquid chromatograph for speciation of metal-containing macromolecules (185) 563

Org NC 45
Determination of Org NC 45 (a myoneural blocking agent) in human plasma using normal-phase HPLC (183) 459

Overpressured thin-layer chromatography
Resolution and retention behaviour of some dyes in overpressured TLC (191) 293

Ovomucoid
Purification of ovomucoid by hydroxyapatite chromatography (187) 224

Oxalate
Complexation with ferric ions: a potential source of error in the isotachophoretic determination of oxalate (188) 266

Oxalic acid
Decreased accuracy in isotachophoretic analysis due to septum bleed (188) 262

Oxalic acid
Determination of multifunctional carboxylic acids in the presence of iron by GC (190) 188

Oxazoles
Separation of isomeric ureas and oxazoles by TLC (193) 338

Oxepinac
Quantitative analysis of oxepinac in human plasma, urine and saliva by GC-mass fragmentography (182) 47

Oxipurinol
Improvements to a HPLC assay for allopurinol and oxipurinol in plasma (181) 286

Oxpentifylline
Determination of oxpentifylline and a metabolite, 1-(5'-hydroxyhexyl)-3,7-dimethylxanthine, by GLC using a nitrogen-selective detector (181) 355

Oxprenolol
Quantification of oxprenolol in biological fluids using HPLC (181) 135

Oxygen
Response of electron-capture detector to hydrogen, oxygen, nitrogen, carbon dioxide, nitric oxide and nitrous oxide (193) 142

Oxymyoglobin
Human oxymyoglobin: isolation and characterization (195) 277

Oxyphenbutazone
Simultaneous determination of phenylbutazone and its metabolites in plasma and urine by HPLC (183) 331

Oxytetracycline
Reversed-phase ion pair chromatography of oxytetracycline, epioxytetracycline and anhydrooxytetracycline (190) 486

P

Packings
Isocratic multi-column HPLC as a technique for qualitative analysis and its application to the characterisation of basic drugs using an aqueous methanol solvent (187) 65

Packings
Support treated with Carbowax for a standard, non-polar packing in GLC (194) 145

Pairing ions
Functional group behaviour in ion-pair reversed-phase HPLC using surface-active pairing ions (185) 197

Palladium
GC of volatile metal dialkyldithiophosphates (193) 53

Papaverine
Determination of papaverine in plasma and urine by HPLC (182) 482

Paracetamol
Measurement of the formation of paracetamol and *p*-nitrophenol glucuronides *in vitro,* by ion-pair HPLC (192) 247

Parathion
Application of a direct aqueous acetylation technique to the GC quantitation of nitrophenols and 1-naphthol in environmental water samples (195) 105

Particle parameters
Influence of particle parameters on the efficiency of LC systems (191) 9

Particle size
Simplified HPLC phase characterization: particle size, surface area, mean pore diameter, amount of bonded phase and activity (186) 435

Partition behaviour
Anomalous partition behaviour of sodium dodecyl sulphate monomer in Sephadex gels with high dextran concentrations. Possible role of water of hydration in the gel phase (193) 345

Partition coefficients
Effect of adsorption on solute retention in GLC (186) 47

Partition coefficients
Evaluation of octanol-water partition coefficients by using HPLC (192) 315

Pattern separation
Numerical analysis of chromatographic-spectrometric data. Purity of partially resolved peaks and pattern separation of the pure component spectra (186) 443

PCBs
GC method for the analysis of polychlorinated biphenyls in transformer oil (189) 425

PCBs
Analysis of alkylated mixtures of polychlorinated biphenyls by capillary GC-MS (191) 207

PCBs
Evaluation of macroreticular resins for the determination of low concentrations of chlorinated hydrocarbons in sea water and tap water (193) 357

PCBs
GC-MS method for identifying and determining polychlorinated biphenyls (196) 109

PCBs
General approach to the fractionation and class determination of complex mixtures of chlorinated aromatic compounds (197) 155

Peak identification
Peak identification in capillary GC by simultaneous flame ionization detection and ^{14}C-detection (198) 389

Peak resolution
Resolution of overlapping chromatographic peaks by parameter estimation of their model (191) 157

Peak shape
Chromatographic peak shape. III. Influence of kinetics parameters on solute profiles (186) 117

Peltatin
Isolation of α- and β-peltatin and podophyllotoxin by LC and analysis by HPLC (189) 276

Penicillic acid
HPLC analysis of the mycotoxin penicillic acid and its application to biological fluids (194) 387

Penicillin
Fast and simple method for the separation of intermediates and cofactors involved in the biosynthesis of cephalosporin C using chemically bonded C_{12} reversed-phase TLC (193) 511

Penicillins
Analysis of isoxazolyl penicillins and their metabolites in body fluids by HPLC (183) 339

Pentachlorophenol
Analysis of pentachlorophenol in waste water using HPLC (190) 203

Pentachlorophenol
Simple TLC method for detection of pentachlorophenol in sawdust and woodshavings (195) 441

Pentadecene isomers
Comparison of capillary columns coated with C_{87} hydrocarbon and squalane in the analysis of *n*-pentadecene isomers (191) 199

Pentafluorobenzyloxyamine
GC determination of low-molecular-weight carbonyl compounds in aqueous solution as their O-(2,3,4,5,6-pentafluorobenzyl) oximes (187) 413

Pentafluorophenyldialkylchlorosilanes
New electron-capturing pentafluorophenyldialkylchlorosilanes as versatile derivatizing reagents for GC (199) 123

Pentafluorophenylhydrazine
GC determination of low-molecular-weight carbonyl compounds in aqueous solution as their O-(2,3,4,5,6-pentafluorobenzyl) oximes (187) 413

Pentazocine
Syringe and column adsorption of tertiary amines in GC (196) 489

Peppermint oils
Quantitative determination of minor components in essential oils: determination of pulegone in peppermint oils (190) 471

Peptidase
Rapid chromatographic purification of dipeptidyl peptidase IV in human submaxillary gland (189) 233

Peptides
Fluorescence micro disc electrophoresis in sodium dodecyl sulfate. A simple and sensitive method applicable for routine laboratory analysis (181) 259

Peptides
HPLC determination of cyclosporin A in human plasma and urine (182) 454

Peptides
Separation of peptides and amino acids by ion-exchange chromatography of their copper complexes (183) 9

Peptides
Determination of specific urinary hydroxyproline-containing peptides in patients with lung cancer (183) 221

Peptides
Separation and quantification of angiotensins and some related peptides by HPLC (183) 303

Peptides
New developments in isoelectric focusing (Review) (184) 415

Peptides
Reversed-phase separation of optical isomers of Dns-amino acids and peptides using chiral metal chelate additives (185) 323

Peptides
HPLC of amino acids, peptides and proteins. XX. Investigation of the effect of pH and ion-pair formation on the retention of peptides on chemically-bonded hydrocarbonaceous stationary phases (185) 429

Peptides
Ligand-exchange chromatography of small peptides on copper(II)-modified silica gel: application to the study of the enzymatic degradation of methionine-enkephalin (186) 677

Peptides
Peptide separations on substituted polystyrene resins. Effect of cross-linkage (187) 27

Peptides
Separation of cyanogen bromide peptides of collagen by means of HPLC (190) 393

Peptides
HPLC of amino acids, peptides and proteins. XXI. The application of preparative reversed-phase HPLC for the purification of a synthetic underivatised peptide (192) 222

Peptides
Separation of alkylaminonaphthylenesulfonyl peptides and amino acids by HPLC. Methods for measuring melanotropin inhibiting factor breakdown (192) 341

Peptides
New horizontal flow-through coil planet centrifuge for counter-current chromatography. III. Separation and purification of dinitrophenyl amino acids and peptides (193) 37

Peptides
New sensitive technique for the quantitative analysis of initiation peptides (193) 491

Peptides
Hydrophobic chromatography with dynamically coated stationary phases. II. Dynamic cation-exchange separations of tyrosinyl peptides (194) 353

Peptides
Ion-pair HPLC of the insect neuropeptide proctolin and some analogs (194) 421

Peptides
Estimation of glutathione in rat liver by reversed-phase HPLC: separation from cysteine and γ-glutamylcysteine (194) 424

Peptides
Separation of underivatized dipeptides by HPLC on a weak anion-exchange bonded phase (195) 119

Peptides
Perfluoroalkanoic acids as lipophilic ion-pairing reagents in reversed-phase LC of peptides including secretin (195) 181

Peptides
Reversed-phase and soap TLC of peptides (195) 187

Peptides
HPLC and field desorption MS of hypothalamic oligopeptides (195) 369

Peptides
Study of the feasibility of separation of carrier ampholytes from peptides by hydrophobic interaction chromatography on octyl-Sepharose (196) 51

Peptides
Slab gel designed for enzymatic digestion of proteins in polyacrylamide gel slices and direct resolution of peptides (198) 169

Peptides
The isolation of methionine and ethionine by silver ligand chromatography and application to methionine-containing peptides (200) 185

Peptido-aminobenzophenone
Determination of peptido-aminobenzophenone (2-*o*-chlorobenzoyl-4-chloro-N-methyl-N'-glycylglycinanilide) by electron-capture GLC (182) 171

Perazine
Reliable routine method for determination of perazine in serum by TLC with an internal standard (183) 475

Perfluoroalkanoic acids
Perfluoroalkanoic acids as lipophilic ion-pairing reagents in reversed-phase LC of peptides including secretin (195) 181

Perloline
Determination of perloline in grass species (in German) (192) 199

Pesticides
GLC retention indices of 296 non-drug substances on SE-30 or OV-1 likely to be encountered in toxicological analyses (Review) (184) 185

Pesticides
Resolution, sensitivity and selectivity of a HPLC post-column fluorometric labeling technique for determination of carbamate insecticides (185) 615

Pesticides
Improved determination of dalapon residues in plant tissues and natural water by

derivatization and electron capture GC (187) 244

Pesticides
Comparison of the HPLC behaviour of *s*-triazine derivatives on various stationary phases (187) 341

Pesticides
Determination of halogenated anilines in urine by HPLC with an electrochemical detector (188) 412

Pesticides
TLC of 2-methyl-4-chlorophenoxyacetic acid and its soil metabolites (189) 73

Pesticides
GC determination of Lenacil and Pyrazon in soil and bacterial cultures (189) 89

Pesticides
Improved clean-up procedure for the determination of small residues of carbophenothion in mice (190) 201

Pesticides
Analysis of pentachlorophenol in waste water using HPLC (190) 203

Pesticides
Analysis of aminocarb formulations by HPLC (190) 208

Pesticides
HPLC analysis of formulations of phenoxycarboxylic acids, dicamba, ioxynil and bromoxynil mixtures (in French) (190) 401

Pesticides
Stimulated sequential dehydroxy-fluorination: a safe, convenient method for GC analysis of phosphate diesters with possible application to hydrolysates of organophosphorus pesticides and nerve gases (192) 173

Pesticides
Determination of carbofuran residues in potato, onion and turnip: comparative analyses using GC-selected ion monitoring or HPLC (193) 213

Pesticides
Determination of 2,6-dichlorobenzonitrile residues in soil and water (in German) (193) 320

Pesticides
Evaluation of macroreticular resins for the determination of low concentrations of chlorinated hydrocarbons in sea water and tap water (193) 357

Pesticides
Detection of aminocarb and its major metabolites by TLC (194) 100

Pesticides
GC measurement of volatility of herbicide esters (195) 75

Pesticides
Application of a direct aqueous acetylation technique to the GC quantitation of nitrophenols and 1-naphthol in environmental water samples (195) 105

Pesticides
Direct analysis of the herbicide barban in wheat products by reversed-phase LC (195) 113

Pesticides
Analysis of barban formulations by HPLC (195) 151

Pesticides
OV-17-QF-1 capillary column for organochlorine pesticide analysis (195) 265

Pesticides
Direct determination of propoxur in plant tissues by GC with an alkali flame-ionization detector (197) 263

Pesticides
Determination of residues of methazole and its metabolites, 1-(3,4-dichlorophenyl)-3-methylurea and 1-(3,4-dichlorophenyl)urea in soil by HPLC (197) 267

Pesticides
Retention behavior of chlorinated benzenes, chlorinated phenols, Aldrin, and Aroclor 1242 on Tenax-GC, Chromosorb 101, Florisil, and Carbopack C HT (200) 115

Pesticides
Simplified method for the analysis of some carbamate insecticides in foliage, forest soil and fish tissue by direct GLC (200) 179

Pesticides
GLC analysis of aminocarb and seven derivatives (200) 228

Pesticides
Determination of amitrole by ion-pair HPLC (200) 234

Pesticides
GC determination of Metalaxyl (Ridomil) residues in tobacco (200) 313

Pesticides
Estimation of the systemic fungicide Ridomil by TLC (200) 317

Petroleum hydrocarbons
Quantification of petroleum-type hydrocarbons in avian tissue (187) 153

pH effect
HPLC of amino acids, peptides and proteins. XX. Investigation of the effect of pH and ion-pair formation on the retention of peptides on chemically-bonded hydrocarbonaceous stationary phases (185) 429

pH effect
Combined effects of pH and surface-active-ion concentration in reversed-phase LC (199) 307

Phages
Affinity chromatography of cells and cell membranes (Review) (184) 471

Pharmalytes
Gel chromatographic comparison of the molecular weight distributions of Ampholine, Servalyte and Pharmalyte carrier ampholytes used in isoelectric focusing (198) 301

Phase transitions
GC evidence for phase transitions in very compact octadecyl bonded silicas (200) 95

Phenanthrimidazoles
Phenanthrimidazoles as fluorescent derivatives in the analysis of fatty acids by HPLC (189) 359

Phencyclidine
Capillary GC with a nitrogen detector for measurement of phencyclidine, ketamine and other arylcycloalkylamines in the picogram range (193) 157

Phenethylamines
Cactus alkaloids. XL. Identification of mescaline and other β-phenethylamines in *Pereskia, Pereskiopsis* and *Islaya* by use of fluorescamine conjugates (189) 79

Phenethylamines
Microdetermination of stimulant drugs in urine by HPLC (196) 334

Phenobarbital
Microassay for primidone and its metabolites phenylethylmalondiamide, phenobarbital, and *p*-hydroxyphenobarbital in human serum, saliva, breast milk and tissues by GC-MS using selected ion monitoring (182) 71

Phenobarbital
Improved GLC method for the simultaneous determination of phenobarbital, phenytoin, carbamazepine and primidone in biological fluids (182) 116

Phenobarbital
A quantitative method for the simultaneous determination of carbamazepine, mephenytoin, phenylethylmalonamide, phenobarbital, phenytoin and primidone in serum by thin-layer chromatography. Additional comments to the analysis of carbamazepine-10,11-epoxide (183) 387

Phenobarbital
Rapid method for simultaneous determination of phenobarbital, diphenylhydantoin and their main hydroxylated metabolites by nitrogen-selective GC (183) 505

Phenol derivatives
Structure and sorption properties of porous copolymers of styrene, acrylonitrile and divinylbenzene (196) 75

Phenolic acid esters
HPLC separation of naturally occurring esters of phenolic acids (187) 181

Phenolic acids
GC-MS identification of phenolic acids in recent sediments (193) 89

Phenolic aldehydes
Chromatography in aqueous solution on styrene-divinylbenzene resins (193) 197

Phenolic amines
GC of phenolic amines, 3-methoxycatechol-amines, indoleamines and related amines as their N,O-ethyloxycarbonyl derivatives (194) 399

Phenolic compounds
HPLC, GLC and TLC of naturally occurring flavonoids, phenolic and related compounds (187) 255

Phenolic compounds
Alkaline chloramine-T reagent for the detection of phenolic compounds on thin-layer plates (196) 309

Phenol-impregnated thin layers
Chromatographic behaviour of aliphatic amines on phenol-impregnated thin layers (196) 225

Phenols
Chromatography in aqueous solution on styrene-divinylbenzene resins (193) 197

Phenols
Reversed-phase and soap TLC of phenols (195) 339

Phenols
Relationship between the retention indices of phenols on polar and non-polar stationary phases (196) 217

Phenols
Reversed-phase HPLC. I. Relationship between R_M values and Hansch's π parameters for a series of phenols (197) 43

Phenols
Study of the interactions between substituted phenols and alcohols and dibenzyl sulphoxide by means of GC (198) 495

Phenothiazines
Rapid, sensitive determination of unchanged promethazine in biological material using a nitrogen-selective flame ionization detector. Identification of metabolites by GC-MS (192) 375

Phenothiazines
HPLC separation of the N- and S-oxides of fluphenazine and fluphenazine decanoate (194) 416

Phenothiazines
Use of PTFE coils in post-column photochemical reactors for LC: application to pharmaceuticals (199) 239

Phenoxycarboxylic acids
HPLC analysis of formulations of

phenoxycarboxylic acids, dicamba, ioxynil and bromoxynil mixtures (in French) (190) 401

Phenylacetic acid
Comparative study of derivatisation procedures for the quantitative determination of the auxin, phenylacetic acid, by GC (192) 429

Phenylalanine
Rapid automated ion-exchange analysis of plasma tyrosine and phenylalanine with data print-out (181) 337

Phenylalanine
Presence of sodium azide during acid hydrolysis of protein samples causes the destruction of tyrosine, phenylalanine and histidine (189) 106

Phenylarsine oxide
Derivatization of dithiols and certain monothiols with phenylarsine oxide for GC (200) 171

Phenylbutazone
Reversed-phase HPLC of phenylbutazone in body fluids (181) 132

Phenylbutazone
Simultaneous determination of phenylbutazone and its metabolites in plasma and urine by HPLC (183) 331

Phenylethylene glycol
Improved GC method for measuring phenylethylene glycol. Application to the determination of styrene monooxygenase and epoxide hydrase activities (188) 400

Phenylethylmalonamide
A quantitative method for the simultaneous determination of carbamazepine, mephenytoin, phenylethylmalonamide, phenobarbital, phenytoin and primidone in serum by thin-layer chromatography. Additional comments to the analysis of carbamazepine-10,11-epoxide (183) 387

Phenylethylmalonidiamide
Microassay for primidone and its metabolites phenylethylmalondiamide, phenobarbital, and *p*-hydroxyphenobarbital in human serum, saliva, breast milk and tissues by GC-MS using selected ion monitoring (182) 71

Phenytoin
Improved GLC method for the simultaneous determination of phenobarbital, phenytoin, carbamazepine and primidone in biological fluids (182) 116

Phenytoin
A quantitative method for the simultaneous determination of carbamazepine, mephenytoin, phenylethylmalonamide, phenobarbital, phenytoin and primidone in serum by thin-layer chromatography. Additional comments to the analysis of carbamazepine-10,11-epoxide (183) 387

Pheromones
GC detection and structure analysis of volatile pheromones in insects (186) 605

Phlorizin
Purification of phlorizin by CC on Sephadex LH-20 with aqueous propan-2-ol (188) 439

Phloroglucinols
Separation of naturally occurring acylphloroglucinols by HPLC (188) 213

Phosphate
Trimethylsilyl esters of plant acids and their application in GLC. Preparation, kinetics of silylation and solvent effects on quantitative reaction and stability of the derivatives (in German) (194) 33

Phosphate diesters
Stimulated sequential dehydroxy-fluorination: a safe, convenient method for GC analysis of phosphate diesters with possible application to hydrolysates of organophosphorus pesticides and nerve gases (192) 173

Phosphate esters
Paper ionophoresis of pancreatic islet phosphate esters in a magnesium-containing buffer (187) 235

Phosphates
Chromatographic separation and identification of short-chain acid-soluble polyphosphates from *Saccharomyces cerevisiae* (189) 440

Phosphates
Ion-exclusion chromatography of condensed phosphates on a cation-exchange resin (190) 77

Phosphates
Separation and determination of the degradation products of tributyl phosphate by high-speed analytical isotachophoresis (195) 303

Phosphates
Optimization of HPLC of inorganic polyphosphates for routine analyses (196) 471

Phosphates, organic
Sintered TLC (Review) (184) 37

Phosphatidylcholine
Comparative determination of plasma phospholipids by automated GLC and manual colorimetric phosphorus methods (182) 1

Phosphatidylcholine
GC determination of the fatty acid pattern of red cell membrane plasmalogens in healthy children (182) 27

Phosphatidyl ethanolamine
GC determination of the fatty acid pattern of red cell membrane plasmalogens in healthy children (182) 27

Phosphoglycerides
Isocratic HPLC separation of phosphoglycerides and lysophosphoglycerides (197) 79

Phospholipase
Assay of phospholipase D on a TLC plate (188) 292

Phospholipids
Comparative determination of plasma phospholipids by automated GLC and manual colorimetric phosphorus methods (182) 1

Phospholipids
GC determination of the fatty acid pattern of red cell membrane plasmalogens in healthy children (182) 27

Phospholipids
Lipid content of human platelets quantitated by TLC in combination with flame ionization detection (182) 301

Phospholipids
Miniature two-dimensional TLC separation of lecithin and sphingomyelin (182) 311

Phospholipids
Isocratic HPLC separation of phosphoglycerides and lysophosphoglycerides (197) 79

Phospholipids
HPLC of diglyceride *p*-nitrobenzoates. An approach to molecular analysis of phospholipids (198) 520

Phosphonacetylaspartic acid
Quantitation of the antitumor agent N-(phosphonacetyl)-L-aspartic acid in human plasma and urine by GC-MS-selected ion monitoring (182) 163

Phosphonic acids
TLC of phosphonic acids (187) 447

Phosphonoacetylaspartic acid
MS technique for the determination of N-phosphonoacetyl-L-aspartic acid in serum (181) 195

Phosphorus oxo anions
Gel chromatographic separations of monomeric oxo anions of phosphorus on Sephadex G-10 and Bio-Gel P-2 (190) 161

Photochemical reaction detectors
Use of PTFE coils in post-column photochemical reactors for LC: application to pharmaceuticals (199) 239

Photochemical reactor
Construction of a simple gas chromatograph with photolytic intersect (189) 119

Photodensitometry
Correction for deviation from the Lambert-Beer law in the quantitation of thin-layer chromatograms by photodensitometry (192) 208

Photoionization detector
The photoionization detector. Theory, performance and application as a low-level monitor of oil vapour (190) 263

Photoionization detector
Application of a unified theory of GC photoionization detector response to a 10.2-eV detector (200) 35

Photolysis
Construction of a simple gas chromatograph with photolytic intersect (189) 119

Photolysis
GC with *in situ* photolysis (189) 127

Phthalate
Empirical approach to the universal calibration of isomers of phthalate esters and similar compounds in size exclusion chromatography (192) 295

Phthalate
Determination by electron-capture GC of mono- and di(2-ethylhexyl) phthalate in intravenous solutions stored in poly(vinyl chloride) bags (193) 405

Phthalate esters
On-line trace enrichment in HPLC using a pre-column (185) 483

Phthalates
Background contamination by phthalates commonly encountered in the chromatographic analysis of lipid samples (189) 421

Phthalazinone
Determination of hydralazine metabolites: 4-hydrazino-phthalazin-1-one and N-acetylhydrazinophthalazin-1-one by GC and *s*-triazolo[3,4-α]phthalazine and phthalazinone by HPLC (183) 167

Phthalocyanines
Silica gel modified with phthalocyanines as a stationary phase in gas-solid chromatography (189) 25

Phylloquinone
Resolution of phylloquinone (vitamin K_1), phylloquinone 2,3-epoxide, 2-chloro-phylloquinone and their geometric isomers by HPLC (200) 293

Physalins
Reversed-phase HPLC separation and analysis of some physalins (13,14-*seco*-16,24-cyclo-steroids) (198) 203

Physostigmine
Determination of pilocarpine, physostigmine, its degradation product rubreserine and preservatives by HPLC (198) 529

Phytanic acid
The use of formic acid in carrier gas. Rapid method for identification and determination of phytanic acid by GC-MS-chemical ionization (182) 269

Phytic acid
Analysis of phytic acid by HPLC (197) 274

Pigments, organic
TLC of slightly soluble substances at elevated temperature (186) 575

Pilocarpine
Determination of pilocarpine, physostigmine, its degradation product rubreserine and preservatives by HPLC (198) 529

Pindolol
Determination of pindolol in human plasma by HPLC (183) 255

Pipecoloxylidide
Simultaneous determination of bupivacaine and 2,6-pipecoloxylidide in serum by GLC (182) 226

Pipemidic acid
Estimation of pipemidic acid and N-nitrosopipemidic acid by HPLC (200) 245

Piperine
High-temperature quantitative glass capillary GC. Analysis of piperine and of quinine-quinidine (199) 105

Pirimicarb
Simplified method for the analysis of some carbamate insecticides in foliage, forest soil and fish tissue by direct GLC (200) 179

Piromidic acid
GC determination of piromidic acid (193) 476

Piroxicam
Analysis of piroxicam in plasma by HPLC (183) 104

Pirquinozol
HPLC of the oral anti-asthmatic agent pirquinozol and its active metabolite (200) 282

Pivmecillinam hydrochloride
Determination of mecillinam in urine by reversed-phase HPLC (181) 272

Plant extracts
Trimethylsilyl esters of plant acids and their application in GLC. Preparation, kinetics of silylation and solvent effects on quantitative reaction and stability of the derivatives (in German) (194) 33

Plant extracts
Separation of benzoic acid derivatives from plant extracts on Sephadex G-10 columns (in German) (198) 373

Plant hormones
HPLC of plant hormones. I. Separation of plant hormones of the indole type (191) 129

Plant hormones
High-performance steric exclusion chromatography of plant hormones (198) 57

Plasticisers
GLC retention indices of 296 non-drug substances on SE-30 or OV-1 likely to be encountered in toxicological analyses (Review) (184) 185

Plasticizers
Determination by electron-capture GC of mono- and di(2-ethylhexyl) phthalate in intravenous solutions stored in poly(vinyl chloride) bags (193) 405

Platinum
GC of volatile metal dialkyldithiophosphates (193) 53

Podophyllotoxin
Analysis of the anticancer drugs VP 16-213 and VM 26 and their metabolites by HPLC (182) 211

Podophyllotoxin
Isolation of α- and β-peltatin and podophyllotoxin by LC and analysis by HPLC (189) 276

Poiseuille flow
The HETP of an unretained component of fluid flow in a coiled round pipe (186) 341

Poiseuille flow
Early phases of the dispersion of a sample injected in Poiseuille flow (186) 353

Polarity
Polarity of some chromatographic materials intended for the separation of biopolymers: spectroscopic label technique (196) 39

Polarity range
"Sequential isocratic step" LC: a high-throughput, problem-solving approach with sensitive, near-universal detection for wide polarity mixtures (199) 219

Polarity variation
Binary liquid phase system for GLC with *in situ* variable polarity (189) 255

Polarographic detector
Design and selective application of a dropping mercury electrode amperometric detector in column LC (186) 489

Polarographic detector
Polarographic detection for HPLC using a flow-through detector (191) 47

Pollutants
Analysis of polynuclear aromatic hydrocarbons in environmental waters by HPLC (185) 655

Pollutants
Analysis of non-volatile organic compounds in water by HPLC (185) 671

Pollutants
Vapor-phase silylation of alcohols for air analysis (186) 273

Pollutants
Direct aqueous injection GC for the analysis of trace organics in water (186) 785

Pollutants
Evaluation of chromatographic sorbents used in air pollution studies (186) 811

Pollutants
Carbon skeleton-GC techniques and their applications (187) 47

Pollutants
Quantification of petroleum-type hydrocarbons in avian tissue (187) 153

Pollutants
Rapid analysis of chlorinated dioxins in complex organic mixtures by application of a GC-MS-calculator system with user-developed software (189) 53

Pollutants
Background contamination by phthalates commonly encountered in the chromatographic analysis of lipid samples (189) 421

Pollutants
GC method for the analysis of polychlorinated biphenyls in transformer oil (189) 425

Pollutants
Determination of benzo[*a*]pyrene and other polyaromatic hydrocarbons in petroleum oils by direct LC (190) 85

Pollutants
Determination of 1,2-dibromoethane, 1,2-dichloroethane and benzene in ambient air using porous polymer traps and GC-MS analysis with selected ion monitoring (190) 97

Pollutants
Determination of airborne free monomeric aromatic and aliphatic isocyanates by HPLC (190) 175

Pollutants
Analysis of alkylated mixtures of polychlorinated biphenyls by capillary GC-MS (191) 207

Pollutants
Direct determination of organic compounds in water using steam-solid chromatography (191) 221

Pollutants
Evaluation of macroreticular resins for the determination of low concentrations of chlorinated hydrocarbons in sea water and tap water (193) 357

Pollutants
TLC of chlorinated guaiacols (194) 55

Pollutants
Application of a direct aqueous acetylation technique to the GC quantitation of nitrophenols and 1-naphthol in environmental water samples (195) 105

Pollutants
Quantitative analysis of polynitrophenols in water in the micro- to nanogram range by reversed-phase ion-pair LC (195) 435

Pollutants
GC-MS method for identifying and determining polychlorinated biphenyls (196) 109

Pollutants
General approach to the fractionation and class determination of complex mixtures of chlorinated aromatic compounds (197) 155

Pollutants
HPLC in the analysis of chorinated dibenzodioxins and dibenzofurans in chicken liver and wood shaving samples (197) 171

Pollutants
Discharge of elemental sulfur and its distribution in surficial sediments in the region of the Buccaneer oil field (199) 161

Polyacetylenes
Visualization reagents for sesquiterpene lactones and polyacetylenes on thin-layer chromatograms (189) 187

Polyacrylamide-boronate beads
Polyacrylamide-boronate beads saturated with biomolecules: a new general support for affinity chromatography of enzymes (189) 225

Polyamines
Automated ion-exchange chromatographic analysis of usual and unusual natural polyamines (187) 442

Polyaromatic hydrocarbons
Determination of benzo[*a*]pyrene and other polyaromatic hydrocarbons in petroleum oils by direct LC (190) 85

Polybrominated biphenyls
GC-MS analysis of polybrominated biphenyl constituents of Firemaster FF-1 (197) 258

Polychlorinated alkanes
Carbon skeleton-GC techniques and their applications (187) 47

Polychlorinated biphenyls, see PCBs

Polycyclic aromatic compounds
Fractionation and GC analysis of aza-arenes in complex mixtures (199) 371

Polycyclic aromatic hydrocarbons, see also polynuclear aromatic hydrocarbons

Polycyclic aromatic hydrocarbons
Retention characteristics of several bonded-phase LC columns for some polycyclic aromatic hydrocarbons (188) 115

Polycyclic aromatic hydrocarbons
Alteration of the electron-capture detector response to polycyclic aromatic hydrocarbons by oxygen doping of the carrier gas (197) 51

Polycyclic aromatic hydrocarbons
Concentration and determination of polycyclic aromatic hydrocarbons in aqueous samples on graphitized carbon black (198) 143

Polydentate phosphonium bromides
Polydentate phosphonium salts for the deactivation of wall-coated open-tubular

columns. Comparison of temperature stability with other deactivation methods (187) 57

Polyethylene
Application of pyrolysis-capillary GC to the characterization of polyethylene (191) 181

Polyethylene glycol
Simple method for GLC analysis of polyethylene glycol 400 in biological fluids (183) 72

Poly(ethylene glycol)
Study of the adsorption effects at the surface of poly(ethylene glycol)-coated column packings (186) 63

Polyethylene glycol
Separation range and separation efficiency in high-speed gel filtration on TSK-GEL SW columns (190) 297

Polyhydroxy compounds
Distribution ratios and elution behavior of carbamates and polyhydroxy compounds from Sephadex G-15 (190) 448

Polymer analysis
Retention characteristics of various polymers in thermal field-flow fractionation (186) 37

Polymer degradation
Comparison between conventional and cross-correlation GC for evolved gas analysis (186) 145

Polymers
Use of water-soluble biospecific polymers for the purification of proteins (Review) (184) 325

Polymers
High-performance exclusion chromatography of water-soluble polymers with chemically bonded stationary phases (185) 305

Polymers
Comprehensive study of porous polymeric stationary phases for chromatography (186) 317

Polymers
Fractionation of polymers by gel permeation chromatography: an experimental and theoretical approach (192) 259

Polymers
Determination of oligomers in vinyl chloride polymers by steric exclusion chromatography (193) 235

Polymers
Determination of the composition of copolymers as a function of molecular weight by pyrolysis GC size-exclusion chromatography (194) 163

Polymers
Reactive polymers. XXXI. Thermally treated polymeric sorbents based on glycidylmethacrylate copolymers (194) 297

Polymer samples
Band dispersion in high-performance exclusion chromatography (185) 289

Polymer solutions
Concentration effects in gel permeation chromatography. VI. Non-Newtonian behaviour of polymer solutions (187) 21

Polymer sorbents
Polyfunctional polymer sorbents (186) 227

Polymethylphenylsiloxanes
Crosslinked polymethylphenylsiloxanes as adsorbents for GC (189) 169

Polynitrophenols
Quantitative analysis of polynitrophenols in water in the micro- to nanogram range by reversed-phase ion-pair LC (195) 435

Polynuclear aromatic hydrocarbons, see also polycyclic aromatic hydrocarbons

Polynuclear aromatic hydrocarbons
Analysis of polynuclear aromatic hydrocarbons in environmental waters by HPLC (185) 655

Polynuclear aromatic hydrocarbons
Adsorption chromatography capillary columns. I. Rubidium chloride (196) 237

Polyolefins
Poly(α-olefins): novel non-polar stationary phases for GC (186) 259

Polyolefin sintered sheets
Sintered TLC (Review) (184) 37

Polyphenoloxidase
Purification of grape polyphenoloxidase with hydrophobic chromatography (192) 232

Polyphosphates
Chromatographic separation and identification of short-chain acid-soluble polyphosphates from *Saccharomyces cerevisiae* (189) 440

Polyphosphates
Optimization of HPLC of inorganic polyphosphates for routine analyses (196) 471

Polysaccharides
Fingerprinting of heparins by low-amperage electrophoresis in barium acetate (195) 398

Polystyrene
High-performance gel permeation chromatography of polystyrene with silica microspheres (188) 333

Polystyrene resins
Peptide separations on substituted polystyrene resins. Effect of cross-linkage (187) 27

Polyurethane prepolymers
Determination of free toluene diisocyanate in polyurethane prepolymers by HPLC (198) 212

Poly(vinyl chloride) resins
Determination of residual vinyl chloride in poly(vinyl chloride) resins (198) 337

Poly(vinylidene chloride)
GC determination of vinylidene chloride monomer in packaging films and in foods (197) 71

Porapak columns
GC analysis of light hydrocarbons for the characterization of Porapak columns (196) 245

Pore size
Empirical correlation of layer thickness and support pore size in "chemically bonded" phases (195) 25

Pore size distributions
Separation of C_6 cyclic hydrocarbons before the inert peak on microporous silica: a case of negative adsorption (199) 23

Porphobilinogen
Determination of 5-aminolaevulinic acid and porphobilinogen by HPLC (185) 605

Porphyrin esters
Quantitation of porphyrin esters on thin-layer chromatograms (198) 227

Porphyrins
HPLC determination of protoporphyrin and zinc protoporphyrin in blood (181) 319

Porphyrins
Second derivative-HPLC-fluorometric detection of porphyrins in chick embryo liver cell culture medium (189) 209

Porphyrins
Diagnosis of porphyrias by ion-pair HPLC (199) 339

Prazosin
HPLC determination of prazosin in human plasma, whole blood and urine (183) 367

Prednisolone
Improved HPLC separation of decomposition products of prednisolone by adding sulphite to the mobile phase (193) 480

Pregnenolone
Pregnenolone separation from cholesterol using Sephadex LH-20 minicolumns (190) 256

Preparative counter-current chromatography
Preparative counter-current chromatography with a slowly rotating glass coil (196) 295

Primidone
Microassay for primidone and its metabolites phenylethylmalondiamide, phenobarbital, and *p*-hydroxyphenobarbital in human serum, saliva, breast milk and tissues by GC-MS using selected ion monitoring (182) 71

Primidone
Improved GLC method for the simultaneous determination of phenobarbital, phenytoin, carbamazepine and primidone in biological fluids (182) 116

Primidone
A quantitative method for the simultaneous determination of carbamazepine, mephenytoin, phenylethylmalonamide, phenobarbital, phenytoin and primidone in serum by thin-layer chromatography. Additional comments to the analysis of carbamazepine-10,11-epoxide (183) 387

Probenecid
Rapid HPLC method for the determination of probenecid in biological fluids (182) 252

Procainamide
Comparison of HPLC with an enzyme multiplied immunoassay technique for the analysis of serum procainamide and N-acetylprocainamide (181) 490

Procarbazine hydrochloride
Application of an electrochemical detector to the determination of procarbazine hydrochloride by HPLC (190) 359

Proctolin
Ion-pair HPLC of the insect neuropeptide proctolin and some analogs (194) 421

Procyanidins
Reversed-phase gradient HPLC of procyanidins and their oxidation products in ciders and wines, optimised by Snyder's procedures (194) 62

Profiles, chromatographic
Simultaneous analysis of ATP, ADP, AMP, and other purines in human erythrocytes by HPLC (181) 103

Profiles, chromatographic
Differentiation of the polysaccharide side-chains of glycoproteins by a fingerprinting technique: heterogeneity of human gastric mucin (181) 108

Profiles, chromatographic
Comparison of HPLC serum profiles of humans and dogs (181) 169

Profiles, chromatographic
High-performance anion-exchange chromatography of UV-absorbing constituents of human urine (181) 311

Profiles, chromatographic
Volatile metabolites in sera of normal and diabetic patients (182) 137

Profiles, chromatographic
Identification and quantitation of nucleosides, bases and other UV-absorbing compounds in serum, using reversed-phase HPLC. I. Chromatographic methodology (186) 647

Profiles, chromatographic
Identification and quantitation of nucleosides, bases and other UV-absorbing compounds in serum, using reversed-phase HPLC. II. Evaluation of human sera (186) 659

Profiles, chromatographic
GC-MS of saturated and unsaturated dicarboxylic acids in urine (199) 181

Profiles, isotachophoretic
Comparison of urinary excretion of UV-absorbing constituents in healthy subjects and patients with rheumatoid arthritis using analytical isotachophoresis (181) 347

Progesterone
HPLC of the reduction products of progesterone (190) 169

Programmed amino acid analysis
Program for processing amino acid data with a programmable pocket calculator (194) 69

Proguanil
HPLC of proguanil, cycloguanil and 4-chlorophenylbiguanide using hydrophobic pairing ion and its application to serum assay (182) 359

Proline
GLC of free amino acids in the hyaloplasm of rat cerebral, cerebellar and ocular tissues, and in skeletal and heart muscle (182) 147

Proline
Employment of GLC for the analysis of collagen amino acids in biopsy tissue (182) 155

Proline
Ligand-exchange chromatography. I. Resolution of L- and D-proline on a copper(II)-proline complex bound to microparticulate silica gel (192) 228

Promethazine
Rapid, sensitive determination of unchanged promethazine in biological material using a nitrogen-selective flame ionization detector. Identification of metabolites by GC-MS (192) 375

Propiomazine
Determination of propiomazine and its N-demethyl metabolite in plasma by GC with alkali flame ionization detection (183) 229

Propionaldehyde
GLC analysis of dimedone derivatives of formaldehyde and other aliphatic aldehydes on capillary columns (191) 239

Propoxur
Direct determination of propoxur in plant tissues by GC with an alkali flame-ionization detector (197) 263

Propoxur
Simplified method for the analysis of some carbamate insecticides in foliage, forest soil and fish tissue by direct GLC (200) 179

Propranolol
Determination of propranolol and its major metabolites in plasma and urine by HPLC without solvent extraction (183) 213

Propylnaphthalenes
Qualitative and quantitative analysis of propylnaphthalenes by GC on silicone stationary phases (197) 92

Prostaglandins
Capillary column GLC-MS assay for 7α-hydroxy-5,11-diketotetranorprostane-1,16-dioic acid, the major human urinary metabolite of prostaglandins E_1 and E_2 (181) 85

Prostaglandins
Prostaglandins E and F, and 19-hydroxylated E and F (series I and II) in semen of fertile men. GC and LC separation with selected ion detection (181) 295

Prostaglandins
Determination of an antisecretory trimethyl prostaglandin E_2 analog in human plasma by combined capillary column GC-negative chemical ionization MS (183) 411

Prostaglandins
Adsorption chromatographic separation of ^{125}I-labelled prostaglandin $F_{2\alpha}$ and prostaglandin E_2 tyrosine methyl ester (189) 433

Prostaglandins
Separation and quantitation of [^{14}C]prostaglandin E_1 from lung effluent metabolites by HPLC (192) 463

Prostaglandins
Simple and rapid separation of certain prostaglandins by reversed-phase HPLC (194) 85

Proteases
Affinity chromatographic interactions of proteases with low-molecular-weight soybean protease inhibitors (195) 385

Protein hydrolysates
Continuous automatic nitrogen determination for gel chromatography of protein enzymatic hydrolysates (193) 496

Protein-protein interactions
Protein-protein interactions studied by counter-current distribution. II. Test of the theoretical model using bovine serum albumin and L-tryptophan (196) 207

Proteins
Use of water-soluble biospecific polymers for the purification of proteins (Review) (184) 325

Proteins
Preparation of a porous microparticulate anion-exchange chromatography support for proteins (185) 375

Proteins
Interactions of proteins with bonded-phase ion exchangers (185) 393

Proteins
Use of hydrophobic interaction methods in the isolation of proteins from endocrine and paraendocrine tissues and cells by HPLC (185) 413

Proteins
HPLC of amino acids, peptides and proteins. XX. Investigation of the effect of pH and ion-pair formation on the retention of peptides on chemically-bonded hydrocarbonaceous stationary phases (185) 429

Proteins
Preparative separation of proteins and enzymes in the mean molecular-weight range of 10,000-100,000 on LiChrosorb Diol packing by high-performance size-exclusion chromatography (185) 445

Proteins
High-speed steric exclusion chromatography of biopolymers (185) 453

Proteins
Application of 1,1'-carbonyldiimidazole-activated agarose for the purification of proteins. II. The use of an activated matrix devoid of additional charged groups for the purification of thyroid proteins (185) 463

Proteins
Covalent coupling of cholic acid to aminohexylamino-Sepharose 4B and its use in affinity chromatography of serum albumin (187) 409

Proteins
Presence of sodium azide during acid hydrolysis of protein samples causes the destruction of tyrosine, phenylalanine and histidine (189) 106

Proteins
Titration curves of proteins by combined isoelectric focusing-electrophoresis in highly porous polyacrylamide matrices (189) 317

Proteins
Trypan blue staining of proteins fractionated by acrylamide disc gel electrophoresis (189) 410

Proteins
Simple method for the purification of C1q, a subcomponent of the first component of complement by affinity chromatography using IgG-Sepharose (189) 428

Proteins
Separation range and separation efficiency in high-speed gel filtration on TSK-GEL SW columns (190) 297

Proteins
Dissolvable disulphide-polyacrylamide gels for the electrophoretic analysis of chromosomal proteins and for affino-electrophoresis of thiol-proteins (190) 437

Proteins
Investigation of vinca alkaloid-plasma membrane interactions by detergent gel chromatography (191) 287

Proteins
HPLC of amino acids, peptides and proteins. XXI. The application of preparative reversed-phase HPLC for the purification of a synthetic underivatised peptide (192) 222

Proteins
Determination of the free electrophoretic mobility of proteins by polyacrylamide gradient gel electrophoresis: A new approach (192) 402

Proteins
Chromatography of chromatin proteins on Cibacron Blue F3G-A-agarose (192) 494

Proteins
Size and charge distribution of macromolecules in living systems (193) 1

Proteins
High-speed gel filtration of proteins in sodium dodecyl sulphate aqueous solution on TSK-GEL SW type (193) 29

Proteins
High-speed gel filtration of proteins in 6 *M* guanidine hydrochloride on TSK-GEL SW columns (193) 458

Proteins
Electrofocusing of stroma-free hemolyzates of human erythrocytes; comparison of fresh, stored and chemically modified samples (193) 515

Proteins
Electrofocusing of stroma-free haemoglobin on a thin layer of agarose gel (196) 142

Proteins
Use of silicone-coated porous glass for exclusion chromatography in an aqueous medium (196) 485

Proteins
HPLC of myosin's subfragment 1 and light chains on Spherogel TSK-type columns (197) 207

Proteins
Quantitative analysis of non-histone chromosomal proteins HMG 14 and HMG 17 by polyacrylamide gel electrophoresis (198) 80

Proteins
Slab gel designed for enzymatic digestion of proteins in polyacrylamide gel slices and direct resolution of peptides (198) 169

Proteins
An approach to ampholyte-displacement chromatography (200) 57

Proteins
Quantitative staining of submicrogram amounts of histone and high-mobility group proteins on sodium dodecylsulphate-polyacrylamide gels (200) 200

Proteolytic enzymes
Solid-phase reagents for LC detection (185) 497

Protoporphyrin
HPLC determination of protoporphyrin and zinc protoporphyrin in blood (181) 319

PR-toxin
Photodensitometric determination of "PR-toxin", metabolite of *Penicillium roqueforti* (in French) (198) 481

Pseudoephedrine
Analysis of perfluoroacyl derivatives of ephedrine, pseudoephedrine and analogues by GC and MS. I. Trifluoroacetyl derivatives (190) 53

Psilocin
Determination of hallucinogenic components of *Psilocybe* mushrooms using HPLC (196) 180

Psilocybin
Determination of hallucinogenic components of *Psilocybe* mushrooms using HPLC (196) 180

Psychedelic drugs
Cactus alkaloids. XL. Identification of mescaline and other β-phenethylamines in *Pereskia, Pereskiopsis* and *Islaya* by use of fluorescamine conjugates (189) 79

Psychotropic drugs
Determination of carpipramine in plasma by HPLC (183) 246

Pterins
Quantitative determination of pterins in biological fluids by HPLC (188) 363

Pulegone
Quantitative determination of minor components in essential oils: determination of pulegone in peppermint oils (190) 471

Purine nucleoside phosphorylase
Optimized assay for purine nucleoside phosphorylase by reversed-phase HPLC (199) 275

Purine ribonucleotides
Analysis of purine ribonucleotides and deoxyribonucleotides in cell extracts by HPLC (198) 435

Purines
Simultaneous analysis of ATP, ADP, AMP, and other purines in human erythrocytes by HPLC (181) 103

Purines
Quantitative HPLC of nucleosides and bases in human plasma (183) 421

Purines
Determination of selected pyrimidines, purines and their metabolites in serum and urine by reversed-phase ion-pair chromatography (199) 345

Purity test
Detection of trace organic impurities in binary solvent systems: a solvent purity test (188) 193

Putrefactive compounds
GLC retention indices of 296 non-drug substances on SE-30 or OV-1 likely to be encountered in toxicological analyses (Review) (184) 185

Pyramin
GC determination of Lenacil and Pyrazon in soil and bacterial cultures (189) 89

Pyrazines
Studies of pyrazines as their n-π charge-transfer complexes with some nitro aromatic compounds (194) 254

Pyrazon
GC determination of Lenacil and Pyrazon in soil and bacterial cultures (189) 89

Pyrethrins
Examination of the conversion products of pyrethrins and allethrin formulations exposed to sunlight by GC and MS (198) 317

Pyridine derivatives
Hydrophobic chromatography and bioanalysis of some polar pyridine derivatives used as antilipolytic agents (187) 189

Pyridopyrimidine
Relation of GC behaviour to the chemical structure of pyrido[1,2*a*]pyrimidine derivatives (191) 245

Pyridostigmine
Reversed-phase, ion-pair LC of quaternary ammonium compounds. Determination of pyridostigmine, neostigmine and edrophonium in biological fluids (183) 193

Pyridoxal
Separation of vitamin B_6 components by HPLC (188) 285

Pyridoxamine
Separation of vitamin B_6 components by HPLC (188) 285

Pyridoxine
HPLC determination of pyridoxine and congeners in biological fluids of man after high-dose therapy (183) 492

Pyridoxine
LC separation and quantification of B_6 vitamers at plasma concentration levels (196) 176

Pyridoxol
Separation of vitamin B_6 components by HPLC (188) 285

Pyrimidine bases
Analysis of pyrimidine bases in biological materials by GC-MS (190) 107

Pyrimidine deoxypentanucleoside tetraphosphates
Separation and sequencing of the sequence isomers of pyrimidine deoxypentanucleoside tetraphosphates by HPLC (188) 273

Pyrimidine dimers
Separation of cyclobutyl dimers of thymine and thymidine by HPLC and TLC (195) 139

Pyrimidinedione derivatives
Quantitative determination of 1,3-bis(tetrahydro-2-furanyl)-5-fluoro-2,4-pyrimidinedione and its metabolites in visceral tissues by HPLC and GC mass fragmentography (188) 270

Pyrimidine nucleosides
HPLC analysis of cytosine arabinoside and metabolites in biological samples (183) 449

Pyrimidine ribotide
Determination of pyrimidine ribotide and deoxyribotide pools in cultured cells and mouse liver by HPLC (188) 149

Pyrimidines
Quantitative HPLC of nucleosides and bases in human plasma (183) 421

Pyrimidines
Determination of selected pyrimidines, purines and their metabolites in serum and urine by reversed-phase ion-pair chromatography (199) 345

Pyrrolizidine alkaloids
Preparative separation of the pyrrolizidine alkaloids, intermedine and lycopsamine, as their borate complexes (195) 379

Pyrrolizidine alkaloids
Chloranil, a sensitive detection reagent for pyrrolizidine alkaloids on thin-layer chromatograms (195) 407

Pyrrolizidine alkaloids
Specific detection of pyrrolizidine alkaloids on thin-layer chromatograms (195) 412

Pyrrolizidine alkloids
Rapid HPLC isolation of monoesters, diesters and macrocyclic diester pyrrolizidine alkaloids from *Senecio jacobaea* and *Amsinckia intermedia* (192) 474

Q

Quantitative analysis
Comparison of different methods of quantitative analysis employing GLC, illustrated by the determination of water in organic solvents (198) 7

Quantitative structure-activity relationships
R_M values from reversed-phase TLC as parameters of lipophilicity in quantitative structure-activity relationships in four series of arylaliphatic acids (195) 329

Quantitative structure-activity relationships
Use of reversed-phase TLC to asses hydrophobicity in studies of quantitative structure-activity relationships for drugs (196) 391

Quantitative structure-retention relationships
Quantitative structure-retention relationships in cation-exchange chromatography (198) 241

Quaternary alkylammonium ions
Ion-pair chromatography in the low concentration range by use of highly absorbing counter ions. III. HPLC of quaternary alkylammonium ions as ion pairs with naphthalene-2-sulphonate, using a silica support of low surface area (193) 225

Quaternary ammonium compounds
Analysis of quaternary ammonium compounds and basic drugs based on ion-pair adsorption HPLC (186) 683

Quaternary ammonium compounds
GLC of straight chain homologues of alkylbenzyldimethylammonium compounds (200) 216

Quaternary ammonium ion
Reversed-phase ion-pair chromatography with an adsorbing stationary phase and a hydrophobic quaternary ammonium ion in the mobile phase. I. Retention studies with tetrabutylammonium as cationic component (185) 225

Quinazolinones
Determination of nitromethaqualone in blood by electron-capture-GC (182) 435

Quinic acid
Quantitative determination of quinic acid and derivatives by HPLC after derivatization with *p*-bromophenacyl bromide (190) 411

Quinidine
Simple and selective HPLC method for estimating plasma quinidine levels (181) 207

Quinidine
HPLC separation and isolation of quinidine and quinine metabolites in rat urine (181) 219

Quinidine
Simultaneous analysis of disopyramide and quinidine in plasma by HPLC (183) 65

Quinidine
An integrated approach to measurements of quinidine and metabolites in biological fluids (183) 514

Quinidine
Reversed-phase HPLC analysis of cinchona alkaloids in pharmaceuticals (189) 241

Quinine
HPLC separation and isolation of quinidine and quinine metabolites in rat urine (181) 219

Quinine
Reversed-phase HPLC analysis of cinchona alkaloids in pharmaceuticals (189) 241

Quinine-quinidine
High-temperature quantitative glass capillary GC. Analysis of piperine and of quinine-quinidine (199) 105

Quinolinic acid
Determination of urinary quinolinic acid by HPLC (182) 425

Quinones
GC identification and determination of chlorinated quinones formed during chlorination of dihydric phenols with hypochlorite in dilute aqueous solution (200) 137

R

Radioactive sources
Comparative investigation of radioactive sources for the electron capture detector (195) 1

Radioisotope detector
Synchronized accumulating radioisotope detector for radio GC (196) 144

Rauwolfia alkaloids
Chromogenic reactions of *Rauwolfia* alkaloids after separation by TLC (187) 199

Reaction kinetics
Study of the kinetics for a catalyzed reaction using GC (197) 129

Reactor, photochemical
Construction of a simple gas chromatograph with photolytic intersect (189) 119

Rebaudioside
Stevia rebaudiana. I. Use of *p*-bromophenacyl bromide to enhance UV detection of water-soluble organic acids (steviolbioside and rebaudioside B) in HPLC analysis (192) 387

Recycle chromatography
New approach to chromatographic optimization in glass capillary GC (186) 189

Redistribution products
GC with plasma emission spectroscopic detection of Friedel-Crafts catalyzed alkyl group redistribution products among Si, Ge, Sn and Pb atoms (196) 265

Relative molar response factor
Dependence of the relative molar response of isomeric alkanes on molecular structure (in German) (194) 213

Relative retention
Relative retention expressions in chromatography (198) 229

Renal contrast media
Rapid determination of renal contrast media in biological fluids by means of HPLC (182) 492

Residual impurities
GC quantitation of residual impurities in raw material matrixes (193) 207

Resins
Peptide separations on substituted polystyrene resins. Effect of cross-linkage (187) 27

Resins
Chromatography in aqueous solution on styrene-divinylbenzene resins (193) 197

Resins
Separation and characterization of by-product oligomers in epoxy resins by reversed-phase HPLC (193) 243

Resins
Preparation and sorption behaviour of cyclodextrin polyurethane resins (194) 153

Resins
Evaluation of Amberlite XAD-4 for recovery of glucuronides and sulfates from aqueous solution (196) 303

Resins
Chromatographic behaviour of aromatic compounds on anion-exchange resins in various carboxylate forms (198) 267

Resolution
Determination of resolution in gel permeation chromatography (191) 279

Response mechanism
Studies and speculations on an alternative mechanism for the response of direct-current electron-capture detectors (188) 1

Retension mechanism
Retension mechanism in reversed-phase ion-pair chromatography of amines and amino acids on bonded phases (185) 241

Retention behaviour
Electron donor-acceptor properties of urea and its role in charge-transfer chromatography (188) 257

Retention behaviour
Influence of temperature on the retention behaviour of members of homologous series in reversed-phase HPLC (196) 1

Retention-carbon number relationships
Linearity of the retention time *versus* carbon number plot for *n*-alkanes (188) 246

Retention data
GC of some alkenyl- and

cycloalkenylnaphthalenes on Bentone 34, silicone fluid DC 550, and Apiezon L phases (198) 1

Retention data
Linearity of retention data plots for *n*-alkanes on porous polymers in GC (198) 156

Retention data
Stationary phase effects in reversed-phase chromatography. I. Comparison of energetics of retention on alkyl-silica bonded phases (199) 35

Retention data, presentation
Evaluation of procedures for the estimation of dead time (Review) (184) 1

Retention expressions
Relative retention expressions in chromatography (198) 229

Retention index
Relationships between GC retention index and molecular structure (187) 1

Retention index
GC of esters. XIII. Interrelationship of equivalent chain length *(ECL)* and retention index values of fatty esters (187) 297

Retention index system
Different bases for the GC retention index system (192) 107

Retention indices
GLC retention indices of 296 non-drug substances on SE-30 or OV-1 likely to be encountered in toxicological analyses (Review) (184) 185

Retention indices
Determination of Kováts retention indices with a capillary column and electron-capture detection: application to the assay of the enzymatic conversion of 3,4-epoxy-1-butene into diepoxybutane (186) 81

Retention indices
Calculation of GC retention indices of isoalkanes based on a topological approach (188) 297

Retention indices
Determination of the energy of intermolecular interactions in GC from retention indices (190) 1

Retention indices
Relationship between the retention indices of phenols on polar and non-polar stationary phases (196) 217

Retention indices
GC retention data for silyl and acyl derivatives of nucleosides (196) 367

Retention measurements
Effect of temperature on precision of retention measurements in LC (194) 285

Retention mechanism
Mechanism of solute retention in liquid-solid chromatography and the role of the mobile phase in affecting separation. Competition *versus* "sorption" (Review) (184) 363

Retention mechanism
Retention mechanism for reversed-phase ion-pair LC (186) 419

Retention mechanism
Adsorption effects in gas-liquid partition chromatography. Quantitative evaluation (190) 21

Retention mechanisms
Retention mechanisms of tetracyclines on a C_8 reversed-phase material (195) 35

Retention order
Retention behaviour of *o*-, *m*- and *p*-isomers of benzene derivatives on a silica gel hydroxylated surface in LC (195) 205

Retention-structure relationships
Relationships between GC retention index and molecular structure (187) 1

Retention volume
Effect of adsorption on solute retention in GLC (186) 47

Retention volumes, specific
Error in the determination of the specific retention volume. Specific retention volumes of *n*-alkanes on Apolane-87 stationary phase as functions of column temperature (198) 85

Retention volumes, specific
Correlation of GC specific retention volumes of homologous compounds with temperature and methylene number (199) 209

Retinals
HPLC analysis of *cis-trans* stereoisomeric 3-dehydroretinals in the presence of retinal isomers (192) 395

Retinoic acid
GC-MS assay for low levels of retinoic acid in human blood (182) 335

Retinoic acid
HPLC of 13-*cis*-retinoic acid and of endogenous retinol in human plasma (183) 346

Retinol
HPLC of 13-*cis*-retinoic acid and of endogenous retinol in human plasma (183) 346

Retinyl acetate
Preparative TLC of the oxidation products of retinyl acetate (191) 301

Reversed-phase thin-layer chromatography
Problems and applications of reversed-phase TLC (186) 563

R_F values
Correction of R_F values according to Galanos and Kapoulas (in German) (198) 357

Rhamnopyranoside
Preparative LCC of methyl ethers of methyl α-L-rhamnopyranoside (196) 331

Rhodium
GC of volatile metal dialkyldithiophosphates (193) 53

Riboflavin
Rapid method for determination of riboflavin in urine by HPLC (182) 285

Ribonucleic acids, see RNA

Ribonucleotides
Determination of pyrimidine ribotide and deoxyribotide pools in cultured cells and mouse liver by HPLC (188) 149

Ribonucleotides
Analysis of purine ribonucleotides and deoxyribonucleotides in cell extracts by HPLC (198) 435

Ridomil
GC determination of Metalaxyl (Ridomil) residues in tobacco (200) 313

Ridomil
Estimation of the systemic fungicide Ridomil by TLC (200) 317

RNA
Fractionation of ^{32}P-labeled animal RNA on Sephadex-Sepharose columns (197) 236

RNA, transfer
HPLC and MS of transfer RNA bases for isotopic abundance (194) 205

Rubidium chloride
Adsorption chromatography capillary columns. I. Rubidium chloride (196) 237

Rubreserine
Determination of pilocarpine, physostigmine, its degradation product rubreserine and preservatives by HPLC (198) 529

S

Saccharide distribution
A rapid and high-resolution method to determine the composition of corn syrups by LC (187) 381

Saccharides, see also sugars

Saccharides
Analysis of cyclodextrins by HPLC (189) 98

Saccharides
Separation of saccharides and their anomers by HPLC (191) 121

Saccharides
LC of non-reducing oligosaccharides: a new detection principle (191) 313

Saccharides
Improved separation of anomers of saccharides by HPLC on macroreticular anion-exchange resin in the sulphate form (192) 452

Saccharides
Separation of monosaccharides as their alditol acetates by capillary column GLC (194) 410

Saccharin
HPLC determination of impurities in commercial saccharin (187) 249

Salicylic acid
Rapid determination of indomethacin and salicylic acid in serum by means of reversed-phase LC (181) 385

Salicylic acid
Specific and sensitive method for the determination of aspirin and salicylic acid in plasma using reversed-phase HPLC (181) 473

Salsoline
Analysis of salsolinol and salsoline in biological samples using deuterium-labelled internal standards and GC-MS (183) 17

Salsolinol
Analysis of salsolinol and salsoline in biological samples using deuterium-labelled internal standards and GC-MS (183) 17

Sample application
Contact spotting. A new approach to sample application in TLC (186) 555

Sample capacity
Isotherm linearity and sample capacity in LC (193) 181

Sample concentration
On-line trace enrichment in HPLC using a pre-column (185) 483

Sample concentration
Chromatographic analysis of major trace components in solvents with concentration prior to sample injection (191) 309

Sample concentration
JADO carbon as an adsorbent in sampling columns (193) 137

Sample dispersion
Early phases of the dispersion of a sample injected in Poiseuille flow (186) 353

Sample handling
GC detection and structure analysis of volatile pheromones in insects (186) 605

Sample injection
Should the septum part of vaporizing injectors be kept at lower temperatures? (198) 64

Sample introduction
Versatile all-glass splitless sample-introduction

system for trace analysis by capillary GC (186) 167

Sample size
Influence of mobile phase composition and nature of sample on retention data from HPLC (191) 81

Sample volume overload
Preparative LC: sample volume overload (186) 457

Sampling method
High-temperature quantitative glass capillary GC. Analysis of piperine and of quinine-quinidine (199) 105

Sapogenins, steroidal
Sintered TLC (Review) (184) 37

Saponins
GLC method for determination of ginsenosides in *Panax ginseng* (196) 121

Scintillation reagents
GLC retention indices of 296 non-drug substances on SE-30 or OV-1 likely to be encountered in toxicological analyses (Review) (184) 185

Secretin
Perfluoroalkanoic acids as lipophilic ion-pairing reagents in reversed-phase LC of peptides including secretin (195) 181

Sedative drugs
Screening procedure for therapeutic benzodiazepines by HPLC of their benzophenones (189) 94

Sedative drugs
Analysis of trace amounts of barbiturates in saliva (190) 113

Sedative drugs
Mass screening and confirmation of methaqualone and its metabolites in urine by radioimmunoassay-TLC (190) 129

Sedative drugs
GC separations of dialkyl barbiturate derivatives (192) 212

Sedative drugs
Glass capillary column GC of barbiturates after flash-heater derivatization with dimethylformamide dimethylacetal (192) 363

Sedative drugs
Direct derivatization and GC determination of barbiturates in autopsy liver tissues (195) 349

Sedative drugs
Use of dual-wavelength UV detection in HPLC for the identification of barbiturates (200) 271

Sediment analysis
Elucidation of geomatrices by laser pyrolysis-GC and pyrolysis-MS (186) 831

Sediment analysis
GLC-MS studies on extracts and distillates from Posidonomia shales (186) 843

Segmented-flow liquid chromatography
Coated-open-tubular chromatography with flow segmentation. I. Theory (185) 43

Segmented-flow liquid chromatography
Coated-open-tubular chromatography with flow segmentation. II. Experimental study and optimization for size-exclusion separation (185) 57

Selectivity
Effect of alkyl chain length of the stationary phase on retention and selectivity in reversed-phase LC. Participation of solvent molecules in the stationary phase (199) 327

Selectivity optimization
Optimization of solvent strength and selectivity for reversed-phase LC using an interactive mixture-design statistical technique (199) 57

Sensitivity setting
Effect of the sensitivity setting of a katharometer on response factors at low concentrations (197) 121

Sensitization, selective
Selective electron-capture sensitization (186) 3

Septum bleed
Decreased accuracy in isotachophoretic analysis due to septum bleed (188) 262

Sequential isocratic step liquid chromatography
"Sequential isocratic step" LC: a high-throughput, problem-solving approach with sensitive, near-universal detection for wide polarity mixtures (199) 219

Serotonin
Simple and sensitive procedure for the assay of serotonin and catecholamines in brain by HPLC using fluorescence detection (182) 277

Serotonin
Comparison of LC-electrochemical and GC-MS methods for brain dopamine and serotonin (183) 483

Serotonin
In vivo studies of the tryptophan-5-hydroxylase system. Quantitation of serotonin and tryptamine using GC-mass fragmentography (199) 171

Serum albumin
Covalent coupling of cholic acid to aminohexylamino-Sepharose 4B and its use in affinity chromatography of serum albumin (187) 409

Serum profiles
Comparison of HPLC serum profiles of humans and dogs (181) 169

Serum proteins
Simple method for the purification of C1q, a subcomponent of the first component of

complement by affinity chromatography using IgG-Sepharose (189) 428

Servalytes
Gel chromatographic comparison of the molecular weight distributions of Ampholine, Servalyte and Pharmalyte carrier ampholytes used in isoelectric focusing (198) 301

Sesquiterpene lactones
Visualization reagents for sesquiterpene lactones and polyacetylenes on thin-layer chromatograms (189) 187

Sesquiterpene lactones
Application of GC to the analysis of sesquiterpene lactones from *Lactarius (Russulaceae)* mushrooms (190) 466

Sesquiterpenes
Study of three types of essential oil of *Valeriana officinalis* L. s. l by combined GC-negative ion chemical ionization MS (190) 321

Sesquiterpenoids
Photodensitometric determination of "PR-toxin", metabolite of *Penicillium roqueforti* (in French) (198) 481

Shikimic acid
Quantitative determination of quinic acid and derivatives by HPLC after derivatization with *p*-bromophenacyl bromide (190) 411

Side-chain determination
Quantitative determination of alkylamino side-chains coupled to agarose beads. Comparison of methods (193) 451

Silatranes
Chromatography of organometallic and organometalloidal derivatives of amino alcohols. II. GC evidence for transannular bonds in solutes (200) 105

Silica gel
Characterization of HPLC stationary phases: the nature of silica gel bonded material (190) 331

Silica gel photoluminescence
UV-induced silica gel GF photoluminescence studied by liquid scintillation spectrophotometry (195) 281

Silicate anions
Gel permeation chromatographic separation and identification of trimethylsilyl derivatives of silicate anions (197) 59

Silicic acid
Gel chromatographic study of the polymerization of silicic acid in acid solutions (198) 429

Silylating reagent
tert.-Butylpentafluorophenylmethyl-chlorosilane as a reagent for the formation of hydrolytically stable alkylsilyl derivatives with electron-capturing properties (187) 331

Sintered thin-layer chromatography
Sintered TLC (Review) (184) 37

Size characterization
Steric field-flow fractionation as a tool for the size characterization of chromatographic supports (185) 261

Size distribution
Size and charge distribution of macromolecules in living systems (193) 1

Soda-lime glass columns
Studies of open-tubular micro-capillary LC. IV. Soda-lime glass columns treated with alkaline solution (185) 73

Sodium azide
Presence of sodium azide during acid hydrolysis of protein samples causes the destruction of tyrosine, phenylalanine and histidine (189) 106

Sodium dodecyl sulphate
Anomalous partition behaviour of sodium dodecyl sulphate monomer in Sephadex gels with high dextran concentrations. Possible role of water of hydration in the gel phase (193) 345

Solasodine
Rapid quantitative analysis of solasodine, solasodine glycosides and solasodiene by HPLC (187) 87

Solid-phase reagents
Solid-phase reagents for LC detection (185) 497

Solute-modifier interactions
Role of organic modifier sorption on retention phenomena in reversed-phase LC (199) 259

Solute-nitrile group interactions
GLC determination of equilibrium constants of interactions between solutes and free nitrile groups (in French) (193) 9

Solute profiles
Chromatographic peak shape. III. Influence of kinetics parameters on solute profiles (186) 117

Solute-solvent interactions
Mechanism of solute retention in liquid-solid chromatography and the role of the mobile phase in affecting separation. Competition *versus* "sorption" (Review) (184) 363

Solute-solvent interactions
Liquid adsorption chromatography with mixed mobile phases. IV. A new equation for the capacity ratio involving the solute-solvent interactions (188) 27

Solute-solvent interactions
Solute-solvent interactions on the surface of silica gel. III. Multilayer adsorption of water on the surface of silica gel (196) 193

Solute-solvent interactions
Solute-solvent interactions on the surface of reversed phases. I. Stationary phase

interactions and their dependence on bonding characteristics (197) 11

Solvent composition-retention relationships
Mobile phase effects in reversed-phase chromatography. I. Concomitant dependence of retention on column temperature and eluent composition (185) 99

Solvent composition-retention relationships
Mobile phase effects in reversed-phase chromatography. III. Changes in conformation and retention of oligo(ethylene glycol) derivatives with temperature and eluent composition (185) 129

Solvent effect
Effect of solvents on the resolution of neutral lipids on chromarods (198) 279

Solvent eluotropic scale
LSC on Amberlite XAD-2 and other styrene-divinylbenzene adsorbents. I. Development of a solvent eluotropic scale (189) 145

Solvent participation
Effect of alkyl chain length of the stationary phase on retention and selectivity in reversed-phase LC. Participation of solvent molecules in the stationary phase (199) 327

Solvent polarity scale
Fundamental, predictive approach to dipole-dipole interactions based on the dipole moment of the solute and the dielectric constant of the solvent (194) 105

Solvent programmer
Single-pump solvent programmer for HPLC using synchronized valve switching (185) 515

Solvent purity test
Detection of trace organic impurities in binary solvent systems: a solvent purity test (188) 193

Solvent strength optimization
Optimization of solvent strength and selectivity for reversed-phase LC using an interactive mixture-design statistical technique (199) 57

Solvent systems
Design of binary solvent systems for LSC on silica gel columns using mono- and difunctional steroids as solutes (200) 85

Sorbents
Polyfunctional polymer sorbents (186) 227

Sorbents
Evaluation of chromatographic sorbents used in air pollution studies (186) 811

Sorbents
Modification of the Rohrschneider method for the classification of the polarity of polymeric sorbents based on methacrylic acid esters (190) 13

Sorbents
Electron-microscopic investigation of the sorbents used in GLC (193) 132

Sorbents
Reactive polymers. XXXI. Thermally treated polymeric sorbents based on glycidylmethacrylate copolymers (194) 297

Sorbents
Structure and sorption properties of porous copolymers of styrene, acrylonitrile and divinylbenzene (196) 75

Sorbents
Model studies of the reactions and non-specific adsorptive properties of cyanogen bromide-activated macroporous agarose (196) 379

Sorbents
Stability of ion exchangers against treatment with nitric acid and radiation. II. Change in sorption properties of Dowex A-1, Chelex 100 and Wof MC 50 after treatment with nitric acid (in German) (198) 257

Sorbents
Retention behavior of chlorinated benzenes, chlorinated phenols, Aldrin, and Aroclor 1242 on Tenax-GC, Chromosorb 101, Florisil, and Carbopack C HT (200) 115

Sorption effect
Influence of sorption on the shape of chromatographic elution curves (198) 377

Soybean protease inhibitors
Affinity chromatographic interactions of proteases with low-molecular-weight soybean protease inhibitors (195) 385

Specific surface areas
Separation of C_6 cyclic hydrocarbons before the inert peak on microporous silica: a case of negative adsorption (199) 23

Spheron
Ion-exchange derivatives of Spheron. III. Carboxylic cation exchangers (192) 159

Sphingolipids
Biochemistry of sphingolipids. XXVII. Thin-layer microchromatography of DNP derivatives of long-chain bases on Silufol $_{UV\ 254}$ sheets (190) 498

Sphingomyelin
Comparative determination of plasma phospholipids by automated GLC and manual colorimetric phosphorus methods (182) 1

Sphingomyelin
Miniature two-dimensional TLC separation of lecithin and sphingomyelin (182) 311

Splitless sample introduction
Versatile all-glass splitless sample-introduction system for trace analysis by capillary GC (186) 167

Sporidesmin
Separation and purification of metabolites of the mycotoxin, sporidesmin, by HPLC (200) 288

Spotting
Contact spotting. A new approach to sample application in TLC (186) 555

Spray reagents
Detection of thiols on thin-layer chromatograms with 3,5-di-*tert.* -butyl-1,2-benzoquinone-iron(III) chloride (187) 418

Spray reagents
Visualization reagents for sesquiterpene lactones and polyacetylenes on thin-layer chromatograms (189) 187

Spray reagents
Optical brighteners as TLC detection reagents for glycoalkaloids and steroid alkaloids in *Solanum* species. II. Blankophor BA 267%, BBU neu and KU, and Tinopal CBS-X and 5 BMS-X (189) 406

Spray reagents
Use of ninhydrin as a spray reagent for the detection of some basic drugs on thin-layer chromatograms (195) 133

Spray reagents
Chloranil, a sensitive detection reagent for pyrrolizidine alkaloids on thin-layer chromatograms (195) 407

Spray reagents
Alkaline chloramine-T reagent for the detection of phenolic compounds on thin-layer plates (196) 309

Spray reagents
Fluorescence detection of secondary amines on thin-layer plates using a fluorogenic reaction with fluorescamine (200) 324

Spreader
New spreader for the preparation of gradient layers in TLC (189) 293

Squalane
Study of mixed solvents: squalane-stearonitrile. I. Comparison of results by GC and surface tensions (in French) (188) 315

Squalane
Presence of squalane in urban aquatic environments (194) 199

Stability constants
Measurement of association constants for complexes by reversed-phase HPLC (186) 371

Staining
Trypan blue staining of proteins fractionated by acrylamide disc gel electrophoresis (189) 410

Staining
Quantitative staining of submicrogram amounts of histone and high-mobility group proteins on sodium dodecylsulphate-polyacrylamide gels (200) 200

Stannatranes
Chromatography of organometallic and organometalloidal derivatives of amino alcohols. II. GC evidence for transannular bonds in solutes (200) 105

Stationary phase activity
Simplified HPLC phase characterization: particle size, surface area, mean pore diameter, amount of bonded phase and activity (186) 435

Stationary phase characterization
Simplified HPLC phase characterization: particle size, surface area, mean pore diameter, amount of bonded phase and activity (186) 435

Stationary phase classification
Elimination of the reference stationary phase in classification schemes using the molecular retention index (199) 199

Stationary phase effects
Stationary phase effects in reversed-phase chromatography. I. Comparison of energetics of retention on alkyl-silica bonded phases (199) 35

Stationary phase polarity
Modification of the Rohrschneider method for the classification of the polarity of polymeric sorbents based on methacrylic acid esters (190) 13

Stationary phases
High-performance exclusion chromatography of water-soluble polymers with chemically bonded stationary phases (185) 305

Stationary phases
Ligand-exchange chromatography of enantiomeric amino acids on copper-loaded chiral bonded silica gel and of amino acids on copper(II)-modified silica gel (185) 345

Stationary phases
Evaluation of a thermally stable hydrocarbon as a non-polar base stationary phase for use in Rohrschneider-McReynolds-type schemes (186) 89

Stationary phases
Utility of high-temperature thermotropic liquid crystals as stationary phases for novel GLC separations (186) 237

Stationary phases
Poly(α-olefins): novel non-polar stationary phases for GC (186) 259

Stationary phases
Comprehensive study of porous polymeric stationary phases for chromatography (186) 317

Stationary phases
LC resolution of enantiomers on normal-phase chiral amide-bonded silica gel. Retentions of optically active α-amino acid derivatives on N-acyl homologues of L-valylaminopropylsilanized silica phases (186) 543

Stationary phases
Synthesis and properties of a novel chiral stationary phase for the resolution of amino acid enantiomers (186) 611

Stationary phases
Reversed-phase ion-pair HPLC of drugs and related compounds using underivatized silica as the stationary phase (186) 705

Stationary phases
Effect of the support surface on the structure of the film of liquid crystalline stationary phase (188) 107

Stationary phases
Coordination polymers as adsorbents and stationary phases in GC. VI. Separation properties of alkaline-earth metal di-*n*-hexylphosphinates (188) 323

Stationary phases
Synthesis and GC use of metal-containing surfaces (189) 13

Stationary phases
Silica gel modified with phthalocyanines as a stationary phase in gas-solid chromatography (189) 25

Stationary phases
LC on metal oxides with silica geometry (189) 139

Stationary phases
4-(*p*-methoxycinnamyloxy)-4'-methoxy-azobenzene: a nematic liquid crystal for the GLC analysis of the stereochemistry of lepidopterous sex pheromones and related unsaturated fatty alcohols and derivatives (190) 35

Stationary phases
Use of Carbowax 20M glass capillary columns below their lower temperature limit (190) 283

Stationary phases
Calcium complex grease as a low-polarity stationary phase (190) 291

Stationary phases
Separation range and separation efficiency in high-speed gel filtration on TSK-GEL SW columns (190) 297

Stationary phases
Characterization of HPLC stationary phases: the nature of silica gel bonded material (190) 331

Stationary phases
Comparison of linear and branched non-polar chemically bonded stationary phases in HPLC (191) 71

Stationary phases
Liquid crystals as the stationary phase in GC. Adsorption phenomena caused by an electric field applied across the column (192) 89

Stationary phases
Solution behaviour of liquid crystal phases in capillary GC. II. Solution properties of smectic A-, B- and C-modifications (in German) (192) 97

Stationary phases
Studies on open-tubular microcapillary LC. VI. Styrene-divinylbenzene copolymer stationary phase (192) 127

Stationary phases
Broad spectrum resolution of optical isomers using chiral HPLC bonded phases (192) 143

Stationary phases
Ion-exchange derivatives of Spheron. III. Carboxylic cation exchangers (192) 159

Stationary phases
Use of chemically bonded stationary phases in HPTLC. II (192) 331

Stationary phases
Separation of xylenol isomers on nitrofluorenes and nitro-9-fluorenones by GLC. I. Mixed columns and effect of temperature on retention behaviour (193) 300

Stationary phases
Comparison of TSK-GEL PW type and SW type in high-speed aqueous gel-permeation chromatography (193) 311

Stationary phases
GLC separation of bile acids and steroids on a nematic liquid crystal (193) 444

Stationary phases
Quantitative determination of alkylamino side-chains coupled to agarose beads. Comparison of methods (193) 451

Stationary phases
Empirical correlation of layer thickness and support pore size in "chemically bonded" phases (195) 25

Stationary phases
Liquid crystals. I. Synthesis and application as stationary phases in GLC (195) 309

Stationary phases
N-Docosanoyl-L-valine-2(2-methyl)-*n*-heptadecylamide as a stationary phase for the resolution of optical isomers in GLC (195) 317

Stationary phases
Role of the chain length of chemically bonded phases and the retention mechanism in reversed-phase LC (196) 21

Stationary phases
GC analysis of light hydrocarbons for the characterization of Porapak columns (196) 245

Stationary phases
Applications of Dexsil and other GC porous-layer open tubular columns to metal compound separations and pyrolysis (196) 403

Stationary phases
GC study of grafted alkyl chains used as stationary phases (196) 481

Stationary phases
Solute-solvent interactions on the surface of reversed phases. I. Stationary phase interactions and their dependence on bonding characteristics (197) 11

Stationary phases
GC and LC on silica-supported chromia layers (197) 21

Stationary phases
GC study of solution and adsorption of hydrocarbons on glycols. I. Diethylene glycol and triethylene glycol (197) 135

Stationary phases
Separation behaviour of some isomeric organic compounds on sugars, sugar alcohols and their mixed phases by GLC (197) 251

Stationary phases
GC stationary phases analysed by capillary GC (198) 176

Stationary phases
Stationary phases for HPLC (198) 421

Stationary phases
Separation of C_6 cyclic hydrocarbons before the inert peak on microporous silica: a case of negative adsorption (199) 23

Stationary phases
Applications of surface-modified porous silicas to glass capillary column preparation (199) 97

Stationary phases
Effect of alkyl chain length of the stationary phase on retention and selectivity in reversed-phase LC. Participation of solvent molecules in the stationary phase (199) 327

Stationary phases
"Bonded phases" from cyclic organosilicons for GC and LC (200) 3

Stationary phases
Properties of liquid crystalline cyanoazoxybenzene alkyl carbonates as stationary phases in GC (200) 65

Stationary phases
GC evidence for phase transitions in very compact octadecyl bonded silicas (200) 95

Stationary phases, optically active
Structural requirements for enantioselectivity in GC of chiral α-hydroxy acids (200) 189

Stationary phases, optically active
Resolution of amino acid enantiomers by glass capillary GC on easily prepared optically active stationary phases (200) 195

Stearonitrile
Study of mixed solvents: squalane-stearonitrile. I. Comparison of results by GC and surface tensions (in French) (188) 315

Steric field-flow fractionation
Steric field-flow fractionation as a tool for the size characterization of chromatographic supports (185) 261

Steroidal alkaloids
HPLC of *Solanum* and *Veratrum* alkaloids (198) 363

Steroidal glucosiduronic esters
Liquid ion exchanges in PC of steroidal glucosiduronic esters (189) 33

Steroid alkaloids
Optical brighteners as TLC detection reagents for glycoalkaloids and steroid alkaloids in *Solanum* species. II. Blankophor BA 267%, BBU neu and KU, and Tinopal CBS-X and 5 BMS-X (189) 406

Steroidal sapogenins
Sintered TLC (Review) (184) 37

Steroid dehydrogenase
Efficient purification of a microbial steroid-1-dehydrogenase by electrophoretic desorption from the affinity matrix on a preparative scale (189) 279

Steroid digitonides
Fractionation of steroid digitonides by TLC and CC on silica gel (188) 445

Steroid glucuronides
Separation of steroid glucuronides by reversed-phase LCC (194) 80

Steroid hormones
Separation of metabolites of testosterone hydroxylation by HPLC (196) 494

Steroids
Enzymatic color development of 3α-hydroxysteroids on thin-layer chromatograms for determination of excretion pattern of 3α-hydroxysteroids in patients with some adrenogenital syndrome (182) 430

Steroids
Rapid quantitative assay of plasma 11-deoxycortisol and cortisol by HPLC for use in the metyrapone test (183) 87

Steroids
Quantitative analysis of corticosteroids in adrenal cell cultures by capillary column GC combined with MS (186) 581

Steroids
Strong non-polar cation exchangers for the separation of steroids in mixed chromatographic systems (186) 725

Steroids
HPLC of the reduction products of progesterone (190) 169

Steroids
Pregnenolone separation from cholesterol using Sephadex LH-20 minicolumns (190) 256

Steroids
GC of dimethylalkylsilyl ether derivatives of bile acid methyl esters (190) 339

Steroids
HPLC of the topical anti-inflammatory steroid halcinonide (190) 481

Steroids
HPLC and GLC of some norgestrel intermediates. Physical properties of the isolated *syn*- and *anti*-isomers of oximes (191) 145

Steroids
Capillary GC studies of cholesterol biosynthesis in rats treated with EGYT-1299 (191) 261

Steroids
GC-MS properties of boronate esters of 24*R*,25-dihydroxycholecalciferol (192) 434

Steroids
Separation of C_{21}-, C_{19}- and C_{18}-steroids on Sephadex LH-20 microcolumns (193) 322

Steroids
Quantitative HPLC determination of Δ^4-3-ketosteroids in adrenocortical extracts (193) 413

Steroids
GLC separation of bile acids and steroids on a nematic liquid crystal (193) 444

Steroids
Improved HPLC separation of decomposition products of prednisolone by adding sulphite to the mobile phase (193) 480

Steroids
Practical aspects of the preparation and chromatography of the trimethylsilyl ethers of ecdysteroids (194) 43

Steroids
Intermediate- and large-scale reversed-phase preparative HPLC on an axially compressed column: a facile, quantitative separation of 7*α*- and 7β-methyl-17β-acetoxy-3-oxoandrost-4-enes (194) 239

Steroids
Comparison of HPLC and GC for the analysis of ecdysteroids (194) 343

Steroids
HPLC separation and analysis of steroidal constituents of two solanaceous plants (195) 425

Steroids
Concentration-dependent changes in Methylene Units in steroid analysis (in German) (197) 232

Steroids
Reversed- and normal-phase HPLC of 18-hydroxylated steroids and their derivatives. Comparison of selectivity, efficiency and recovery from biological samples (198) 23

Steroids
Reversed-phase HPLC separation and analysis of some physalins (13,14-*seco*-16,24-cyclo-steroids) (198) 203

Steroids
Design of binary solvent systems for LSC on silica gel columns using mono- and difunctional steroids as solutes (200) 85

Steroids
Separation of urinary steroids by GLC on packed and mixed-bed columns (200) 221

Sterols
Simple mass chromatographic procedure for the detection and identification of sterols using derivative correlations (189) 267

Sterols
Separation of sterols using digitonin-impregnated TLC plates (194) 429

Sterols
Metabolic profiles of penaeid shrimp: dietary lipids and ovarian maturation (195) 359

Sterols
Marine sterols. XIII. Evaluation of glass support-coated open tubular columns, prepared by the mercury-plug dynamic technique, for marine sterol analysis (198) 111

Steviolbioside
Stevia rebaudiana. I. Use of *p*-bromophenacyl bromide to enhance UV detection of water-soluble organic acids (steviolbioside and rebaudioside B) in HPLC analysis (192) 387

Stigmasterol
Fractionation of steroid digitonides by TLC and CC on silica gel (188) 445

Stilbene derivatives
Simple rapid TLC method for the detection of optical brighteners in polymers (194) 434

Stimulant drugs
Determination of caffeine in serum by straight-phase HPLC (182) 121

Stimulant drugs
Quantitation of doxapram in blood, plasma and urine (182) 191

Stimulant drugs
Microdetermination of stimulant drugs in urine by HPLC (196) 334

Stimulant drugs
Qualitative screening procedure for the detection of benzoyl ecgonine in the urine of greyhounds (196) 337

Structure-activity relationships
R_M values from reversed-phase TLC as parameters of lipophilicity in quantitative structure-activity relationships in four series of arylaliphatic acids (195) 329

Structure-activity relationships
Reversed-phase HPLC. I. Relationship between R_M values and Hansch's π parameters for a series of phenols (197) 43

Structure-response relationships
Structure-response relationship of GC-flame photometric detection to some organophosphorus compounds (189) 331

Structure-response relationships
Electron capture GC of sulphonamides. Effects of structure and temperature on detector response (189) 351

Structure-retention correlations
Identification of *n*-hepta- and *n*-octadienes by high-resolution GC using structure-retention correlations and MS (191) 187

Structure-retention relationships
Relation of GC behaviour to the chemical structure of pyrido[1,2*a*]pyrimidine derivatives (191) 245

Structure-retention relationships
Quantitative structure-retention relationships in cation-exchange chromatography (198) 241

Styrene
Structure and sorption properties of porous copolymers of styrene, acrylonitrile and divinylbenzene (196) 75

Styrene
Determination of styrene in foods by reversed-phase HPLC (198) 198

Styrene-divinylbenzene adsorbents
LSC on Amberlite XAD-2 and other styrene-divinylbenzene adsorbents. I. Development of a solvent eluotropic scale (189) 145

Styrene-divinylbenzene copolymer
Studies on open-tubular microcapillary LC. VI. Styrene-divinylbenzene copolymer stationary phase (192) 127

Styrene-divinylbenzene resins
Chromatography in aqueous solution on styrene-divinylbenzene resins (193) 197

Styrene monooxygenase
Improved GC method for measuring phenylethylene glycol. Application to the determination of styrene monooxygenase and epoxide hydrase activities (188) 400

Styrene oxide
Quantitative determination of styrene-7β-oxide in blood by combined GC-multiple ion detection mass fragmentography (196) 314

Sugar alcohols
Estimation of neutral sugars and sugar alcohols in biological fluids by GLC (Review) (184) 457

Sugars, see also saccharides

Sugars
Estimation of neutral sugars and sugar alcohols in biological fluids by GLC (Review) (184) 457

Sugars
Relative effectiveness of ion-exchange and lead acetate precipitation methods in isolating pear sugars and acids (190) 452

Sugars
Separation and quantitative determination of fructose as the O-methyloxime by GLC using glass capillary columns (192) 446

Sugars
Thermal degradation of sugars in a GC injection port (193) 83

Sugars
Fluorodensitometric microdetermination of reducing sugars on thin-layer chromatograms (193) 333

Sugars
Separation of partially methylated sugars by reversed-phase HPLC (196) 171

Sulfinpyrazone
HPLC analysis of sulfinpyrazone and its metabolites in biological fluids (181) 417

Sulpha drugs
Determination of sulfanilic acid in the presence of sulfanilamide and some sulfa drugs by reversed-phase ion-pair HPLC (196) 498

Sulphamoylbenzoic acid
HPLC determination of impurities in commercial saccharin (187) 249

Sulphanilamide
Determination of sulfanilic acid in the presence of sulfanilamide and some sulfa drugs by reversed-phase ion-pair HPLC (196) 498

Sulphanilic acid
Determination of sulfanilic acid in the presence of sulfanilamide and some sulfa drugs by reversed-phase ion-pair HPLC (196) 498

Sulphate conjugates
HPLC of glucuronide and sulfate conjugates using ion-pair chromatography (195) 431

Sulphates
Evaluation of Amberlite XAD-4 for recovery of glucuronides and sulfates from aqueous solution (196) 303

Sulphates, organic
Sintered TLC (Review) (184) 37

Sulphilimines
HPLC of diastereomeric sulphoxides and sulphilimines (198) 207

Sulphite liquor
Determination of furfural in spent sulfite liquor by GC (194) 228

Sulphocysteine
Separation of S-sulphocysteine and related compounds by anion-exchange chromatography and electrophoresis (188) 442

Sulphonamides
Quantitation of non-halogenated compounds by electron-capture GC (Review) (184) 313

Sulphonamides
Electron capture GC of sulphonamides. Effects of structure and temperature on detector response (189) 351

Sulphonamides
Improved GC method for the quantitative determination of secondary amines as sulphonamides formed by reaction with benzenesulphonyl chloride (190) 462

Sulphonic acids
Reversed-phase LC of aromatic sulphonic acids and other strongly polar compounds without addition of an ion-pairing counter-ion (197) 181

Sulphonylureas
Electron-capture GC of plasma sulphonylureas after extractive methylation (181) 17

Sulphoxides
HPLC of diastereomeric sulphoxides and sulphilimines (198) 207

Sulphur
Discharge of elemental sulfur and its distribution in surficial sediments in the region of the Buccaneer oil field (199) 161

Sulphur hexafluoride
Adsorption-desorption GC-IR determination of trace disulfur decafluoride in sulfur hexafluoride (193) 265

Supports
Steric field-flow fractionation as a tool for the size characterization of chromatographic supports (185) 261

Supports
Preparation of a porous microparticulate anion-exchange chromatography support for proteins (185) 375

Supports
Interactions of proteins with bonded-phase ion exchangers (185) 393

Supports
Non-bonded affinity chromatography: a biospecific oil-soluble coated support for concanavalin A (188) 61

Supports
Polyacrylamide-boronate beads saturated with biomolecules: a new general support for affinity chromatography of enzymes (189) 225

Supports
Bis(octadecyl phosphate)zirconium(IV). Novel support for reversed-phase chromatography (190) 145

Supports
Use of volcanic slag as a support in GC (191) 323

Support surface
Effect of the support surface on the structure of the film of liquid crystalline stationary phase (188) 107

Surface-active agent effect
Combined effects of pH and surface-active-ion concentration in reversed-phase LC (199) 307

Surface analysis
Glass surface analytical studies in the preparation of open tubular columns for GC (199) 355

Surface area
Simplified HPLC phase characterization: particle size, surface area, mean pore diameter, amount of bonded phase and activity (186) 435

Surface treatments
Preparation of glass capillary columns for GC (Review) (184) 235

Surfactants
Improved HPLC analysis of ethylene oxide condensates by their esterification with 3,5-dinitrobenzoyl chloride (187) 261

Surfactants
Analysis of dipeptide mixtures by the combination of ion-pair reversed-phase HPLC and GC-MS techniques (197) 31

Surfactants
Hydrophobic chromatography with dynamically coated stationary phases. III. Non-ionic surfactant effects (198) 397

Switching systems
Information content of multi-dimensional switching systems in GC (186) 129

Sympathicomimetic drugs
Simplified LC-electrochemical determination of norepinephrine and dopamine in rat brain (181) 453

Sympathicomimetic drugs
Simple and sensitive procedure for the assay of serotonin and catecholamines in brain by HPLC using fluorescence detection (182) 277

Sympathicomimetic drugs
Simple method for isolating free urinary catecholamines by boric acid gel chromatography (183) 78

Sympathicomimetic drugs
Investigation of the performance of a HPLC system with an electrochemical detector (183) 261

Sympathicomimetic drugs
Semi-automated fluorimetric method for the estimation of urinary catecholamines using HPLC (183) 295

Sympathicomimetic drugs
Analysis of perfluoroacyl derivatives of ephedrine, pseudoephedrine and analogues by

GC and MS. I. Trifluoroacetyl derivatives (190) 53

Sympathicomimetic drugs
GC of epinephrine and norepinephrine after derivatization with chloroformates in aqueous media (190) 347

T

Tamoxifen
Paired-ion chromatographic analysis of tamoxifen and two major metabolites in plasma (181) 41

Taurine
TLC of citric acid cycle intermediates, lactic acid, and the amino acid taurine (189) 286

Taurine conjugates
Rapid determination of glycine- and taurine-conjugated bile acids in human bile by HPLC (181) 177

Taxonomy, experimental
Experimental taxonomy of *Biomphalaria* (gastropoda: planorbidae). I. Methods for experimental taxonomic studies on *Biomphalaria* carried out by horizontal starch gel electrophoresis and staining of twelve enzymes (188) 169

Temperature effect
Effect of temperature on precision of retention measurements in LC (194) 285

Temperature effect
Influence of temperature on the retention behaviour of members of homologous series in reversed-phase HPLC (196) 1

Temperature effect
Influence of temperature on separation processes in adsorption LC systems (196) 11

Temperature effect
Error in the determination of the specific retention volume. Specific retention volumes of *n*-alkanes on Apolane-87 stationary phase as functions of column temperature (198) 85

Temperature effects
Systematic approach to optimizing resolution in reversed-phase LC with emphasis on the role of temperature (185) 153

Temperature effects
Electron capture GC of sulphonamides. Effects of structure and temperature on detector response (189) 351

Temperature-retention relationships
Mobile phase effects in reversed-phase chromatography. I. Concomitant dependence of retention on column temperature and eluent composition (185) 99

Temperature-retention relationships
Mobile phase effects in reversed-phase chromatography. III. Changes in conformation and retention of oligo(ethylene glycol) derivatives with temperature and eluent composition (185) 129

Temperature selection
Influence of different carrier gases and temperatures on the overall performance of a packed column (186) 249

Tenuazonic acid
HPLC determination of major mycotoxins produced by *Alternaria* molds (194) 89

Terpenoids
Applications of reversed-phase chromatography and nephelometric detection to analysis of non-polar mixtures at microgram level (187) 111

Tertiary amines
Syringe and column adsorption of tertiary amines in GC (196) 489

Testosterone
Separation of metabolites of testosterone hydroxylation by HPLC (196) 494

Tetrabutylammonium
Reversed-phase ion-pair chromatography with an adsorbing stationary phase and a hydrophobic quaternary ammonium ion in the mobile phase. I. Retention studies with tetrabutylammonium as cationic component (185) 225

Tetracycline
LC determination of tetracycline in plasma and urine (185) 583

Tetracyclines
Mechanism of reversed-phase separation of tetracyclines by HPLC (186) 763

Tetracyclines
Rapid analysis of doxycycline from biological samples by HPLC (187) 439

Tetracyclines
Reversed-phase ion pair chromatography of oxytetracycline, epioxytetracycline and anhydrooxytetracycline (190) 486

Tetracyclines
Retention mechanisms of tetracyclines on a C_8 reversed-phase material (195) 35

Tetrahydrocannabinol
Simple, rapid method for extraction of urinary cannabinoids by LSC (196) 342

Tetrahydrocannabinol metabolites
Combined HPLC and radioimmunoassay method for the analysis of Δ^9-

tetrahydrocannabinol metabolites in human urine (186) 595

Tetraiodothyronine
Separation of enantiomeric iodinated thyronines by LC of diastereomers (198) 471

Tetrapyrroles
Separation and characterization of bilirubin conjugates by HPLC (182) 100

Tetrapyrroles
Accurate and sensitive analysis of ethyl anthranilate azopigments from bile by reversed-phase HPLC (182) 105

Tetrazolium derivatives
Separation of tetrazolium derivatives by reversed-phase HPLC (196) 185

Theophylline
New rapid assay of theophylline in plasma by isotachophoresis (181) 147

Theophylline
Micro estimation of plasma theophylline by GLC with on-column butylation and nitrogen-specific detection (182) 232

Therapeutic drugs
Analysis of therapeutic and commonly abused drugs in serum and urine by GLC using a photoionization detector (186) 637

Thermal conductivity detector
Effect of the sensitivity setting of a katharometer on response factors at low concentrations (197) 121

Thermal degradation
Thermal degradation of sugars in a GC injection port (193) 83

Thermal energy analyzer
Response of nitramines in the thermal analyzer (187) 229

Thermoprogrammed reduction
Thermoprogrammed reduction of cobalt oxide catalysts (188) 254

Thiamine
Modified procedure for determining vitamin B_1 by GC (193) 470

Thiamine phosphates
Differential fluorimetric determination of picogram levels of thiamine, thiamine monophosphate, diphosphate and triphosphate using HPLC (188) 417

Thin-layer chromatograms
Instant copy technique for documentation of thin-layer chromatograms (189) 284

Thin-layer plate preparation
Sintered TLC (Review) (184) 37

Thiocyanate
Determination of cyanide and thiocyanate in biological fluids by GC-MS (188) 357

Thiol peptides
Estimation of glutathione in rat liver by reversed-phase HPLC: separation from cysteine and γ-glutamylcysteine (194) 424

Thiol-proteins
Dissolvable disulphide-polyacrylamide gels for the electrophoretic analysis of chromosomal proteins and for affino-electrophoresis of thiol-proteins (190) 437

Thiols
Solid-phase reagents for LC detection (185) 497

Thiols
Detection of thiols on thin-layer chromatograms with 3,5-di-*tert.* -butyl-1,2-benzoquinone-iron(III) chloride (187) 418

Thiols
Derivatization of dithiols and certain monothiols with phenylarsine oxide for GC (200) 171

Thiols, hydrophilic
Counter-current distribution of hydrophilic thiols in the presence of tributyltin hydroxide (190) 156

Thiolsulphonates
GC and GC-MS characterisation of methane thiolsulphonates carrying further functional groups (198) 287

Thiourea
Inclusion compounds in chromatography (Review) (184) 347

Thymidine dimers
Separation of cyclobutyl dimers of thymine and thymidine by HPLC and TLC (195) 139

Thymine
Analysis of pyrimidine bases in biological materials by GC-MS (190) 107

Thymine dimers
Separation of cyclobutyl dimers of thymine and thymidine by HPLC and TLC (195) 139

Thyroidal amino acids
Quantitative elution of thyroidal amino acids from a cation-exchange column: analysis of tablet formulations containing dyes (196) 512

Thyroid lysosome
Separation of rat thyroid lysosome subpopulations on Sepharose 2B (189) 101

Thyroid proteins
Application of 1,1'-carbonyldiimidazole-activated agarose for the purification of proteins. II. The use of an activated matrix devoid of additional charged groups for the purification of thyroid proteins (185) 463

Thyronines
Separation of enantiomeric iodinated thyronines by LC of diastereomers (198) 471

Tin tetraalkyls
Comparative assessment of GLC and HPLC

for the separation of tin tetraalkyls and alkyltin halides (200) 305

Titration curves
Titration curves of proteins by combined isoelectric focusing-electrophoresis in highly porous polyacrylamide matrices (189) 317

Tocainide
HPLC assay for tocainide in human plasma: comparison with GLC assay (183) 109

Tocopherol
The estimation of α-tocopherol in biological material by GC (183) 487

Tocopherol
Preparation, separation and characterization of vitamin E quinone (196) 506

Tocopherols
Quantitation of serum tocopherols by HPLC with fluorescence detection (181) 242

Tolbutamide
Electron-capture GC of plasma sulphonylureas after extractive methylation (181) 17

Toluene diisocyanate
Determination of free toluene diisocyanate in polyurethane prepolymers by HPLC (198) 212

Toluene sulphonamide
HPLC determination of impurities in commercial saccharin (187) 249

Topological analysis
Topological analysis of the behaviour of linear alkenes up to tetradecenes in GLC on squalane (194) 121

Toroidal coil planet centrifuge
Toroidal coil planet centrifuge for counter-current chromatography (192) 75

Trace analysis
On-line trace enrichment in HPLC using a pre-column (185) 483

Trace analysis
Versatile all-glass splitless sample-introduction system for trace analysis by capillary GC (186) 167

Trace components
Chromatographic analysis of major trace components in solvents with concentration prior to sample injection (191) 309

Trace organics
Direct aqueous injection GC for the analysis of trace organics in water (186) 785

Tranquilizers
Determination of chlordiazepoxide and its metabolites in human plasma by reversed-phase HPLC (181) 141

Tranquilizers
Determination of some benzodiazepines and metabolites in serum, urine and saliva by HPLC (181) 227

Tranquilizers
GC determination of bromo and fluoro derivatives of benzodiazepine in human body fluids (182) 81

Tranquilizers
Determination of peptido-aminobenzophenone (2-*o*-chlorobenzoyl-4-chloro-N-methyl-N'-glycylglycinanilide) by electron-capture GLC (182) 171

Tranquilizers
LC identification of clorazepate in pharmaceutical products (188) 426

Tranquilizers
Screening procedure for therapeutic benzodiazepines by HPLC of their benzophenones (189) 94

Tranquilizers
TLC of the acid hydrolysis products of nineteen benzodiazepine derivatives (194) 262

Transannular bonds
Chromatography of organometallic and organometalloidal derivatives of amino alcohols. II. GC evidence for transannular bonds in solutes (200) 105

Transition metals
Resolution of neutral complexes of transition metals by stereoselective adsorption on optically active complexes. I. Partial resolution of tris(acetylacetonato)cobalt(III) and tris(acetylacetonato)chromium(III) on cobalt(III) complexes (198) 172

Transport relationships
Mass, charge, momentum and energy conservation in electrophoretic fractionation systems (Review) (184) 21

Tranylcypromine
Determination of tranylcypromine in human plasma and urine using high-resolution GLC with nitrogen-sensitive detection (183) 25

Triacylglycerols
Fractionation of oil triacylglycerols by reversed-phase HPLC (196) 279

Triazine derivatives
Comparison of the HPLC behaviour of *s*-triazine derivatives on various stationary phases (187) 341

Triazines
Separation and behaviour of *s*-triazine derivatives on a NH_2-chemically bonded stationary phase by HPLC (191) 115

Triazolophthalazine
Determination of hydralazine metabolites: 4-hydrazino-phthalazin-1-one and N-acetylhydrazinophthalazin-1-one by GC and *s*-triazolo[3,4-α]phthalazine and phthalazinone by HPLC (183) 167

Tributyl phosphate
Separation and determination of the

degradation products of tributyl phosphate by high-speed analytical isotachophoresis (195) 303

Tributyltin hydroxide
Counter-current distribution of hydrophilic thiols in the presence of tributyltin hydroxide (190) 156

Tricyclic antidepressant amines
Peak tailing and retention behaviour of tricyclic antidepressant amines and related hydrophobic ammonium compounds in reversed-phase ion-pair LC on alkyl-bonded phases (189) 299

Tridecenes
Use of Carbowax 20M glass capillary columns below their lower temperature limit (190) 283

Triethylene glycol
GC study of solution and adsorption of hydrocarbons on glycols. I. Diethylene glycol and triethylene glycol (197) 135

Triglycerides
Applications of reversed-phase chromatography and nephelometric detection to analysis of non-polar mixtures at microgram level (187) 111

Triglycerides
Investigation of the use of argentation HPLC for the analysis of triglycerides (188) 205

Triglycerides
Fatty acids. XIX. A quantitative treatment of saturated triglycerides by reversed-phase HPLC (192) 457

Triiodothyronine
Separation of enantiomeric iodinated thyronines by LC of diastereomers (198) 471

Trinucleoside diphosphates
Rapid purification of radiolabeled trinucleoside diphosphates by reversed-phase HPLC (192) 485

Tris(acetylacetonato)chromium
Resolution of neutral complexes of transition metals by stereoselective adsorption on optically active complexes. I. Partial resolution of tris(acetylacetonato)cobalt(III) and tris(acetylacetonato)chromium(III) on cobalt(III) complexes (198) 172

Tris(acetylacetonato)cobalt
Resolution of neutral complexes of transition metals by stereoselective adsorption on optically active complexes. I. Partial resolution of tris(acetylacetonato)cobalt(III) and tris(acetylacetonato)chromium(III) on cobalt(III) complexes (198) 172

Tris(dibromopropyl)phosphate
Quantitative GC determination of tris(2,3-dibromopropyl)phosphate in the 10-ng range by using a 0.8-mm I.D. column packed with a high liquid loaded support (196) 133

Trisoxalatochromium
Paper electrophoretic study of ion-pair formation. XIII. Behaviour of anionic trisoxalatocobalt(III) and trisoxalatochromium(III) in optically active electrolytes (198) 76

Trisoxalatocobalt
Paper electrophoretic study of ion-pair formation. XIII. Behaviour of anionic trisoxalatocobalt(III) and trisoxalatochromium(III) in optically active electrolytes (198) 76

Triterpenoids
TLC of lantadene A and some related triterpenoids (196) 515

Trithimotide
Inclusion compounds in chromatography (Review) (184) 347

Trypan blue
Trypan blue staining of proteins fractionated by acrylamide disc gel electrophoresis (189) 410

Trypsin
Interaction of trypsin with immobilized *p*-aminobenzamidine derivatives studied by means of affinity electrophoresis (194) 175

Trypsin
Affinity chromatographic interactions of proteases with low-molecular-weight soybean protease inhibitors (195) 385

Tryptamine
Trimethylsilylation of biogenic indoleamines. Practical kinetic and analytical aspects related to tryptamine and 5-hydroxytryptamine (186) 619

Tryptamine
In vivo studies of the tryptophan-5-hydroxylase system. Quantitation of serotonin and tryptamine using GC-mass fragmentography (199) 171

Tryptophan
Method for the determination of tryptophan in serum and cerebrospinal fluid (181) 100

Tryptophan
Determination of tryptophan and metabolites in rat brain and pineal tissue by reversed-phase HPLC with electrochemical detection (181) 187

Tryptophan
TLC method for the quantitative analysis of L-tryptophan in human plasma (182) 110

Tryptophan
GLC of free amino acids in the hyaloplasm of rat cerebral, cerebellar and ocular tissues, and in skeletal and heart muscle (182) 147

Tryptophan
Protein-protein interactions studied by counter-current distribution. II. Test of the

theoretical model using bovine serum albumin and L-tryptophan (196) 207

Tryptophanhydroxylase
In vivo studies of the tryptophan-5-hydroxylase system. Quantitation of serotonin and tryptamine using GC-mass fragmentography (199) 171

Tryptophan metabolites
Determination of urinary quinolinic acid by HPLC (182) 425

TSK-GEL
Comparison of TSK-GEL PW type and SW type in high-speed aqueous gel-permeation chromatography (193) 311

TSK-GEL
High-speed gel filtration of proteins in 6 *M* guanidine hydrochloride on TSK-GEL SW columns (193) 458

TSK-GEL SW
Separation range and separation efficiency in high-speed gel filtration on TSK-GEL SW columns (190) 297

Tube deformation
Mass transfer in ideal and geometrically deformed open tubes. III. Deformed metal and plastic tubes (199) 3

Tuberculostearic acid
Detection of tuberculostearic acid in biological specimens by means of glass capillary GC-electron and chemical ionization MS, utilizing selected ion monitoring (182) 402

Tubes with elliptical cross-section
The dynamic diffusion constant within fluid flow in an open straight tube with an elliptical cross-section (196) 349

Tunicamycin homologues
Separation of tunicamycin homologues by reversed-phase HPLC (198) 506

Two-dimensional chromatography
Two-dimensional (Letter to the Editor) (187) 466

Tyrosine
Rapid automated ion-exchange analysis of plasma tyrosine and phenylalanine with data print-out (181) 337

Tyrosine
Presence of sodium azide during acid hydrolysis of protein samples causes the destruction of tyrosine, phenylalanine and histidine (189) 106

Tyrosine monooxygenase
Rapid and sensitive assay of tyrosine 3-monooxygenase activity by HPLC using the native fluorescence of DOPA (198) 511

Tyrosines
Simple and fast separation of the iodotyrosines by TLC (194) 270

Tyrosinyl peptides
Hydrophobic chromatography with dynamically coated stationary phases. II. Dynamic cation-exchange separations of tyrosinyl peptides (194) 353

Tyrosyltyrosine
Separation of iodinated compounds of L-tyrosyl-L-tyrosine from iodothyronines by Biogel P-2 CC (188) 431

U

Ultra-micro high-performance liquid chromatography
Ultra-micro HPLC (190) 150

Uracil
Analysis of pyrimidine bases in biological materials by GC-MS (190) 107

Uracil derivatives
Separation of halogenated uracil derivatives from nucleobases and nucleosides by TLC on silica gel (187) 453

Uracils
GC determination of 5-alkyl-uracils and 5-alkyl-deoxyuridines using flash methylation and glass capillary columns (191) 253

Urea
Inclusion compounds in chromatography (Review) (184) 347

Urea
Electron donor-acceptor properties of urea and its role in charge-transfer chromatography (188) 257

Urea
Separation of methylol derivatives of imidazolidines, urea and carbamates by LC (190) 226

Urea
TLC method for the separation and determination of the products of the reaction of amides with formaldehyde (195) 158

Urea-formaldehyde addition products
Separation of urea-formaldehyde addition products by gel permeation chromatography (195) 154

Ureas
Separation of isomeric ureas and oxazoles by TLC (193) 338

Urease
Purification of enzymes by high-speed gel filtration on TSK-GEL SW columns (190) 305

Urethane prepolymers
Determination of free toluene diisocyanate in polyurethane prepolymers by HPLC (198) 212

Uric acid
Determination of uric acid in serum using isotachophoresis (182) 317

Uricosuric agents
HPLC analysis of sulfinpyrazone and its metabolites in biological fluids (181) 417

Uridine derivatives
Study of rearrangement reactions occurring during GC of *tert.* -butyldimethylsilyl ether derivatives of uridine (194) 379

Uridine diphosphate glucose pyrophosphorylase
Polyacrylamide-boronate beads saturated with biomolecules: a new general support for affinity chromatography of enzymes (189) 225

Uridine diphosphate glucuronic acid
New method for the quantitative assay of the enzymatic hydrolysis of UDPglucuronic acid using analytical capillary isotachophoresis (188) 235

Uridylic acid
Application of a weakly basic dimethylamino-modified silica ion exchanger to the separation of oligonucleotides (185) 403

Urinary constituents
High-performance anion-exchange chromatography of UV-absorbing constituents of human urine (181) 311

Urinary constituents
Comparison of urinary excretion of UV-absorbing constituents in healthy subjects and patients with rheumatoid arthritis using analytical isotachophoresis (181) 347

Urocanic acid
Analysis of *Z* and *E* isomers of urocanic acid by HPLC (183) 83

Urokinase
Rapid method for the determination of urokinase activity by HPLC (188) 185

Urokinase
Assay method for urokinase activity by capillary-tube isotachophoresis using a synthetic substrate (188) 383

Uroporphyrin
Improved separation of uroporphyrin isomers by HPLC (190) 221

Urushiol congeners
HPLC separation of urushiol congeners in poison ivy and poison oak (200) 163

Urushiol diacetate
Separation of Japanese lac urushiol diacetate on silver nitrate-coated silica gel columns by HPLC (198) 49

UV-absorbing compounds
Identification and quantitation of nucleosides, bases and other UV-absorbing compounds in serum, using reversed-phase HPLC. I. Chromatographic methodology (186) 647

UV-absorbing compounds
Identification and quantitation of nucleosides, bases and other UV-absorbing compounds in serum, using reversed-phase HPLC. II. Evaluation of human sera (186) 659

V

Vacuum equipment
Optimization of the instrumental parameters of a combined liquid chromatograph-mass spectrometer coupled by an interface for direct liquid introduction. I. Performance of the vacuum equipment (185) 529

Valine
GLC of free amino acids in the hyaloplasm of rat cerebral, cerebellar and ocular tissues, and in skeletal and heart muscle (182) 147

Valproate
Simple, rapid procedure for the determination of valproate and ethosuximide in plasma by GLC (182) 237

Valproate
Direct determination of valproate in serum by isotachophoresis. Comparison with a GC method (182) 496

Valproic acid
GC determination of valproic acid in human plasma (182) 241

Vanillylmandelic acid
Determination of homovanillic, isohomovanillic and vanillylmandelic acids in human urine by means of glass capillary GLC with temperature-programmed electron-capture detection (183) 391

Vaporizing injectors
Should the septum part of vaporizing injectors be kept at lower temperatures? (193) 64

Vasodilator drugs
Determination of oxpentifylline and a metabolite, 1-(5'-hydroxyhexyl)-3,7-dimethylxanthine, by GLC using a nitrogen-selective detector (181) 355

Vasodilator drugs
HPLC analysis of verapamil. II. Simultaneous quantitation of verapamil and its active metabolite, norverapamil (181) 484

Vasodilator drugs
Micro estimation of plasma theophylline by GLC with on-column butylation and nitrogen-specific detection (182) 232

Vasodilator drugs
Determination of the antispasmodic agent ethaverine in human plasma by HPLC (182) 379

Vasodilator drugs
GLC determination of acephylline in urine (182) 445

Vasodilator drugs
Determination of papaverine in plasma and urine by HPLC (182) 482

Vasodilator drugs
HPLC analysis of dipyridamole in plasma and whole blood (183) 57

Vasopressin analogues
Separation of vasopressin analogues by reversed-phase HPLC (193) 427

Vegetable oils
HPLC determination of *tert.* -butylhydroquinone in vegetable oils (187) 436

Venoms
Application of high-speed aqueous gel permeation chromatography to insect venoms (197) 278

Venzar
GC determination of Lenacil and Pyrazon in soil and bacterial cultures (189) 89

Verapamil
HPLC analysis of verapamil. II. Simultaneous quantitation of verapamil and its active metabolite, norverapamil (181) 484

Verapamil
Analysis of verapamil in plasma by LC (182) 473

Vicine
Chromogenic reagent for vicine and convicine on thin-layer plates (194) 251

Vinca alkaloids
Investigation of vinca alkaloid-plasma membrane interactions by detergent gel chromatography (191) 287

Vinyl chloride
Determination of residual vinyl chloride in poly(vinyl chloride) resins (198) 337

Vinyl chloride polymers
Determination of oligomers in vinyl chloride polymers by steric exclusion chromatography (193) 235

Vinylidene chloride
GC determination of vinylidene chloride monomer in packaging films and in foods (197) 71

Viruses
Affinity chromatography of cells and cell membranes (Review) (184) 471

Viruses
High-speed steric exclusion chromatography of biopolymers (185) 453

Vitamers
LC separation and quantification of B_6 vitamers at plasma concentration levels (196) 176

Vitamin A
HPLC of 13-*cis*-retinoic acid and of endogenous retinol in human plasma (183) 346

Vitamin A
Preparative TLC of the oxidation products of retinyl acetate (191) 301

Vitamin A
HPLC analysis of *cis-trans* stereoisomeric 3-dehydroretinals in the presence of retinal isomers (192) 395

Vitamin B_2
Rapid method for determination of riboflavin in urine by HPLC (182) 285

Vitamin B_6
HPLC determination of pyridoxine and congeners in biological fluids of man after high-dose therapy (183) 492

Vitamin B_6
Separation of vitamin B_6 components by HPLC (188) 285

Vitamin B_6
LC separation and quantification of B_6 vitamers at plasma concentration levels (196) 176

Vitamin B_1
Modified procedure for determining vitamin B_1 by GC (193) 470

Vitamin B_{12}
TLC of acid-labile cobalamins (194) 273

Vitamin D
HPLC of vitamin D: enhanced UV absorbance by prior conversion to isotachysterol derivatives (197) 271

Vitamin D_2/D_3
Rationalisation of the chromatographic behaviour of vitamin D_2/D_3 and related compounds in adsorption HPLC (190) 141

Vitamin D_3
GC-MS properties of boronate esters of 24*R*,25-dihydroxycholecalciferol (192) 434

Vitamin E
Quantitation of serum tocopherols by HPLC with fluorescence detection (181) 242

Vitamin E
The estimation of α-tocopherol in biological material by GC (183) 487

Vitamin E quinone
Preparation, separation and characterization of vitamin E quinone (196) 506

Vitamin K
HPLC assay of vitamin K in human serum (186) 749

Vitamin K_1
Resolution of phylloquinone (vitamin K_1), phylloquinone 2,3-epoxide, 2-chloro-phylloquinone and their geometric isomers by HPLC (200) 293

Vitamins
Determination of 5-methyltetrahydrofolic acid in plasma and spinal fluid by HPLC, using on-column concentration and electrochemical detection (182) 35

Vitamins
Application of HPLC to the separation of ascorbic acid from isoascorbic acid (196) 163

Vitamins
Comparison of reversed-phase and adsorption modes of HPLC for the assay of fat soluble vitamins in multivitamin tablets (200) 300

VM 26
Analysis of the anticancer drugs VP 16-213 and VM 26 and their metabolites by HPLC (182) 211

Volatility
GC measurement of volatility of herbicide esters (195) 75

Volcanic slag
Use of volcanic slag as a support in GC (191) 323

VP 16-213
Analysis of the anticancer drugs VP 16-213 and VM 26 and their metabolites by HPLC (182) 211

W

Water
Comparison of different methods of quantitative analysis employing GLC, illustrated by the determination of water in organic solvents (198) 7

Water
Determination of water at the parts per million level in buta-1,3-diene and associated solvents by GC (199) 149

Water adsorption
Solute-solvent interactions on the surface of silica gel. III. Multilayer adsorption of water on the surface of silica gel (196) 193

Werner complexes
Inclusion compounds in chromatography (Review) (184) 347

X

Xanthines
Determination of oxpentifylline and a metabolite, 1-(5'-hydroxyhexyl)-3,7-dimethylxanthine, by GLC using a nitrogen-selective detector (181) 355

Xipamide
Fluorimetric determination of xipamide in body fluids with a new fluorescent reagent (in German) (183) 124

Xylene isomers
Properties of liquid crystalline cyanoazoxybenzene alkyl carbonates as stationary phases in GC (200) 65

Xylenol
TLC estimation of 2,4-xylenol in 2,5-xylenol (192) 250

Xylenol
Separation of xylenol isomers on nitrofluorenes and nitro-9-fluorenones by GLC. I. Mixed columns and effect of temperature on retention behaviour (193) 300

Xylenols
Separation behaviour of some isomeric organic compounds on sugars, sugar alcohols and their mixed phases by GLC (197) 251

Xylitol
GLC and HPLC analyses of the acid-catalyzed dehydration reaction of xylitol (187) 131

Z

Zeolites
Inclusion compounds in chromatography (Review) (184) 347

Zeolites
Stabilized Y-type zeolites for GC separation (191) 175

Zimelidine
Reversed-phase chromatography of zimelidine and similar dibasic amines. I. Analysis in biological material (185) 593

Zinc
Quantitative analysis of zinc, copper and nickel diethyldithiocarbamates by GLC (187) 363

Zinc
Binding of zinc and copper to some gel filtration media (188) 405

Zinc
GC of volatile metal dialkyldithiophosphates (193) 53

Zinc protoporphyrin
HPLC determination of protoporphyrin and zinc protoporphyrin in blood (181) 319

Zirconium phosphate micro-crystals
Surface ion exchange and adsorption of some dyes on α-$Zr(HPO_4)_2 \cdot H_2O$ micro-crystals (195) 270

Zomepirac sodium dihydrate
Purity determination of zomepirac sodium dihydrate by HPLC (190) 429

Zymograms
Experimental taxonomy of *Biomphalaria* (gastropoda: planorbidae). I. Methods for experimental taxonomic studies on *Biomphalaria* carried out by horizontal starch gel electrophoresis and staining of twelve enzymes (188) 169

PUBLICATION SCHEDULE FOR 1981

Journal of Chromatography (incorporating *Chromatographic Reviews*) and *Journal of Chromatography, Biomedical Applications*

MONTH	N 1980	D 1980	J	F	M	A	M	J	J	A	S	O	N	D
Journal of Chromatography			203 204 205/1 205/2	206/1 206/2 206/3		The publication schedule for further issues will be published later.								
Chromatographic Reviews														
Biomedical Applications	221/1	221/2	222/1	222/2	222/3									

INFORMATION FOR AUTHORS

(Detailed *Instructions to Authors* were published in Vol. 193, No. 3, pp. 529–532. A free reprint can be obtained by application to the publisher)

Types of Contributions. The following types of papers are published in the *Journal of Chromatography* and the section on *Biomedical Applications:* Regular research papers (Full-length papers), Short communications and Notes. Short communications are preliminary announcements of important new developments and will, whenever possible, be published with maximum speed. Notes are usually descriptions of short investigations and reflect the same quality of research as Full-length papers, but should preferably not exceed four printed pages. For reviews, see page 2 of cover under Submission of Papers.

Submission. Every paper must be accompanied by a letter from the senior author, stating that he is submitting the paper for publication in the *Journal of Chromatography*. Please do not send a letter signed by the director of the institute or the professor unless he is one of the authors.

Manuscripts. Manuscripts should be typed in double spacing on consecutively numbered pages of uniform size. The manuscript should be preceded by a sheet of manuscript paper carrying the title of the paper and the name and full postal address of the person to whom the proofs are to be sent. Authors of papers in French or German are requested to supply an English translation of the title of the paper. As a rule, papers should be divided into sections, headed by a caption (e.g., Summary, Introduction, Experimental, Results, Discussion, etc.). All illustrations, photographs, tables, etc. should be on separate sheets.

Introduction. Every paper must have a concise introduction mentioning what has been done before on the topic described, and stating clearly what is new in the paper now submitted.

Summary. Full-length papers and Review articles should have a summary of 50–100 words which clearly and briefly indicates what is new, different and significant. In the case of French or German articles an additional summary in English, headed by an English translation of the title, should also be provided. (Short communications and Notes are published without a summary.)

Illustrations. The figures should be submitted in a form suitable for reproduction, drawn in Indian ink on drawing or tracing paper. Each illustration should have a legend, all the *legends* being typed (with double spacing) together on a *separate sheet*. If structures are given in the text, the original drawings should be supplied. Coloured illustrations are reproduced at the author's expense, the cost being determined by the number of pages and by the number of colours needed. The written permission of the author and publisher must be obtained for the use of any figure already published. Its source must be indicated in the legend.

References. References should be numbered in the order in which they are cited in the text, and listed in numerical sequence on a separate sheet at the end of the article. Please check a recent issue for the lay-out of the reference list. Abbreviations for the titles of journals should follow the system used by *Chemical Abstracts*. Articles not yet published should be given as "in press", "submitted for publication", "in preparation" or "personal communication".

Proofs. One set of proofs will be sent to the author to be carefully checked for printer's errors. Corrections must be restricted to instances in which the proof is at variance with the manuscript. "Extra corrections" will be inserted at the author's expense.

Reprints. Fifty reprints of Full-length papers, Short communications and Notes will be supplied free of charge. Additional reprints can be ordered by the authors. An order form containing price quotations will be sent to the authors together with the proofs of their article.

News. News releases of new products and developments, and information leaflets of meetings should be addressed to: The Editor of the News Section, Journal of Chromatography/Journal of Chromatography, Biomedical Applications, Elsevier Scientific Publishing Company, P.O. Box 330, 1000 AH Amsterdam, The Netherlands.

Advertisements. Advertisement rates are available from the publisher on request. The Editors of the journal accept no responsibility for the contents of the advertisements.

The merits and scope of

modern thin-layer chromatography

can, of course, be fully exploited only with a complete system of compatible instruments f
sample application, chromatogram development, and quantitative evaluation.

Whether you choose conventional TLC, high-performance T
with linear chromatogram development or circular TLC,

CAMAG has the right system for you.

1. Sample application



CAMAG LINOMAT III

narrow band sample application, 5—500 µl

conventional TLC · linear HPTLC



CAMAG NANOMAT

application with fixed-volume sample capillary, 100 nl, 200 nl or 0.5—5 µl

conventional TLC · linear HPTLC · circular HPTLC



CAMAG Nano-Applicator Micro-Applicator

variable volume sample application with micrometer syringe, 50—230 nl, or 0.5—2.3 µl

conventional TLC · linear HPTLC · circular HPTLC

2. Chromatogram development

CAMAG Twin-Trough Chamber

versatile, economical and reliable

conventional TLC



CAMAG HPTLC Linear Chamber

separation of 32 samples on a single 10 x 10 cm HPTLC plate, or 72 samples on a 20 x 10 cm plate

linear HPTLC



CAMAG U-Chamber Chromatograph acc. to R. E. Kaiser

Most efficient HPTLC method based on circular developing technique; also suitable for HPTLC/HPLC transfer

circular HPTLC

3. Quantitative evaluation



CAMAG TLC/HPTLC Scanner

for conventional and high-performance th
layer chromatograms

Three scanning modes for circular chromatograms

Automatic evaluation of all scanning tracks
a complete chromatogram

conventional TLC · linear HPTLC · circular HP

If you are planning to set
a new TLC laboratory, or
extend your existing T
equipment

CAMAG specialists will hel

CAMAG Sonnenmattstr. 11 · CH-4132 Muttenz (Switzerland) · Tel. 061-613434 · Telex 62
CAMAG has agents in almost all countries

PERIODICAL
QD
271
JG
Index
1980

LIBRARY USE ONLY